Birational rigidity and alpha invariants of Fano varieties

Ivan Cheltsov, Arman Sarikyan, Ziquan Zhuang

arXiv:2304.11333

http://arxiv.org/abs/2304.11333v1

How small can the alpha invariant of a rigid Fano be?

For a terminal Fano variety X with $\operatorname{rk}\operatorname{Cl}(X)=1$, birational super-rigidity means that X has no other Mori-fibre-space model and every birational self-map is regular. Its alpha invariant is

$$\alpha(X)=\sup\big\{c\in\mathbf{Q}:\ (X,cD)\text{ is log canonical for every effective }D\sim_{\mathbf{Q}}-K_X\big\}.$$

The question motivating the paper is whether super-rigidity forces $\alpha(X)\geq\frac{1}{2}$ and, if so, whether this threshold can be improved.

Main theorem (Cheltsov–Sarikyan–Zhuang, Theorem 1.4)**.** *For every $\varepsilon>0$ there is a singular terminal Fano variety X with $\operatorname{rk}\operatorname{Cl}(X)=1$ which is birationally super-rigid and satisfies*

$$\frac{1}{2}\leq\alpha(X)\leq\frac{1}{2}+\varepsilon.$$

The paper's second headline result is equivariant and sharply different. For every $\varepsilon>0$, Theorem 1.8 constructs a *smooth* Fano variety X and a finite subgroup $G\subset\operatorname{Aut}(X)$ such that $\operatorname{rk}\operatorname{Pic}^G(X)=1$, X is G-birationally super-rigid, and $\alpha_G(X)<\varepsilon$. Here α_G tests log canonicity only against G-invariant effective anticanonical $\mathbf{Q}$-divisors. The detailed proof roadmap below is deliberately scoped to Theorem 1.4; Theorem 1.8 uses the different high-dimensional construction developed in Sections 4–5.

For an integer $a\geq 2$, take the general quasi-smooth well-formed hypersurface

$$X=\{y^2x_1+f_{2a+1}(x_1,\dots,x_{a+2})=0\}\subset\mathbf{P}(1^{a+2},a).$$

It has dimension $a+1$, $-K_X\cong\mathcal{O}_X(1)$, class-group rank one, and a unique terminal quotient singularity O_y of type $\frac{1}{a}(1,\dots,1)$. The alpha estimate used in the paper gives

$$\alpha(X)\leq\frac{a+1}{2a+1}=\frac{1}{2}+\frac{1}{4a+2};$$

choosing a large yields the desired upper bound. Section 2 proves the new geometric input: every such X is birationally super-rigid.

Ingredient 1: Noether–Fano produces a forbidden centre

If a nontrivial birational map from X to a Mori fibre space existed, the Noether–Fano inequality would give a mobile system $\mathcal{M}$ and $\lambda>0$ with $K_X+\lambda\mathcal{M}\sim_{\mathbf{Q}}0$ such that $(X,\lambda\mathcal{M})$ is not canonical. Let Z be a non-canonical centre. The proof excludes every possible Z.

Blow up O_y with weights $(1,\dots,1)$, with exceptional divisor $E\cong\mathbf{P}^a$, and write

$$K_{\widetilde{X}}\sim_{\mathbf{Q}}f^*K_X+\frac{1}{a}E,\qquad\lambda\widetilde{\mathcal{M}}\sim_{\mathbf{Q}}f^*(\lambda\mathcal{M})-\mu E.$$

If O_y were a centre, Lemma 2.3 would force $\mu>1/a$. But projection from O_y resolves to a morphism g whose contracted curves cover the transform of a divisor, and $g^*\mathcal{O}(1)\sim_{\mathbf{Q}}f^*(-K_X)-E/a$. For $\mu>1/a$, a general member of the mobile transform would have negative intersection with all those curves. Lemma 2.5 therefore gives $\mu\leq 1/a$, excluding the singular point.

Ingredient 2: the centre has exactly codimension two

The centre now lies in the smooth locus. If $\dim Z<a-1$, the $4/\lambda^2$ multiplicity inequality for two general members of $\mathcal{M}$ contradicts their total intersection unless a general point of Z lies on the divisor contracted by projection. In that remaining case, cut to a surface S through the contracted curve L. Writing $\lambda\mathcal{M}|_S=mL+\lambda\Delta$, one has

$$H\cdot L=\frac{1}{a},\qquad L^2=-2+\frac{1}{a}.$$

Inversion of adjunction and Corti's inequality then reduce the alleged non-log-canonicity to

$$0>\frac{(2a-1)(m-1)^2}{a},$$

an impossibility. Hence $\dim Z=a-1$, and a second multiplicity estimate shows that its anticanonical degree is 1 or 2.

Ingredient 3: a threefold slice finishes the exclusion

Intersect with general anticanonical divisors to obtain a threefold

$$V \subset \mathbf{P}(1^4, a), \qquad y^2 x_1 + f_{2a+1}(x_1, \dots, x_4) = 0,$$

and a curve $C = V \cap Z$ of degree $C \cdot H \in \{1, 2\}$. Put $\nu = \lambda \operatorname{mult}_C(\mathcal{M}|_V) > 1$ and blow up C. The transformed mobile system has numerical class $\beta^* H - \nu E$.

For $C \cdot H = 1$, the paper computes $E^3 = a - 1$ and proves $a\beta^* H - E$ nef. Positivity against the square of a mobile class would give

$$0 \leq (a\beta^* H - E)(\beta^* H - \nu E)^2 = (\nu - 1)^2 - 2a(\nu^2 - 1) < 0,$$

a contradiction. When $C \cdot H = 2$, C has one of two explicit normal forms. Their exceptional cubes are respectively $2a - 4$ and -2; the nef classes $a\beta^* H - E$ and $2a\beta^* H - E$ yield the same impossible "nonnegative $<$ zero" intersection. Thus no Noether–Fano centre exists, proving super-rigidity.

Toy principle and the hardest point

The recurring toy principle is that a mobile divisor class must intersect a suitable nef class nonnegatively. Each geometric reduction—weighted blow-up, surface slice, then threefold slice—converts an elusive maximal singularity into a numerical expression that is strictly negative.

The hard point is making this reduction exhaustive. One must first exclude the quotient singularity with the precise discrepancy $1/a$, force a smooth centre into codimension two, classify its degree-1 or degree-2 threefold section, and choose a nef divisor adapted to every normal form. The theorem shows that the one-half scale is asymptotically sharp within this singular family; it does not construct an example with $\alpha(X) < 1/2$. Theorem 1.8 instead shows that the equivariant invariant α_G can be arbitrarily small even under smoothness and G-birational super-rigidity.