

Canonical heights on hyper-Kähler varieties and the Kawaguchi-Silverman conjecture

John Lesieutre, Matthew Satriano

arXiv:1802.07388

<http://arxiv.org/abs/1802.07388v1>

The arithmetic growth problem

Let X be a smooth projective variety over $\overline{\mathbf{Q}}$ and $f : X \rightarrow X$ a surjective endomorphism. The first dynamical degree $\lambda_1(f)$ measures the exponential growth of $(f^n)^*H$, whereas the arithmetic degree of $P \in X(\overline{\mathbf{Q}})$ is

$$\alpha_f(P) = \lim_{n \rightarrow \infty} h_H^+(f^n P)^{1/n}$$

when the limit exists, where $h_H^+ = \max\{h_H, 1\}$. The Kawaguchi–Silverman conjecture predicts $\alpha_f(P) = \lambda_1(f)$ whenever the forward orbit of P is Zariski dense.

Main theorem (Lesieutre–Satriano, Theorem 1.2). *The Kawaguchi–Silverman conjecture holds for every surjective endomorphism of a projective hyper-Kähler variety.*

Here hyper-Kähler means that the complex analytification is simply connected and $H^0(X, \Omega_X^2)$ is spanned by a symplectic form. The statement concerns regular maps; it is not asserted for arbitrary rational self-maps.

Ingredient I: two nef expanding directions

A surjective endomorphism of a hyper-Kähler variety is an automorphism: after base change to $\mathbf{C}$ it is finite étale, and $\chi(\mathcal{O}_X) = 1 + \frac{1}{2} \dim X \neq 0$ forces its degree to be one (Lemma 3.3). The case $\lambda_1(f) = 1$ is already known, so put $\lambda = \lambda_1(f) > 1$.

Perron–Frobenius on the nef cone gives classes ν_+ and ν_- with

$$f^*\nu_+ = \lambda\nu_+, \quad (f^{-1})^*\nu_- = \lambda_1(f^{-1})\nu_-. \quad (1)$$

For a hyper-Kähler automorphism, Oguiso’s equality $\lambda_{\dim X-1}(f) = \lambda_1(f)$, together with dynamical-degree duality, gives $\lambda_1(f^{-1}) = \lambda$. Write $\dim X = 2m$. The Beauville–Bogomolov–Fujiki form q is preserved by f^* , has signature $(1, \rho(X) - 1)$ on $N^1(X)_{\mathbf{R}}$, and satisfies $D^{2m} = c_X q(D)^m$. Thus $q(\nu_+) = q(\nu_-) = 0$; the two rays cannot span a totally isotropic plane, so $\nu = \nu_+ + \nu_-$ has positive top self-intersection. Hence ν is big as well as nef (Lemma 3.8). Simple connectedness gives $h^1(X, \mathcal{O}_X) = 0$, allowing unique lifts to real divisors $D_{\pm}$ with the corresponding linear-equivalence eigenrelations.

Ingredient II: a two-sided canonical height

Tate’s limiting argument produces

$$\widehat{h}_+(P) = \lim_{n \rightarrow \infty} \lambda^{-n} h_{D_+}(f^n P), \quad \widehat{h}_-(P) = \lim_{n \rightarrow \infty} \lambda^{-n} h_{D_-}(f^{-n} P). \quad (2)$$

They transform under f by λ and λ^{-1} , respectively, and $\widehat{h} = \widehat{h}_+ + \widehat{h}_-$ is a height for the big and nef divisor $D = D_+ + D_-$. Its augmented base locus $\mathbf{B}_+(D)$ is proper and f -invariant: changing the positive coefficients of the two nef eigenclasses does not change this locus (Lemmas 3.14–3.16). Explicitly,

$$f(\mathbf{B}_+(\nu_+ + \nu_-)) = \mathbf{B}_+(\lambda^{-1}\nu_+ + \lambda\nu_-) = \mathbf{B}_+(\nu_+ + \nu_-). \quad (3)$$

This isolates a fixed proper algebraic locus on which a big height need not be positive and outside which a dense orbit must remain.

Ingredient III: positivity and Northcott off the bad locus

The central height theorem (Theorem 3.20) is valid more generally for an automorphism satisfying the three conditions just verified: equal forward and inverse dynamical degrees, $D_+ + D_-$ big, and $h^1(\mathcal{O}_X) = 0$. Outside $\mathbf{B}_+(D)$ one has

$$\widehat{h}(P) \geq 0, \quad \widehat{h}_\pm(P) \geq 0, \quad \widehat{h}(f^n P) + \widehat{h}(f^{-n} P) = (\lambda^n + \lambda^{-n})\widehat{h}(P). \quad (4)$$

The first assertion follows by combining a uniform lower bound for a big height with (4) and letting $n \rightarrow \infty$; applying the same idea separately to the two summands gives their non-negativity. A big height has the Northcott property away from its augmented base locus: choose a small ample A such that $D - A$ is rational and $\mathbf{B}_+(D) = \mathbf{B}(D - A)$. Then h_{D-A} is bounded below off this locus, so bounded $h_D = h_A + h_{D-A} + O(1)$ bounds the ample height h_A . Consequently, for P outside that locus,

$$\widehat{h}_+(P) = 0 \iff \widehat{h}_-(P) = 0 \iff \widehat{h}(P) = 0 \iff P \text{ is } f\text{-periodic}. \quad (5)$$

Indeed, (4) makes the whole two-sided orbit have bounded height when the sum vanishes; Northcott makes that orbit finite. If only one summand vanishes, the other decays along the appropriate half-orbit and gives the same conclusion.

Proof roadmap and the decisive growth estimate

A dense orbit cannot meet the proper invariant set $\mathbf{B}_+(D)$. It is not periodic, so (5) makes both $\widehat{h}_+(P)$ and $\widehat{h}_-(P)$ positive. Along the forward orbit,

$$h_D(f^n P) = \lambda^n \widehat{h}_+(P) + \lambda^{-n} \widehat{h}_-(P) + O(1),$$

and therefore $\alpha_f(P) \geq \lambda$. The general upper bound $\alpha_f(P) \leq \lambda_1(f)$ supplies the reverse inequality. This proves the theorem.

A useful toy model is the scalar sequence $a\lambda^n + b\lambda^{-n}$ with $a, b > 0$: its exponential rate is λ , while looking simultaneously forward and backward prevents cancellation and detects precisely when both coefficients vanish. The geometric difficulty suppressed by this model is proving that the sum of the two eigenclasses is big and obtaining Northcott only after removing $\mathbf{B}_+(D)$.

Geometric byproduct and scope

For an f -periodic positive-dimensional subvariety V , define $\lambda_1(f|_V)$ using the induced map on its normalization and set

$$E(f) = \bigcup \{V : \lambda_1(f|_V) < \lambda_1(f), \lambda_1(f^{-1}|_V) < \lambda_1(f^{-1})\}. \quad (6)$$

Spectral-radius comparison gives $\nu_+|_V = \nu_-|_V = 0$ for every such V (Lemma 3.23), hence $E(f) \subset \mathbf{B}_+(\nu)$ and $E(f)$ is not dense (Theorem 3.25).

The contraction statement in Theorem 1.3 requires more than this containment. Choose an effective real divisor $\Delta \equiv \varepsilon\nu$ such that (X, Δ) is klt and take its log canonical model. Because $K_X \equiv 0$ and ν is big and nef, the model map is a morphism $\pi : X \rightarrow Y$ with $\varepsilon\nu \equiv \pi^*A$ for an ample A ; it contracts every component of $E(f)$. The positive-coefficient expressions for ν and $(f^{-1})^*\nu$ show that π and $\pi \circ f$ contract the same subvarieties. The rigidity lemma then descends f to Y , and the same argument for f^{-1} gives an inverse. Thus every connected component of $E(f)$ is contracted to a point (Proposition 3.29).

The paper does not claim $E(f) = \mathbf{B}_+(\nu)$; canonical-height positivity is asserted specifically on $X(\overline{\mathbf{Q}}) \setminus \mathbf{B}_+(D)$. This digest focuses on the title hyper-Kähler results and omits the later Calabi–Yau threefold, Mori-fiber-space, and rational-scroll applications.