On lower bounds for canonical heights of the map phi(X,Y)=(Y,X+Y^D+b)

Jorge Mello

Jorge Mello, arXiv:2607.12668v3 (2026), 10 pages
https://arxiv.org/abs/2607.12668v3

A height gap for a Hénon automorphism

Let K be a number field or a function field of characteristic zero and

$$\phi(x, y) = (y, x + y^d + b), \qquad d \geq 2.$$

Writing h for the usual height on $\mathbf{A}^2(K)$, the forward and backward canonical heights are

$$\widehat{h}_\phi^+(P) = \lim_{n\to\infty} d^{-n} h(\phi^n P), \qquad \widehat{h}_\phi^-(P) = \lim_{n\to\infty} d^{-n} h(\phi^{-n} P),$$

and $\widehat{h}_\phi = \widehat{h}_\phi^+ + \widehat{h}_\phi^-$. This height vanishes precisely on periodic points.

Main theorem (Mello, Theorem 1.1)**.** *There is an integer $B > 0$ such that, unless $P \in \mathbf{A}^2(K)$ is ϕ-periodic of period at most B,*

$$\widehat{h}_\phi(P) \geq \varepsilon \max\{h(b), 1\}.$$

Here $\varepsilon > 0$ depends on d and on the places dividing b.

Thus the parameter height cannot be large while a non-exceptional K-rational orbit has very small canonical height. The dependence on the bad places is essential to the stated result: this is not a uniform lower bound depending only on d.

Ingredient 1: local escape and root clustering

For a place v, the paper introduces forward and backward escape regions

$$\mathcal{B}_v^+ = \{|y|_v^d > (3)_v \max(|x|_v, |b|_v, 1)\}, \quad \mathcal{B}_v^- = \{|x|_v^d > (3)_v \max(|y|_v, |b|_v, 1)\}.$$

They are invariant in the appropriate time direction. On them the local canonical height is the logarithm of the escaping coordinate, exactly at non-archimedean places and up to a bounded archimedean error (Lemma 2.1).

Outside both regions, $|x|_v$ and $|y|_v$ are at most a constant times $\max(1, |b|_v)^{1/d}$. If the neighboring forward and backward iterates also fail to escape, Lemma 2.2 gives roots $\gamma_1^d = \gamma_2^d = -b$ such that y and x lie close to γ_1 and γ_2. This is the local rigidity behind the proof: a long slowly escaping orbit must keep returning to one of only finitely many root clusters.

Ingredient 2: thin one orbit simultaneously at every place

Put $P_j = (x_j, y_j) = \phi^j(P)$ and consider indices in $[-M, M]$. Lemma 2.3 repeatedly applies a pigeonhole argument. At a bad place one keeps a definite fraction of the indices for which either the same escape estimate applies or the same choices of roots of $-b$ occur. For distinct surviving i, j, the two coordinate differences and the local size of b are then bounded by

$$3d^M \widehat{\lambda}_{v,\phi}(P) + \alpha_v,$$

in the precise logarithmic form of Lemma 2.3. The constants α_v vanish globally except for a controlled contribution depending on d. At good non-archimedean places, Lemma 2.6 gives the needed bounds directly.

If s is the number of places dividing b, the recursively defined numbers $B_{m,n}$ in Section 3 choose M large enough that thinning across all s bad places still leaves enough indices. This finite combinatorial step is the useful toy model for the whole proof: many local alternatives are synchronized by discarding indices, not by trying to control every iterate at once.

Take the exceptional period bound $B \geq 2M$. If P is not periodic of period at most B, then the iterates P_i with $i \in [-M, M]$ are distinct: $P_i = P_j$ would give a period $|i - j| \leq 2M$. Consequently, the coordinate differences to which the product formula is applied are genuinely nonzero.

Ingredient 3: the product formula forces global height

If two retained iterates differ in both coordinates, summing the local inequality with the standard place weights and using the product formula gives

$$h(b) \le d^{M+2}\widehat{h}_\phi(P) + C(d).$$

There are two degenerate alternatives: all retained x-coordinates agree, or all retained y-coordinates agree. Lemmas 2.4 and 2.5 supply one-coordinate versions of the same thinning estimate; distinctness of the retained iterates makes the other coordinate differences nonzero, so the identical global conclusion follows. This inequality gives the proportional lower bound when $h(b)$ is large. In the bounded-$h(b)$ regime, discreteness of the canonical height supplies a positive minimum. Together these give ε; the bounded-period exception entered earlier through $B \ge 2M$.

The hardest point is Lemma 2.3. Near roots of $-b$ one must estimate $y_i^d - y_j^d$ (and similarly for x) uniformly at archimedean places and at primes dividing d; those estimates are what make the later product-formula sum quantitative.

Function-field consequence and scope

Let $F = K(C)$ for a curve C over a number field, let η be the pole divisor of $b \in F$, and write the two-sided orbit as

$$\mathcal{O}_\phi(R) = \{\phi^n(R) : n \in \mathbf{Z}\}.$$

If $P, Q \in \mathbf{A}^2(F)$ have distinct two-sided ϕ-orbits, Corollary 1.2 states that, for every $s \ge 1$, only finitely many s-integral $t \in C(K)$ with respect to η satisfy

$$\mathcal{O}_{\phi_t}(P(t)) = \mathcal{O}_{\phi_t}(Q(t)).$$

The paper obtains this by inserting Theorem 1.1 into Ingram's specialization argument; it does not reprove that argument in detail. Likewise, the constants B and ε are existential rather than effective in the note's final statement.