Periods of rational maps modulo primes

Robert L. Benedetto, Dragos Ghioca, Benjamin Hutz, Pär Kurlberg, Thomas Scanlon, Thomas J. Tucker

arXiv:1107.2816

http://arxiv.org/abs/1107.2816v1

Problem and setup

For a rational map $\varphi : \mathbf{P}^1_K \to \mathbf{P}^1_K$ of degree at least two over a number field, reduction modulo a prime turns every orbit into a finite directed graph. A global wandering orbit can therefore become periodic or meet a prescribed point after reduction. The paper asks for a positive-density set of primes on which finitely many such orbit–target incidences are eventually excluded, simultaneously.

Main theorem (Theorem 3.1)**.** *For $1 \le i \le g$, let $\varphi_i : \mathbf{P}^1_K \to \mathbf{P}^1_K$ have degree at least two, and let $A_i, T_i \subset \mathbf{P}^1(K)$ be finite. Assume no T_i contains a φ_i-preperiodic point. Among all the A_i, at most one set contains a non-preperiodic point, and that set contains at most one such point. Then there are $M \ge 1$ and a positive-density set $\mathcal{P}$ of primes of K such that, for every $\mathfrak{p} \in \mathcal{P}$, every i, $\gamma \in T_i$, $\alpha \in A_i$, and $m \ge M$,*

$$\varphi_i^m(\gamma) \not\equiv \alpha \pmod{\mathfrak{p}}.$$

In particular, if α is not periodic for one map φ, then $\alpha \bmod \mathfrak{p}$ is nonperiodic for a positive density of good-reduction primes (Corollary 1.2).

The strength is uniformity: one M and one positive-density prime set work for every listed map, source, target, and later iterate. The theorem is also the arithmetic input for the paper's new cases of dynamical Mordell–Lang and for a positive-density failure statement for Newton iteration at finite places.

Proof mechanism

1. Normalize the periodic targets. After a finite extension of K, replace each map by a common iterate and enlarge each T_i by finitely many intermediate iterates. Every preperiodic target in A_i can then be treated as a fixed point. This reduction preserves the desired conclusion for the original maps.

2. Control deep inverse images at one auxiliary prime. Choose a prime $\mathfrak{p}$ of good reduction, with residue characteristic larger than all degrees and with unchanged ramification indices at the fixed targets. If the exceptional wandering target α_0 occurs, a cited one-dimensional result supplies such a $\mathfrak{p}$ for which α_0 is not periodic modulo $\mathfrak{p}$. Lemma 3.2 says that ramification in the field of definition of $\beta \in \varphi^{-m}(\alpha)$ forces β to reduce to a ramification point of φ^m. Because a nonperiodic residue class can be hit at most once along any reduced forward orbit, only finitely many critical backward branches cause trouble; adjoining them removes that trouble.

Lemma 3.3 then gives, at every sufficiently deep inverse level, both unramifiedness and residue degree > 1. Proposition 3.4 extends this to a finite target set: for a fixed target, one applies the lemma to its nontrivial first preimages, using preservation of the target's ramification index to keep those preimages nonperiodic modulo $\mathfrak{p}$.

3. Turn local control into positive density. Adjoin all points whose first entrance into some A_i occurs at time M. The resulting finite Galois extension is unramified at the chosen prime, and its Frobenius fixes none of those first-hit points because their residue degrees exceed one. Chebotarev therefore gives a positive-density set of primes with the same no-fixed-point property.

4. Propagate from one block to the whole tail. Claim 3.5 is the finite-state step: at any Chebotarev prime, a hit at time $m \ge M$ would force a hit during the first M iterates. Otherwise the point at time $m - M$ would give a forbidden first-hit preimage fixed by Frobenius. Removing finitely many primes excludes these early hits for periodic targets; a second finite exclusion handles the sole wandering target. This yields the simultaneous avoidance statement.

Toy model, difficulty, and scope

The combinatorial core can be pictured in a finite functional graph: if no residue-class vertex outside A enters A for the first time after exactly M steps, then every later visit to A has an earlier visit less than M steps away. The proof's hard part is to manufacture a positive-density family of primes with precisely this missing first-entry layer; backward-orbit ramification and Chebotarev provide it.

The restriction to one wandering target is genuine for this method: the auxiliary prime is obtained from a result controlling one nonperiodic point, and the paper explicitly leaves two unrelated wandering points open. Its higher-dimensional discussion is heuristic and numerical, not a theorem: random-map estimates suggest that periodic orbit pieces may meet a ramification hypersurface too often for the same strategy to work.

Codex mathematical paper digest, 2026-10-28. Source paper should be consulted for full statements and proofs.