

Arithmetic dynamics of random polynomials

Pierre Le Boudec, Niki Myrto Mavraki

arXiv:2112.12005

<http://arxiv.org/abs/2112.12005v1>

The family and its statistical invariants

Fix $d \geq 2$. Away from a lower-dimensional exceptional locus, affine conjugacy classes of degree- d polynomials over $\mathbb{Q}$ have a unique representative

$$\psi_{\mathbf{a}}(z) = \frac{a_d}{a_0} z^d + \frac{a_{d-2}}{a_0} z^{d-2} + \cdots + \frac{a_1}{a_0} z + 1, \quad \mathbf{a} = (a_d, a_{d-2}, \dots, a_0) \in \mathbb{Z}_{\text{prim}}^d, \quad (1)$$

with $a_d \neq 0$ and $a_0 > 0$. Write $\mathcal{P}_d$ for this family, $\mathbb{Z}_{\text{prim}}^d$ meaning that the displayed coefficients have greatest common divisor 1, $H(\psi_{\mathbf{a}}) = \max_i |a_i|$, and $\mathcal{P}_d(X) = \{\psi_{\mathbf{a}} : H(\psi_{\mathbf{a}}) \leq X\}$. Let $\text{Prep}_{\mathbb{Q}}(\psi)$ be the rational preperiodic set and $\hat{h}_{\psi}$ the canonical height. Its first positive value is

$$\lambda_1(\psi) = \min\{\hat{h}_{\psi}(z) : z \in \mathbb{Q} \setminus \text{Prep}_{\mathbb{Q}}(\psi)\}. \quad (2)$$

Main theorem (Le Boudec–Mavraki, Theorems 1.1 and 1.2). *For $\varepsilon > 0$, put $\vartheta_2 = 1/2$ and $\vartheta_d = 2(d+1)/(5d+1)$ for $d \geq 3$. Then*

$$\frac{1}{\#\mathcal{P}_d(X)} \sum_{\psi \in \mathcal{P}_d(X)} \#\text{Prep}_{\mathbb{Q}}(\psi) \ll_{d,\varepsilon} X^{-\vartheta_d + \varepsilon}. \quad (3)$$

Moreover,

$$\lim_{X \rightarrow \infty} \frac{1}{\#\mathcal{P}_d(X)} \#\left\{\psi \in \mathcal{P}_d(X) : \left|\lambda_1(\psi) - \frac{\log H(\psi)}{d(d-1)}\right| \leq \varepsilon \log H(\psi)\right\} = 1. \quad (4)$$

Thus 100% of these classes have no rational preperiodic point, and the generic smallest nonzero canonical height has the sharp scale predicted by the dynamical Lang philosophy. The result is an average statement over the normalized coefficient box, not a pointwise uniform-bound theorem.

Local escape becomes an arithmetic sieve

The decisive input is that infinity is a superattracting fixed point of a polynomial. For every place v , the local escape rate

$$\hat{\lambda}_{\psi,v}(z) = \lim_{n \rightarrow \infty} d^{-n} \log \max\{|\psi^n(z)|_v, 1\}$$

has controlled escape estimates outside a coefficient-dependent radius: Lemma 2.1 is an exact formula at non-Archimedean places, while Lemma 2.3 is the required Archimedean lower inequality. The product formula gives $\hat{h}_{\psi} = \sum_p \hat{\lambda}_{\psi,p} + \hat{\lambda}_{\psi,\infty}$. Consequently a rational point of small canonical height imposes divisibility conditions at many primes dividing a_0 .

For $k \geq 2$, let $s_k(a_0)$ be the part of a_0 supported at primes whose exponent in a_0 is less than k . In degree two, Lemma 2.4 gives the clean inequality

$$\hat{h}_{\psi}(z) \geq \frac{1}{2} \log s_2(a_0). \quad (5)$$

Hence a rational preperiodic point forces a_0 to be squareful. Since squareful integers up to X are $O(X^{1/2})$, this already supplies the $d = 2$ power saving after Benedetto's pointwise estimate $\#\text{Prep}_{\mathbb{Q}}(\psi) \ll_d 1 + \log H(\psi)$.

For $d \geq 3$ and $k \in \{2, 3\}$, write $z = x/y$ in lowest terms and let $\sigma_{\mathbf{a}}(x, y)$ be the nonconstant numerator contribution in $a_0 y^d \psi_{\mathbf{a}}(x/y)$. Lemma 2.7 says, schematically but with effective exponents, that small $\hat{h}_{\psi}(z)$ forces a large divisor of $s_k(a_0)$ to divide $\sigma_{\mathbf{a}}(x, y)$, apart from factors shared with a_d . The potentially troublesome shared part is isolated by

$$\Delta_{a_0}(a_d) = \text{the largest divisor of } a_d \text{ supported on primes dividing } a_0.$$

Lemma 2.10 shows that the subfamily $\Delta_{a_0}(a_d) \leq X^{\delta}$ has density one. On this subfamily, Lemma 2.11 counts coefficient vectors together with rational points of bounded canonical height: one first sums the resulting congruence in a coefficient, then uses the $O(Y^{1/k})$ count for k -full integers. This is the hardest part of the argument; it converts local dynamical information into a uniform geometry-of-numbers estimate.

How the two sides of the height asymptotic meet

For (3), take height bound $T = 1$ in Lemma 2.11. Degree two uses (5); for $d \geq 3$, the authors use $k = 3$ and optimize the cutoff δ . The density-zero complementary subfamily and Benedetto's logarithmic bound are then harmless.

The upper side of (4) is unusually concrete. If h denotes logarithmic Weil height, Lemma 2.8 bounds

$$\widehat{h}_\psi(z) - h(z) \leq \frac{\log H(\psi) + \log d}{d-1}.$$

Because $\psi(0) = 1$ and $\widehat{h}_\psi(\psi(z)) = d\widehat{h}_\psi(z)$, whenever 0 is not preperiodic this yields

$$\lambda_1(\psi) \leq \widehat{h}_\psi(0) \leq \frac{\log H(\psi) + \log d}{d(d-1)}. \quad (6)$$

Theorem 1.1 makes the excluded case negligible. For the lower side, any violation supplies at least one rational point below the target height. When $d = 2$, (5) and a separate sum over the squareful part of a_0 show that these exceptions have density zero. Only when $d \geq 3$ do the authors apply Lemma 2.11 with $k = 2$ to obtain the same conclusion.

Higher minima and scope

The paper also defines λ_N using N -tuples satisfying no algebraic relation preserved by diagonal application of ψ . For fixed $N \geq 3$ and $d \geq 3$, Theorem 1.4 shows, with density one,

$$d \left(1 - \frac{4}{d+3} \right) - \varepsilon < \frac{\lambda_2}{\lambda_1} \leq d + \varepsilon, \quad 1 \leq \frac{\lambda_N}{\lambda_2} \leq 1 + \frac{4}{d-1} + \varepsilon. \quad (7)$$

The more general Proposition 4.4 says that, if $d \geq 3$, $N \geq 2$, and 0 is not preperiodic, then

$$\frac{1}{2} \log \frac{N}{H(\psi)} + O(1) < \lambda_N(\psi) \leq (1 + \varepsilon) \log N + \left(\frac{1}{d-1} + \varepsilon \right) \log H(\psi) + O(1). \quad (8)$$

Even N may grow: for $d \geq 3$, $t \geq 0$, $\varepsilon > 0$, and $N_t(\psi) = 1 + \lceil H(\psi)^t \rceil$, Proposition 4.5 proves with density one that $\lambda_{N_t(\psi)}(\psi) > ((1 + 2t)/(d + 3) - \varepsilon) \log H(\psi)$.

For the upper bound in (8), Medvedev–Scanlon’s classification is used only after assuming 0 is not preperiodic; reciprocals of distinct primes coprime to $a_0 a_d$ then form a dynamically independent tuple (Lemmas 4.2–4.3). Theorem 1.1 makes the non-preperiodicity hypothesis generic in the statistical consequences. This classification ingredient is not needed for (3)–(4), whose core is local-height decomposition and the arithmetic sieve above.