

Rigid Gorenstein toric Fano varieties arising from directed graphs

Selvi Kara, Irem Portakal, Akiyoshi Tsuchiya

arXiv:2103.06404

<http://arxiv.org/abs/2103.06404v2>

From a directed graph to a rigid Fano variety

Let G be a finite directed graph on $[n]$. For an edge (i, j) put $\rho(i, j) = e_i - e_j$ and form its directed edge polytope

$$\mathcal{A}_G = \text{conv}\{\rho(i, j) : (i, j) \in A(G)\} \subset \mathbb{R}^n. \quad (1)$$

Assume every directed edge lies on a directed cycle. Proposition 2.1 says that $\mathcal{A}_G$ is terminal and reflexive, so its spanning fan defines a Gorenstein toric Fano variety X_G with terminal singularities. The problem is to recognize from G when Totaro's criterion applies: smoothness in codimension 2 and $\mathbb{Q}$ -factoriality in codimension 3 imply rigidity.

Main theorem (Kara–Portakal–Tsuchiya, Theorem 1.4). *Under the cycle hypothesis above, the following are equivalent.*

1. X_G is smooth in codimension 2 and $\mathbb{Q}$ -factorial in codimension 3.
2. Both graph conditions hold:

(a) G has no directed subgraph C_1 with edge set

$$\{(i_1, i_2), (i_1, i_4), (i_3, i_2), (i_3, i_4)\}; \quad (2)$$

(b) whenever G contains a directed subgraph C_2 with edge set

$$\{(i_1, i_2), (i_2, i_3), (i_1, i_4), (i_4, i_3)\}, \quad (3)$$

either (i_1, i_3) is an edge, or there is $j \notin \{i_2, i_4\}$ with edges (i_1, j) and (j, i_3) .

When these conditions hold, X_G is rigid.

For a symmetric directed graph, Corollary 1.5 collapses the answer to a familiar statement: the underlying undirected graph contains no 4-cycle as a subgraph.

Why only two-dimensional faces matter

Terminality and reflexivity already make X_G smooth in codimension 2. Moreover, $\mathbb{Q}$ -factoriality in codimension 3 is exactly the assertion that every 2-face of $\mathcal{A}_G$ is a triangle. Thus the proof must classify all possible nontriangular 2-faces; Proposition 2.3 then invokes Totaro to obtain rigidity.

The first reduction is elementary but crucial. Lemma 3.1 says every proper face of $\mathcal{A}_G$ is $\mathcal{A}_D$ for an acyclic directed subgraph $D \subset G$. Indeed, if D contained a directed cycle, the average of its root vectors would be 0, forcing the origin into a proper face although it lies in the relative interior of $\mathcal{A}_G$.

For an acyclic D , enlarge to $\tilde{\mathcal{A}}_D = \text{conv}(0, \mathcal{A}_D)$. The imported face criterion, Theorem 3.17, says that $\mathcal{A}_H$ is a face of $\tilde{\mathcal{A}}_D$ exactly when $H \subset D$ is *path consistent* and *admissible*. Path consistency means that the signed net length between two vertices is independent of the undirected path. It gives a vertex weight ω . After contracting each connected component of H^{un} , every component-graph edge e is induced by a unique omitted edge $(u, v) \in A(D) \setminus A(H)$; set $\text{wd}(e) = \omega(u) - \omega(v)$. Admissibility requires every directed cycle C in the component graph to satisfy

$$\sum_{e \in C} \text{wd}(e) > -|C|, \quad (4)$$

This converts the global question “is this quadrilateral a face?” into finite inequalities on paths and weights.

The two square mechanisms

For C_1 , the four root vectors in (2) satisfy the parallelogram relation

$$\rho(i_1, i_2) + \rho(i_3, i_4) = \rho(i_1, i_4) + \rho(i_3, i_2). \quad (5)$$

Lemma 4.1 builds a supporting hyperplane using the cycle-height function μ_{C_1} , with values 1 on the two sources and 0 on the two sinks. The resulting subgraph is path consistent, while every edge of its component graph has weight decrease between -1 and 1 and a cycle contains at most one negative edge. Condition (4) follows, so (5) is genuinely a square 2-face.

For C_2 , (3) consists of two directed length-two paths from i_1 to i_3 . If there is neither a direct shortcut nor another such intermediate vertex, Lemma 4.2 repeats the supporting-face argument with weights 0, 1, 2. A violation of (4) could only be a two-edge component cycle with decreases -2 and 0 , which would produce precisely the excluded extra two-step path. Hence this unfilled diamond also cuts out a square face.

The difficult converse

Suppose instead that $\mathcal{A}_G$ has a nontriangular 2-face. Lemma 4.3 first classifies the acyclic graphs D with $\dim \mathcal{A}_D = 2$. If D has d nonisolated vertices and r connected components, incidence-matrix rank gives

$$2r \leq d \leq r + 3,$$

so $r \leq 3$ and $3 \leq d \leq 6$. Checking these few component types shows that every nontriangle is a square of type C_1 or C_2 .

The C_2 case carries the real global difficulty. If (i_1, i_3) were an edge, the segment from 0 to $\rho(i_1, i_3)$ would meet the interior of the putative square, so it could not be a face. If extra two-step paths through $j_1, \dots, j_k$ existed, the authors enlarge C_2 to the subgraph G' made of all these paths. A supporting hyperplane and the 0, 1, 2 weight function show, via (4), that $\mathcal{A}_{G'}$ is itself a face of $\mathcal{A}_G$. But C_2 is not admissible inside G' , hence cannot be a face of $\mathcal{A}_{G'}$ —contradicting the fact that an intersection of two faces is a face of each. This proves that a square face forces exactly one of the two obstructions in the theorem.

Toy model and proof roadmap

Think of C_1 as the oriented complete bipartite graph with two sources and two sinks. Equation (5) is the affine dependence that makes its four edge vectors a square. The second pattern is a diamond: two parallel directed routes of length two. A shortcut from source to target, or a third route of length two, destroys the exposed square. The full proof is

$$\begin{aligned} \text{cycle condition} &\Rightarrow \mathcal{A}_G \text{ terminal reflexive} \Rightarrow \text{rigidity if all 2-faces are triangles,} \\ &\text{face criterion} \Rightarrow (\text{square faces} \iff C_1 \text{ or unfilled } C_2). \end{aligned} \tag{6}$$

For disconnected G , Proposition 2.2 decomposes $\mathcal{A}_G$ as a free sum and X_G as the product of the varieties of its connected components; the triangular-face property passes through this reduction.

Consequences and limits

The criterion produces rigid but nonsmooth examples. Proposition 5.1 shows that the symmetric directed $2k$ -cycle gives a variety that is $\mathbb{Q}$ -factorial in codimension $2k - 2$ and rigid for $k \geq 3$; Proposition 5.3 gives further rigid families, under its connected and no-multiple-edge hypotheses, by gluing directed even cycles. The theorem is not a converse to rigidity itself. When $\dim \mathcal{A}_G \geq 3$, a square 2-face makes the affine cone's deformation space large, but its degree-zero part need not survive to a projective deformation. Example 5.4 is nonrigid, while Problem 5.5 leaves the general square-face case open. Thus the paper completely classifies when Totaro's sufficient criterion applies to these graph Fano varieties, not all rigid X_G .