Canonical Height Functions For Monomial Maps

Jan-Li Lin, Chi-Hao Wang

arXiv:1205.2020

http://arxiv.org/abs/1205.2020v1

Problem and setup

For an integral matrix $A = (a_{ij})$ with $\det A \neq 0$, the associated monomial map on the torus is

$$\varphi_A(x_1, \ldots, x_N) = \left(\prod_j x_j^{a_{1j}}, \ldots, \prod_j x_j^{a_{Nj}} \right).$$

Write δ_A for the spectral radius of A, and let $\ell_A + 1$ be the largest Jordan-block size among eigenvalues of modulus δ_A. Silverman's forward height is

$$\widehat{h}_A^+(P) = \limsup_{n\to\infty} \frac{h(\varphi_A^n(P))}{n^{\ell_A}\delta_A^n}.$$

It can have infinitely many rational points of arbitrarily small positive height (Proposition 4), so it need not satisfy Northcott finiteness. Set

$$A' = |\det A| A^{-1} \in \mathrm{Mat}_N(\mathbf{Z}), \qquad \widehat{h}_A^- := \widehat{h}_{A'}^+, \qquad \widehat{h}_A := \widehat{h}_A^+ + \widehat{h}_A^-.$$

The adjugate-type matrix A' supplies an integral substitute for inverse iteration.

The total height also has a functional characterization. If $\delta_A > 1$, order the eigenvalues so that $|\lambda_1| \geq \cdots \geq |\lambda_N|$ and put $D = |\det A|$. Theorem 7 proves

$$\frac{\widehat{h}_A(\varphi_A(P))}{|\lambda_1\lambda_N|} + \frac{\widehat{h}_A(\varphi_{A'}(P))}{D} = \left(\frac{1}{|\lambda_1|} + \frac{1}{|\lambda_N|} \right) \widehat{h}_A(P).$$

Moreover, $\widehat{h}_A$ is the unique function satisfying this relation among functions differing from it by $O(1)$. Thus the construction is not merely a sum of two limsups: its interaction with the forward and adjugate maps determines it up to no bounded ambiguity.

The mechanism is the commutation identity $\varphi_A \circ \varphi_{A'} = \varphi_{DI_N} = \varphi_{A'} \circ \varphi_A$. Together with $\delta_A = |\lambda_1|$ and $\delta_{A'} = |\lambda_1 \cdots \lambda_{N-1}| = D/|\lambda_N|$, it gives the required cross-scaling identities for the two forward heights. For uniqueness, the difference g of two candidate functions is bounded and satisfies the homogeneous relation. The strict coefficient inequality

$$\frac{1}{|\lambda_1\lambda_N|} + \frac{1}{D} < \frac{1}{|\lambda_1|} + \frac{1}{|\lambda_N|}$$

forces $\sup|g| = 0$. This is why a bounded error cannot survive the two-map equation.

Main theorem (Theorem 10)**.** *Suppose that A is diagonalizable over $\mathbf{C}$.*

1. If all eigenvalues of A have the same modulus, then $\widehat{h}_A^+(P) \geq h(P)$ for every $P \in \mathbf{G}_m^N(\overline{\mathbf{Q}})$.

2. If the eigenvalues of A have exactly two distinct moduli, then $\widehat{h}_A(P) \geq h(P)$ for every such P.

Consequently the total height has Northcott finiteness: bounded degree and bounded total height define a finite set (Corollary 11). If moreover $\delta_A > 1$, then $\widehat{h}_A \asymp h$ (Corollary 13). In particular, every diagonalizable 2×2 integral matrix with spectral radius greater than one is covered (Corollary 14).

Proof mechanism

1. Extract the extremal spectral projector. Write $A = B\Lambda B^{-1}$. If the eigenvalues of largest modulus $|\lambda|$ occupy the first i_0 diagonal entries, compactness of their phases gives a subsequence n_r for which

$$\frac{A^{n_r}}{|\lambda|^{n_r}} \longrightarrow \alpha := B\,\mathrm{diag}(I_{i_0}, 0)B^{-1}.$$

Evaluating the local-max formula for projective height along this subsequence yields

$$\widehat{h}_A^+(P) \geq \sum_v \max_i \left\{ 0, \sum_j \alpha_{ij} \log |x_j|_v \right\}.$$

If every eigenvalue has modulus $|\lambda|$, then $\alpha = I_N$, and the right-hand side is exactly $h(P)$. This proves the first assertion.

2. Recover the complementary directions by backward growth. Assume now that the two moduli are $|\lambda| > |\mu|$. The eigenvalues of A' are $|\det A|/\lambda_i$, so its dominant eigenspaces are precisely the $|\mu|$-eigenspaces of A. A second subsequence therefore gives the complementary projector

$$\beta = B\,\mathrm{diag}(0, I_{N-i_0})B^{-1}, \qquad \alpha + \beta = I_N,$$

and the same local calculation bounds $\widehat{h}_A^-$ by the corresponding β-expression.

3. Reassemble the ordinary height. At every place, the sum of the two maxima is at least the maximum of the sums. Hence

$$\widehat{h}_A(P) \geq \sum_v \max_i \left\{ 0, \sum_j (\alpha_{ij} + \beta_{ij}) \log |x_j|_v \right\} = h(P).$$

Northcott's theorem for h now gives Corollary 11. The reverse comparison $\widehat{h}_A \ll_A h$ comes from the standard upper bounds for the two forward canonical heights.

The functional equation also makes the height growth along an orbit explicit. Corollary 9 sets $H_n = \widehat{h}_A(\varphi_A^n(P))$ and obtains

$$H_{n+2} - (|\lambda_1| + |\lambda_N|)H_{n+1} + |\lambda_1\lambda_N|H_n = 0.$$

Thus only the largest and smallest eigenvalue moduli govern this second-order recurrence, even when A has many intermediate eigenspaces.

Toy model, difficulty, and scope

For $A = \mathrm{diag}(d, e)$ with integers $d > e > 1$, forward normalization by d^n retains the x-direction and loses the y-direction. Since $A' = \mathrm{diag}(e, d)$, the backward height retains exactly the missing direction. The two spectral projectors add to the identity; this is the linear-algebraic heart of the proof. The real difficulty is that global height is a sum of local maxima, not a linear functional, so the complementary projectors must be combined through the max inequality place by place.

The construction is not universal. Suppose A is non-diagonalizable with its sole eigenvalue $\lambda \in \mathbf{Z}$, let $\ell + 1$ be its largest Jordan-block size, and set $\pi = \varphi_{(A-\lambda I)^\ell}$. Theorem 18 gives, for $\lambda > 0$,

$$\widehat{h}_A^+(P) = \frac{h(\pi(P))}{\ell!\lambda^\ell}, \qquad \widehat{h}_A^-(P) = \frac{h(\pi(P)^{-1})}{\ell!\lambda^\ell};$$

for $\lambda < 0$, each is the corresponding maximum of these two numerators divided by $\ell!|\lambda|^\ell$. Hence the heights factor through a lower-dimensional monomial fibration and are constant on its fibers, so Northcott fails.

In dimension at least three, Proposition 20 produces infinitely many rational points of arbitrarily small positive *total* height when the characteristic polynomial is irreducible and the eigenvalues are distinct and positive. Its approximation argument chooses integral exponent vectors almost annihilated by both extreme spectral covectors; points $(2^{y_1}, \ldots, 2^{y_N})$ then make both the forward and backward contributions small. Thus Theorem 10 is a strong repair in its spectral range, not a final canonical-height theory for all monomial maps.