Preperiodic points for rational functions defined over a rational function field of characteristic zero

J. K. Canci

arXiv:1505.07753

http://arxiv.org/abs/1505.07753v2

Uniformity over an infinite constant field

Let k be algebraically closed of characteristic 0, $K = k(t)$, and S a nonempty finite set of places of K, with $|S| = s$. The paper calls $\phi \in K(z)$ *isotrivial over* K if a conjugate by some $A \in \mathrm{PGL}_2(K)$ is defined over k. Its notion of good reduction at a place likewise allows a K-rational conjugation to a model with simple good reduction. These conjugacy-invariant conventions are part of the statement.

Main theorem (Theorem 1)**.** *For every positive integer d there is a number $B(d, s)$ such that, if $\phi \in K(z)$ has degree d, is not isotrivial over K, and has good reduction outside S, then*

$$\#\,\mathrm{PrePer}(\phi, K) \leq B(d, s).$$

This is a function-field analogue of dynamical uniform boundedness. The main obstacle is that every residue field is the infinite field k: reduction can no longer bound an orbit by counting residue classes. The proof replaces that counting argument by a rigidity statement for S-unit equations. It applies to rational maps, not just polynomials, and even permits $d = 1$ under the non-isotriviality hypothesis.

Ingredient 1: distances turn dynamics into divisibility

Write R_S for the ring of S-integers. Because R_S is a PID, every point of $\mathbb{P}^1(K)$ has S-coprime coordinates. For a place $\mathfrak{p} \notin S$, the logarithmic distance $\delta_{\mathfrak{p}}(P, Q)$ measures congruence of two such points. Simple good reduction gives

$$\delta_{\mathfrak{p}}(\phi(P), \phi(Q)) \geq \delta_{\mathfrak{p}}(P, Q),$$

while on a periodic cycle the relevant distances are invariant under a common shift (Propositions 3.1–3.3). A global minimal model converts the theorem's conjugacy-invariant good reduction into simple good reduction outside S, so these local inequalities may be used simultaneously.

Ingredient 2: the normalized orbit and the S-unit bottleneck

Zannier's theorem bounds by 9^{s-1} the solutions of $\lambda x + \mu y = 1$ in S-units for which $\lambda x/(\mu y)$ is not constant. If some place outside S distinguishes λ and μ, there are no constant-ratio solutions, giving the finite form used repeatedly in the proof (Lemma 4.1).

The basic model is an orbit segment whose pairwise outside-S distances all equal the first distance. Conjugating its first three points gives

$$0 \longmapsto \infty \longmapsto 1 \longmapsto \lambda_3 \longmapsto \cdots .$$

The distance equalities force both λ_i and $1-\lambda_i$ to be S-units. Zannier bounds the nonconstant λ_i's. Constant blocks are controlled by non-isotriviality: Lemma 4.2 proves that there are at most $2d$ constant pairs $(a,b) \in k^2$ with $\phi(a) = b$. Hence each constant block has bounded length, and Lemma 4.3 bounds the whole segment by

$$A(d,s) = \frac{9^{s-1}+1}{2}(2d+1) + 2.$$

This normalized four-point picture is the toy model for every later counting argument: local equal-distance constraints become one additive equation in a multiplicative group.

Ingredient 3: cycles are assembled prime by prime

For a cycle of prime length, Proposition 3.3 puts all its pairwise distances in the setting of Lemma 4.3. For prime-power length p^r, the proof writes the cycle in S-coprime coordinates. If every new coefficient is an S-unit, Lemma 4.3 applies directly; at the first nonunit coefficient, many indices produce distinct solutions of a fixed equation $\lambda x + y = 1$, and Lemma 4.1 bounds their number. Factoring a general period and applying this argument to suitable iterates gives Theorem 2: every minimal period is at most an explicit $C(d,s)$. The large displayed product in the paper matters only through its dependence on d and s.

Ingredient 4: tails satisfy the same equation

After passing to an iterate bounded by $C(d,s)$, the eventual periodic point may be taken to be fixed. Along a backward tail

$$P_{-m+1} \longmapsto \cdots \longmapsto P_{-1} \longmapsto P_0 \longmapsto P_0,$$

good reduction makes the distances to P_0 monotone and supplies the divisibilities in Lemma 4.6. The initial equal-distance portion has length at most $A(d,s)$. At the first strict divisibility, comparing each later point with it again yields one S-unit equation with a nonunit coefficient, so there are at most 9^{s-1} remaining possibilities. This proves the orbit bound $D(d,s)$ of Theorem 3.

Finally all periods divide

$$N(d,s) = \prod_{\substack{p \le C(d,s) \\ p \text{ prime}}} p^{\max_{1 \le n \le C(d,s)} \operatorname{ord}_p(n)},$$

where $\operatorname{ord}_p(n)$ is the exponent of p in n. Thus there are at most $d^{N(d,s)}+1$ periodic points, and at most $d^{D(d,s)}$ points above each; the product is a valid $B(d,s)$.

Scope and hardest point

The difficult step is not merely invoking an S-unit theorem, but separating its uncontrolled constant solutions and proving that non-isotriviality bounds their consecutive dynamical blocks. The resulting bound is explicit but huge and is not presented as sharp. The argument also uses essentially that $R_S \subset k(t)$ is a PID; the paper predicts an analogue for finite extensions of $k(t)$, but does not prove it for function fields of positive-genus curves.