

Bounds for preperiodic points for maps with good reduction

Sebastian Troncoso

arXiv:1608.05849

<http://arxiv.org/abs/1608.05849v3>

Good reduction as a counting principle

Let K be a number field, let S contain all archimedean places, and let $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be a degree- $d \geq 2$ morphism over K with good reduction outside S . Write $s = |S|$, and let $\text{Per}(\phi, K)$ and $\text{Tail}(\phi, K)$ denote the K -rational periodic and strictly preperiodic points.

Main theorem (Troncoso, Theorem 1.2 and Corollary 1.3). *If there are at least three K -rational tail points, then*

$$|\text{Per}(\phi, K)| \leq 2^{16s} + 3.$$

If there are at least four K -rational periodic points, then

$$|\text{Tail}(\phi, K)| \leq 4 \cdot 2^{16s}.$$

Without either cardinality hypothesis one still has

$$\begin{aligned} |\text{Per}(\phi, K)| &\leq 2^{16sd^3} + 3, \\ |\text{Tail}(\phi, K)| &\leq 4 \cdot 2^{16sd^3}, \\ |\text{PrePer}(\phi, K)| &\leq 5 \cdot 2^{16sd^3} + 3. \end{aligned} \tag{1}$$

The striking feature is the first pair of bounds: once one side of the finite orbit graph has enough marked vertices, the other side is bounded independently of d .

Ingredient I: good reduction forces integral separation

For $P = [x_1 : y_1]$ and $Q = [x_2 : y_2]$, let $\delta_{\mathfrak{p}}(P, Q)$ be their $\mathfrak{p}$ -adic logarithmic distance. Proposition 2.22 and Corollary 2.23 say that if R is a tail point, n is the period of its eventual cycle, and P is periodic with $P \neq \phi^{mn}(R)$ for every $m \geq 0$, then

$$\delta_{\mathfrak{p}}(P, R) = 0 \quad (\mathfrak{p} \notin S). \tag{2}$$

Indeed, after replacing ϕ by ϕ^n , the phase-aligned eventual cycle point $\phi^{mn}(R)$ is fixed (once m is large enough). The linear forms defining that fixed point and R both divide the pullback of the fixed-point linear form. Good reduction keeps all normalized coefficients integral, so comparing valuations along a periodic orbit makes both nonnegative summands vanish. Thus a dynamical statement becomes S -integrality.

Ingredient II: three anchors give one S -unit equation

A class-group obstruction prevents simultaneous coprime S -integral coordinates over K . Lemma 3.1 kills it in a fixed radical extension L : every $P \in \mathbf{P}^1(K)$ has coordinates $[x_P : y_P]$ that are coprime over the places above S , while every determinant $x_P y_Q - y_P x_Q$ lies in the radical of K^* . This radical S -unit group has free rank $s - 1$.

Fix three tail points P_1, P_2, P_3 , let n_i be the period of the eventual cycle of P_i , and take a periodic point $P = [x : y]$ distinct from every $\phi^{mn_i}(P_i)$. By (2),

$$x_i y - y_i x = u_i \quad (i = 1, 2, 3) \quad (3)$$

with radical S -units u_i . Eliminating x, y from (3) gives $A(u_1/u_3) + B(u_2/u_3) = 1$, with $A, B \neq 0$ fixed by the anchors. Beukers–Schlickewei bounds this equation by 2^{16s} solutions, and two of (3) recover $[x : y]$. For each anchor, the excluded expressions $\phi^{mn_i}(P_i)$ contain at most one periodic point: the single phase-aligned point on its eventual cycle. Thus there are at most three exceptions. Reversing the roles and using four periodic anchors splits tail points into four classes, giving the second bound.

Ingredient III: remove the hypotheses and improve the d -dependence

To deduce (1), one adjoins a chain of at most three preimages, or enough period-1, 2, 3 points, in an extension E/K of degree at most d^3 . The number of places above S is at most sd^3 , so Theorem 1.2 applies over E and descends to K .

Section 4 avoids putting this d^3 inside the exponent. For a non-rational periodic point $P_* = [a : b]$ of period 1, 2, or 3 and degree $r = [K(P_*) : K] \geq 3$, one has $r \leq d^3$, and the norm form

$$F(X, Y) = N_{K(P_*)/K}(aY - bX) \quad (4)$$

is irreducible of degree r . After one prime is adjoined to S to normalize P_* , (2) makes $F(x, y)$ an S' -unit with $|S'| \leq s + 1$. Evertse’s Thue–Mahler theorem bounds its solution classes by $(5 \cdot 10^6 r)^{s+1}$. If every periodic point of periods 1, 2, and 3 has degree at most two, an extension of degree at most four supplies four periodic anchors and Theorem 1.2 takes over.

Further finite enlargements of S principalize the remaining coordinate ideals. In the single- K -rational-periodic-point case, a dynatomic polynomial of formal period 2 or 3 and a gcd of ideal exponents perform the same normalization. This yields Theorem 1.4:

$$|\text{Tail}(\phi, K)| \leq d \max \left\{ (5 \cdot 10^6 (d^3 + 1))^{s+4}, 4 \cdot 2^{64(s+3)} \right\}. \quad (5)$$

The parallel periodic-point argument uses algebraic tail preimages and gives Theorem 1.5: if $\text{Tail}(\phi, K) \neq \emptyset$, then

$$|\text{Per}(\phi, K)| \leq \max \left\{ (5 \cdot 10^6 (d - 1))^{s+3}, 4 \cdot 2^{128(s+2)} \right\} + 1. \quad (6)$$

Toy model, hardest point, and scope

With three fixed anchors moved to $0, 1, \infty$, condition (2) says that a fourth point has both coordinate and complementary coordinate in an S -unit group: the whole counting problem is the classical equation $u + v = 1$. The full proof’s hardest point is making this picture uniform when the S -class group is nontrivial and the anchors are not K -rational; Lemma 3.1 and the norm form (4) are the two repairs.

The results concern morphisms of $\mathbf{P}^1$ over number fields and retain a good-reduction set S ; their explicit constants are not a proof of uniform boundedness in terms of $[K : \mathbf{Q}]$ and d alone. Section 5 constructs families with fixed S and respectively only two tail points or three periodic points for which the opposite set grows with d . Hence the thresholds in Theorem 1.2 cannot be lowered in a degree-independent statement.