

Moduli spaces of sheaves on K3 surfaces and Galois representations

Sarah Frei

arXiv:1810.06735

<http://arxiv.org/abs/1810.06735v3>

From a K3 surface to all cohomology of its sheaf moduli

Let k be any field, $\bar{k}$ an algebraic closure, and $\ell \neq \text{char } k$. For a k -variety X , write $\bar{X} = X \times_k \bar{k}$. If S is a K3 surface and $M = M_H(S, v)$ is a smooth proper moduli space of stable sheaves, then M need not be birational to the Hilbert scheme $S^{[n]}$ of the same dimension. The paper nevertheless shows that their rational ℓ -adic cohomologies carry the same Galois representation.

Main theorem (Theorems 1 and 2). *Let S_1, S_2 be K3 surfaces over k and suppose*

$$H_{\text{et}}^2(\bar{S}_1, \mathbf{Q}_\ell) \simeq H_{\text{et}}^2(\bar{S}_2, \mathbf{Q}_\ell)$$

as $\text{Gal}(\bar{k}/k)$ -representations. If M_i is a smooth proper moduli space of stable sheaves on S_i and $\dim M_1 = \dim M_2$, then for every $j \geq 0$,

$$H_{\text{et}}^j(\bar{M}_1, \mathbf{Q}_\ell) \simeq H_{\text{et}}^j(\bar{M}_2, \mathbf{Q}_\ell)$$

as Galois representations. Consequently, over a finite field, $Z(S_1, t) = Z(S_2, t)$ implies $Z(M_1, t) = Z(M_2, t)$.

For the standard construction, v is effective and geometrically primitive, H is v -generic, and $v^2 \geq 0$; Proposition 1.8 then gives a nonempty smooth projective geometrically irreducible M of dimension $v^2 + 2$. Geometric primitivity and v -genericity are important because the argument excludes properly semistable sheaves, but it does not require M to be a fine moduli space.

Step 1: the Mukai lattice controls degree two

The weight-zero ℓ -adic Mukai lattice is

$$\tilde{H}(\bar{S}, \mathbf{Z}_\ell) = H^0(\bar{S}, \mathbf{Z}_\ell) \oplus H^2(\bar{S}, \mathbf{Z}_\ell(1)) \oplus H^4(\bar{S}, \mathbf{Z}_\ell(2)),$$

with pairing $(\alpha, \beta) = -\alpha_0\beta_4 + \alpha_2\beta_2 - \alpha_4\beta_0$. A quasi-universal sheaf on $S \times M$ defines the Mukai map. Proposition 2.1 proves the Galois-equivariant isometries

$$H^2(\bar{M}, \mathbf{Z}_\ell(1)) \simeq \begin{cases} v^\perp, & v^2 > 0, \\ v^\perp / \langle v \rangle, & v^2 = 0. \end{cases}$$

Existence of a quasi-universal rather than universal sheaf is what lets this include non-fine moduli spaces. The proof establishes the known complex isometry first, then uses the Lefschetz principle in characteristic zero and a mixed-characteristic lift with smooth base change in positive characteristic.

When $v^2 = 0$, the moduli space has dimension two. Propositions 2.1 and 3.1 compare its degree-two representation with that of S , while its odd cohomology vanishes; degrees zero and four are the canonical representations of a geometrically irreducible smooth proper surface. This is the separate argument of Remark 3.3 and completes the isotropic branch of the theorem.

Step 2: comparison with the Hilbert scheme

Put $n = (v^2 + 2)/2$ and $w = (1, 0, 1 - n)$, the Mukai vector for $S^{[n]}$. Proposition 3.1 constructs a Galois-equivariant rational isometry $v^\perp \simeq w^\perp$ when $v^2 > 0$: reflection through $v - w$ or $v + w$ sends v to $\pm w$, with separate elementary reflections handling the isotropic exceptions. For $v^2 = 0$ it similarly identifies $v^\perp / \langle v \rangle$ with $H^2(\bar{S}, \mathbf{Q}_\ell(1))$. The use of $\mathbf{Q}_\ell$ is essential; Example 3.2 exhibits an integral discriminant obstruction over $\mathbf{Z}_5$.

For $v^2 > 0$, degree two is not enough by itself. Proposition 5.1 promotes the chosen isometry $g: \tilde{H}(\bar{S}, \mathbf{Q}_\ell) \rightarrow \tilde{H}(\bar{S}, \mathbf{Q}_\ell)$ with $g(v) = w$ to a correspondence

$$\gamma_g: H^*(\bar{M}, \mathbf{Q}_\ell) \xrightarrow{\sim} H^*(\bar{S}^{[n]}, \mathbf{Q}_\ell)$$

that is both Galois equivariant and a ring isomorphism. Its class is built from the normalized Mukai classes of the quasi-universal families, the isometry g , and the universal polynomial converting Chern characters to Chern classes.

The hardest point is proving that this rational correspondence is an isomorphism. Over $\mathbf{C}$, Markman's theorem gives the claim for integral isometries. Ring multiplicativity and invertibility are closed conditions on the isometry torsor $I = \text{Isom}(\tilde{H}(S), v, w)$. Lemma 5.3 uses Borel density to show that the integral points of I are Zariski dense, so the closed conditions hold at the desired $\mathbf{Q}_\ell$ -point. Lefschetz and lifting arguments then carry the result to an arbitrary field, while the construction itself makes γ_g Galois equivariant.

Step 3: change the K3 surface

For two surfaces, Proposition 4.1 uses the de Cataldo–Migliorini motivic decomposition of $S^{[n]}$ into Tate twists of symmetric powers $S^{(m)}$. Since the cohomology of a K3 surface is concentrated in degrees 0, 2, 4, its full Galois representation is determined by H^2 ; hence the assumed isomorphism for S_1, S_2 propagates to their Hilbert schemes. In the positive-square case, composing with the two maps γ_g proves Theorem 2; the isotropic case was already closed above. The Lefschetz trace formula then gives the finite-field zeta-function statement.

The toy case is $v = w$, where $M = S^{[n]}$ and the comparison is already visible. The general proof shows that changing the Mukai vector does not change rational Galois cohomology, provided the dimension stays fixed. It does not assert birationality, derived equivalence, or a natural geometric map between M_1 and M_2 ; nor does an arbitrary degree-two representation isomorphism necessarily yield a ring isomorphism between their cohomologies. The conclusion is precisely an isomorphism of graded Galois representations, which is already enough to identify all finite-field point counts.