

Ample canonical heights for endomorphisms on projective varieties

Takahiro Shibata

arXiv:1710.05278

<http://arxiv.org/abs/1710.05278v2>

A canonical scale for non-polarized dynamics

Let X be a smooth projective variety over $\overline{\mathbf{Q}}$, let $f : X \rightarrow X$ be a surjective endomorphism, and fix an ample height $h_X \geq 1$. Write δ_f for the first dynamical degree. Shibata introduces the smallest integer $l_f \geq 0$ for which

$$\frac{h_X(f^n(x))}{\delta_f^n n^{l_f}}$$

is bounded above for every $x \in X(\overline{\mathbf{Q}})$, and defines the upper and lower ample canonical heights $\overline{h}_f, \underline{h}_f$ by the corresponding limsup and liminf. Their values depend on h_X , but their comparison class and the zero locus

$$Z_f(K) := \{x \in X(K) : \underline{h}_f(x) = 0\}$$

do not. The proposed dynamical Northcott principle (Conjecture 1.3) says that, when $\delta_f > 1$, every number field K admits an f -invariant proper closed subset containing $Z_f(K)$.

Main theorem (Shibata, Theorem 1.5). *Assume $\delta_f > 1$. Conjecture 1.3 holds in each of the following cases:*

1. $f^*H \equiv \delta_f H$ for an ample $\mathbf{R}$ -divisor H (in particular, $\rho(X) = 1$);
2. $\rho(X) \leq 2$ and f is an automorphism;
3. X is an abelian variety;
4. X is a smooth projective surface.

Since preperiodic points lie in Z_f , the theorem implies that $\text{Preper}_f(K)$ is not Zariski dense in all four cases (Theorem 8.2). The conclusion is weaker than finiteness because a general endomorphism can have positive-dimensional exceptional dynamics.

Ingredient I: isolate the expanding eigendirection

If an ample H satisfies $f^*H \equiv \delta_f H$, then $l_f = 0$ and both ample canonical heights are comparable with the usual nef canonical height $\widehat{h}_{H,f}$. Northcott's theorem then gives $Z_f = \text{Preper}_f$ and finiteness over each number field.

For a rank-two automorphism, the two boundary rays of the nef cone give eigendivisors D, D' with eigenvalues δ_f, δ_f^{-1} . Since $D + D'$ is ample, its normalized forward height has the limit

$$\delta_f^{-n} h_{D+D'}(f^n(x)) \longrightarrow \widehat{h}_{D,f}(x); \quad (1)$$

the inverse eigendirection decays like δ_f^{-2n} . On a surface the same idea needs a correction: $D^+ + D^-$ is nef and big, and the curves on which it has degree zero are exactly the finitely many periodic curves. Subtracting an effective divisor supported on them makes the class ample; their contribution still disappears after division by δ_f^n . Consequently $l_f = 0$, $\overline{h}_f \asymp \underline{h}_f \asymp \widehat{h}^+$, and Theorem 6.1 identifies

$$Z_f(\overline{\mathbf{Q}}) = \text{Per}_f(\overline{\mathbf{Q}}) \cup \bigcup_{C \text{ periodic}} C(\overline{\mathbf{Q}}). \quad (2)$$

Ingredient II: Jordan growth on an abelian variety

First let f be a group endomorphism of an abelian variety. The powers of f satisfy a rational linear recurrence in the finite-dimensional algebra $\text{End}(X)_{\mathbf{Q}}$. After Jordan decomposition,

$$f^n = \sum_i \lambda_i^n n^{l_i} g_i,$$

ordered lexicographically by $(|\lambda_i|, l_i)$. For a Néron–Tate quadratic form, the leading term gives

$$\widehat{h}_X(f^n(x)) = |\lambda_1|^{2n} n^{2l_1} \widehat{h}_X(g_1(x)) + o(|\lambda_1|^{2n} n^{2l_1}). \quad (3)$$

Thus $(\delta_f, l_f) = (|\lambda_1|^2, 2l_1)$, and the height vanishes precisely where the real and imaginary parts of g_1 vanish. A linear-algebra lemma replaces these real endomorphisms by finitely many rational ones, so their common kernel is an algebraic subgroup: $B + X_{\text{tor}}$ for a proper abelian subvariety B .

For a general morphism $f = \tau_P \circ \phi$, an isogeny decomposition separates the expanding part of ϕ from a factor of dynamical degree one. A translation conjugates the expanding part back to a group endomorphism. The result is the exact description

$$Z_f(\overline{\mathbf{Q}}) = B(\overline{\mathbf{Q}}) + P_0 + X(\overline{\mathbf{Q}})_{\text{tor}}, \quad (4)$$

where $B + P_0$ is proper and f -invariant (Theorem 5.1). Passing to X/B , the image S of $Z_f(K)$ lies in the finite set $\pi(P_0) + (X/B)(K)_{\text{tor}}$. Since $Z_f(K)$ is f -invariant, S is invariant under the induced quotient map; hence the finite union $\pi^{-1}(S)$ is a proper f -invariant closed subset containing $Z_f(K)$.

Ingredient III: reduce surface maps to curves

For a non-automorphic surface map, first pass to an iterate and contract invariant (-1) -curves; this preserves (δ_f, l_f) and sends Z_f into the zero locus downstairs. The surface classification used in Theorem 7.5, after finite étale covers and further iterates, leaves projective planes, abelian surfaces, ruled or bielliptic surfaces, and properly elliptic surfaces. The nontrivial cases carry a fibration $\pi : X \rightarrow C$ with either $\pi f = \pi$ or $\pi f = g\pi$ and $\delta_g = \delta_f$. The type-(II) ruled and bielliptic cases have $\rho(X) \leq 2$, so $l_f = 0$; functorial height comparison then gives $l_g \leq l_f$, hence $l_g = 0$. Equality of the full pairs $(\delta_f, l_f) = (\delta_g, l_g)$ now yields

$$Z_f(K) \subseteq \pi^{-1}(Z_g(K)), \quad (5)$$

and polarized height theory on the curve makes $Z_g(K)$ finite. Type-(I) ruled surfaces over curves of genus at least two and properly elliptic surfaces are not potentially dense, while projective planes and abelian surfaces were handled earlier. This classification-and-descent step is the global geometric input that completes the fourth case.

Toy model, hardest point, and scope

For $f(x, y) = (ax + by, ay)$ on E^2 , $a \geq 2$, one has $f^n(0, y) = (na^{n-1}by, a^n y)$; the Néron–Tate height grows like $a^{2n}n^2$. The factor n^{l_f} is therefore essential, not cosmetic. The hardest new mechanism is turning the leading analytic term in (3) into the algebraic exceptional translate (4); the surface case then relies on a deep classification theorem. No general result is claimed for $\delta_f = 1$ or for arbitrary rational self-maps, and the paper does not prove Conjecture 1.3 in general beyond the stated classes. It also proves bounded time lag for intersections of two dense orbits when $\delta_f = \delta_g > 1$, $l_f = l_g$, and Conjecture 1.4 holds for both maps (in particular, under the four-class hypotheses of Theorem 8.9). If both maps are étale, the meeting-time pairs form finitely many arithmetic progressions.