Examples of dynamical degree equals arithmetic degree

Shu Kawaguchi, Joseph H. Silverman

arXiv:1212.3015

http://arxiv.org/abs/1212.3015v1

Problem and setup

Let $X/\overline{\mathbf{Q}}$ be a normal projective variety and $f : X \dashrightarrow X$ a dominant rational map. Its first dynamical degree is

$$\delta_f = \lim_{n\to\infty} \rho\big((f^n)^*|_{\mathrm{NS}(X)_{\mathbf{R}}}\big)^{1/n}.$$

For a point P whose forward orbit is defined and an ample Weil height h_X, put $h_X^+ = \max\{1, h_X\}$ and consider

$$\alpha_f(P) = \lim_{n\to\infty} h_X^+(f^n(P))^{1/n},$$

when the limit exists. The known fundamental inequality is $\overline{\alpha}_f(P) \le \delta_f$, where $\overline{\alpha}_f$ and $\underline{\alpha}_f$ denote the corresponding limsup and liminf. Conjecture 1 predicts that the limit always exists; that a Zariski-dense orbit has $\alpha_f(P) = \delta_f$; that every arithmetic degree is an algebraic integer; and that only finitely many arithmetic degrees occur.

Main theorem (Theorems 2 and 3)**.** *All four assertions of Conjecture 1 hold in each of the following cases:*

1. *f is a morphism and $\mathrm{NS}(X)_{\mathbf{R}} = \mathbf{R}$;*
2. *f is the extension to $\mathbf{P}^N$ of a regular affine automorphism of $\mathbf{A}^N$;*
3. *X is a smooth projective surface and f is an automorphism;*
4. *$f : \mathbf{P}^N \dashrightarrow \mathbf{P}^N$ is monomial, for points in the dense torus $\mathbf{G}_m^N$.*

Moreover, if $f : \mathbf{A}^2 \to \mathbf{A}^2$ extends dominantly to $\mathbf{P}^2$ and either f is algebraically stable or $\deg f = 2$, then

$$\{P \in \mathbf{A}^2(\overline{\mathbf{Q}}) : \alpha_f(P) = \delta_f\}$$

contains a Zariski-dense set of points with pairwise disjoint forward orbits.

The second assertion is evidence rather than a proof that every dense orbit has maximal arithmetic degree: it constructs many good orbits, without classifying all of them.

Proof mechanism for the four complete cases

1. Turn an eigendivisor into exponential height growth. If $\delta_f = 1$, the fundamental bounds immediately give $1 \le \underline{\alpha}_f(P) \le \overline{\alpha}_f(P) \le 1$; thus the arithmetic degree exists and equals 1. Assume now that $\delta_f > 1$. If an ample class D satisfies $f^*D \equiv \delta_f D$, its canonical height obeys $\widehat{h}_D \circ f = \delta_f \widehat{h}_D$. For a morphism with one-dimensional Néron–Severi space this applies to every ample class; positivity away from preperiodic points gives $\alpha_f(P) = \delta_f$, while preperiodic points give 1.

For a regular affine automorphism, the forward canonical height is positive exactly at wandering affine points. Points at infinity enter the invariant locus $I_{f^{-1}}$; after normalization this locus has Néron–Severi rank one, so the morphism case controls its arithmetic degrees. This is where regularity $I_f \cap I_{f^{-1}} = \varnothing$ is essential. More precisely, Theorem 8 gives degree 1 at periodic points, δ_f at wandering affine points, and δ_g at wandering boundary points, where $g = f|_{I_{f^{-1}}}$. The boundary value need not equal δ_f, but such an orbit is not Zariski dense in $\mathbf{P}^N$.

2. On surfaces, pair expanding and contracting heights. For a surface automorphism with $\delta_f > 1$, Theorem 9 supplies nef eigendivisors and nonnegative heights $\widehat{h}^{\pm}$ satisfying

$$\widehat{h}^+ \circ f = \delta_f \widehat{h}^+, \qquad \widehat{h}^- \circ f = \delta_f^{-1} \widehat{h}^-.$$

Their sum is a height for a big and nef divisor. Off the finite union E_f of periodic curves, a nonperiodic point has $\widehat{h}^+ > 0$, so the expanding term forces growth δ_f^n. On E_f, restriction to curve automorphisms gives arithmetic degree 1. For monomial maps, earlier toric height results identify arithmetic degrees with eigenvalues of the exponent matrix. Theorem 12 sharpens this: a point in $\mathbf{G}_m^N$ with dense orbit has degree δ_f, while a submaximal value confines the whole orbit to a proper invariant algebraic subgroup. Algebraic integrality and finiteness follow because a fixed integer exponent matrix has only finitely many eigenvalues.

Why the planar construction works

3. A weak lower height converts one local escape rate into the global answer. The case $\delta_f = 1$ is already settled by the same fundamental inequality, so Proposition 14, Theorem 19, and Theorem 16 address the remaining case $\delta_f > 1$. For a divisor D define

$$\widehat{h}^{\circ}_{f,D}(P) = \liminf_{n\to\infty} \delta_f^{-n} h_D(f^n(P)).$$

Proposition 14 proves $\widehat{h}^{\circ}_{f,D}(P) > 0 \Rightarrow \alpha_f(P) = \delta_f$: the liminf supplies the lower bound, an ample height dominating h_D transfers it to $\underline{\alpha}_f(P)$, and the fundamental inequality supplies the matching upper bound. The iterate formulas

$$\delta_{f^m} = \delta_f^m, \qquad \widehat{h}^{\circ}_f(P) = \min_{0 \le i < m} \delta_f^{-i} \widehat{h}^{\circ}_{f^m}(f^i(P))$$

(Lemma 18) justify passing to an iterate when the useful point at infinity is periodic rather than fixed.

4. Detect positivity near a periodic point at infinity. Suppose an iterate fixes a point Q_0 on the hyperplane at infinity and is defined there. After replacing f by that iterate, algebraic stability gives $\delta_f = d = \deg f$, and suitable coordinates put

$$f = [aX_1^d + G_1, G_2, \ldots, G_N, W^d],$$

where every G_i vanishes at $Q_0 = [1, 0, \ldots, 0]$. At a good nonarchimedean place, the neighborhood with X_1 a unit and all other coordinates in the maximal ideal is forward invariant. For an affine point with last coordinate β, no common p-adic factor can be cancelled from the iterates, and the W-coordinate becomes β^{d^n}. Consequently

$$h(f^n(P)) \ge d^n \log |\beta|_p^{-1},$$

so the weak lower height is positive (Theorem 19).

5. Produce the required neighborhood. For algebraically stable maps of $\mathbf{A}^2$, the leading homogeneous terms factor as $(F, G) = H(F_0, G_0)$ and induce $\varphi = [F_0, G_0]$ on the line at infinity. If $\deg \varphi \ge 2$, one chooses a periodic orbit avoiding the finite zero set of H. If $\deg \varphi = 0$, algebraic stability forces the image point to be a genuine fixed point. When $\deg \varphi = 1$, a periodic point can fail to exist outside indeterminacy; the proof instead builds an invariant p-adic sector $|x|_p > |y|_p > |z|_p$ in which one coordinate grows as d^n. For non-algebraically-stable quadratic maps, Guedj's classification reduces the argument to explicit normal forms, each admitting the same kind of local growth estimate. The resulting p-adic open set contains a Zariski-dense collection of algebraic points; choosing successively outside previously chosen orbits gives the disjoint-orbit conclusion.

Toy model, difficulty, and scope

For $f(z) = z^d$ on $\mathbf{P}^1$, $\delta_f = d$ and the usual canonical height satisfies $\widehat{h}(f^n(P)) = d^n \widehat{h}(P)$: preperiodic points have arithmetic degree 1, all others have degree d. This illustrates the canonical-height mechanism used in the paper; the higher-dimensional cases need not retain this two-valued dichotomy, because boundary dynamics or different eigenvalues can produce additional arithmetic degrees.

The hardest step is not the final height comparison but finding a region on which cancellation cannot destroy the predicted exponential growth; the periodic-at-infinity normal form and the quadratic classification do exactly that. The planar theorem does not assert the full conjecture for all polynomial maps, and the monomial conclusion is restricted to the dense torus. Several complete-case arguments also import established canonical-height results rather than rebuilding them in this paper. The converse to Proposition 14 is left open: equality $\alpha_f(P) = \delta_f$ need not presently be known to produce any divisor whose weak lower canonical height is positive (Question 15).