

Fano varieties with large Seshadri constants in positive characteristic

Ziquan Zhuang

arXiv:1707.02682

<http://arxiv.org/abs/1707.02682v2>

Local positivity in a log Fano pair

Unless stated otherwise, the paper works over an algebraically closed field k of characteristic $p > 0$. Let X be a normal projective n -fold, $\Delta \geq 0$ an effective $\mathbf{Q}$ -divisor, and

$$L = -(K_X + \Delta)$$

a $\mathbf{Q}$ -Cartier ample divisor. At a smooth point x , with blowup $\sigma : Y = \text{Bl}_x X \rightarrow X$ and exceptional divisor E , the Seshadri constant is

$$\varepsilon(L, x) = \sup\{t > 0 : \sigma^*L - tE \text{ is ample}\}.$$

It measures positivity concentrated at one point, yet the theorem shows that crossing the dimension forces the entire variety to be projective space.

Main theorem (Zhuang, Theorem 1.3). *Under the hypotheses above, let $x \in X$ be smooth.*

1. *If $\varepsilon(L, x) > n$, then $X \simeq \mathbf{P}^n$.*
2. *If $\varepsilon(L, x) \geq n$ and $p > 2$, then X is globally F -regular.*

Part (1) allows arbitrary singularities away from x and has no restriction beyond positive characteristic. Part (2) supplies the positive-characteristic replacement for the vanishing and singularity control normally obtained from characteristic-zero methods.

Ingredient I: blow up exactly at the dimensional threshold

After a harmless base change one may take a very general point outside $\text{Supp}(\Delta)$. Since

$$K_Y = \sigma^*K_X + (n-1)E, \quad -(K_Y + \Delta_Y + E) = \sigma^*L - nE, \quad (1)$$

the strict inequality makes the right-hand side ample. Moreover $E \simeq \mathbf{P}^{n-1}$, $N_{E/Y} \simeq \mathcal{O}_E(-1)$, and $\Delta_Y \cap E = \emptyset$. Thus the original local inequality produces a log Fano pair containing a smooth projective-space divisor and with $\rho(Y) \geq 2$.

Ingredient II: Frobenius restores vanishing and semiampleness

Lemma 2.3 is the basic engine. If E is a prime divisor contained in the smooth locus of Y , with $E \not\subset \text{Supp}(\Delta_Y)$, $-(K_Y + E + \Delta_Y)$ is nef and big with ample restriction to E , and the restricted pair is globally F -regular, then (Y, Δ_Y) is globally F -regular. The proof compares Frobenius trace maps on Y and E ; Serre vanishing makes the restriction map on sections surjective, so a splitting on E lifts to Y . Global F -regularity then yields the Kodaira-type vanishing

$$H^i(Y, \mathcal{O}_Y(D)) = 0 \quad (i > 0) \quad (2)$$

for an effective Weil divisor D when $D - (K_Y + \Delta_Y)$ is $\mathbf{Q}$ -Cartier, nef, and big.

This vanishing controls the threshold for subtracting a projective-space divisor. Lemmas 3.5–3.8 show that the threshold is rational and that the limiting nef divisor is semiample: if its stable base locus were nonempty, Frobenius methods produce a positive-dimensional subvariety on which it is numerically trivial; such a subvariety must meet E , contradicting the base-point-free restriction to E .

Ingredient III: classify the exceptional divisor

Lemma 3.11 applies to a pair (Y, Δ_Y) containing $E \simeq \mathbf{P}^{n-1}$ with $-(K_Y + \Delta_Y + E)$ ample, $\rho(Y) \geq 2$, and $E \cap \Delta_Y = \emptyset$. The semiample threshold defines a morphism $g : Y \rightarrow Z$. Its restriction embeds E , a Hodge-index argument rules out numerical triviality on E , and the adjoint contraction lemma rules out a birational g . Vanishing from (2) shows every fiber is an integral rational curve. Consequently

$$Z \simeq \mathbf{P}^{n-1}, \quad Y \simeq \mathbf{P}_{\mathbf{P}^{n-1}}(\mathcal{O} \oplus \mathcal{O}(-d)), \quad (3)$$

with E a section. In the blowup situation, $N_{E/Y} \simeq \mathcal{O}_E(-1)$ forces $d = 1$ and identifies E as the unique negative section. Hence Y is the blowup of $\mathbf{P}^n$ at a point, and contracting E gives $X \simeq \mathbf{P}^n$.

The threshold case and proof roadmap

For part (2), $\varepsilon(L, x) \geq n$ implies $L^n \geq n^n$. If the volume is strict, Corollary 2.4 applies Lemma 2.3 directly to the blowup. Suppose instead that $L^n = n^n$, so $D = \sigma^*L - nE$ has $D^n = 0$. The paper first proves that D is semiample and analyzes the resulting rational conic fibration. It then separates two branches. If $\Delta \neq 0$, Step 2 proves that Δ is $\mathbf{Q}$ -Cartier and replaces it by $(1 - c)\Delta$ for $0 < c \ll 1$; this perturbs the problem to strict volume, so Corollary 2.4 applies. In the remaining branch $\Delta = 0$ and $p > 2$, the general conic is reduced and E is a double section. Step 4 reduces the Frobenius calculation according to the finite map $g|_E : E \rightarrow Z$. In the ramified case, $Z \simeq \mathbf{P}(1^{n-1}, 2)$ and $M \subset Z$ is its ramification hyperplane; surjectivity of the trace on Y is reduced to that for the globally F -regular toric pair $(Z, \frac{1}{2}M)$. The quasi-etale case is treated separately by the analogous, simpler trace argument. Thus Y , and hence the required X , is globally F -regular.

The proof of part (1) can therefore be read as the chain

$$\varepsilon(L, x) > n \implies \text{ample divisor (1) on the blowup} \implies \text{classification (3)} \implies X \simeq \mathbf{P}^n.$$

Part (2) replaces the first strict implication by a careful analysis of the nef boundary and its conic fibration.

Toy model, hardest point, and scope

For $X = \mathbf{P}^n$ and $L = -K_X = (n + 1)H$, one has $\varepsilon(L, x) = n + 1$. The blowup is exactly $\mathbf{P}_{\mathbf{P}^{n-1}}(\mathcal{O} \oplus \mathcal{O}(-1))$, and the exceptional divisor is its negative section. This model explains both why the strict bound detects projective space and why the equality value n admits additional geometries classified later in the paper.

The hardest point is manufacturing, without general positive-characteristic MMP or Kawamata–Viehweg vanishing, the semiample contraction needed for the $\mathbf{P}^{n-1}$ -divisor classification; Frobenius trace and global F -regularity provide the substitute. The global F -regularity conclusion excludes $p = 2$. The separate weak-boundedness theorem concerns only threefolds in characteristic $p > 5$ and uses moving Seshadri constants; it is not part of the projective-space criterion proved above.