Linear growth and moduli spaces of rational curves

Jakob Glas

Jakob Glas, arXiv:2505.07645v1 (2025), 16 pages

http://arxiv.org/abs/2505.07645v1

From rational-curve spaces to function-field points

Let X be a smooth Fano variety over $\mathbf{F}_q$. A point of $X(\mathbf{F}_q(t))$ is a morphism $\mathbf{P}^1 \to X$, so height strata are $\mathbf{F}_q$-points of spaces of rational curves. For an embedding $X \subset \mathbf{P}^{n-1}$ and an open $U \subset X$, write $\mathrm{Mor}_{e,U}(\mathbf{P}^1, X)$ for degree-e maps whose image meets U, and

$$N_U(e) = \#\{x \in U(\mathbf{F}_q(t)) : H(x) = q^e\}$$

for naive height. Expected dimension can therefore become a counting bound without computing cohomology or Frobenius traces.

Main theorem (Glas, Theorems 1.2–1.4)**.** *If $X \subset \mathbf{P}^{n-1}$ is smooth Fano and is defined by forms of degrees $d_1, \dots, d_R$, then*

$$N_U(e) = O\Big(C^e q^{\dim \mathrm{Mor}_{e,U}(\mathbf{P}^1, X)}\Big), \qquad C = \prod_{i=1}^{R} d_i^{d_i},$$

with an implied constant depending on $q, d_1, \dots, d_R$.

Over a field K, let $X \subset \mathbf{P}^{n-1}$ instead be either a smooth intersection of two quadrics, or a smooth cubic hypersurface with $\operatorname{char} K > 3$. If $\dim X \geq 3$, then every component of $\mathrm{Mor}_e(\mathbf{P}^1, X)$ has the expected dimension, namely

$$\begin{cases} e(n-4)+n-3, & X \text{ is an intersection of two quadrics,} \\ e(n-3)+n-2, & X \text{ is cubic,} \end{cases} \qquad e \geq 1.$$

Over $\mathbf{F}_q$ this gives respectively $N_X(e) = O(16^e q^{e(n-4)})$ and $N_X(e) = O(27^e q^{e(n-3)})$.

These are upper bounds, not asymptotic formulas. Their extra factor is exponential in e with fixed base 16 or 27; viewed as a power of q^e, that loss becomes arbitrarily small when q is large.

Ingredient 1: a uniform finite-field degree bound

For any locally closed $Y \subset \mathbf{A}^N$ over $\mathbf{F}_q$, Proposition 2.1 proves the elementary but robust inequality

$$\#Y(\mathbf{F}_q) \leq \deg(Y) q^{\dim Y}.$$

One closes Y, separates top-dimensional from smaller components, slices by the parallel hyperplanes $x_i = a$, and inducts on dimension; Bezout controls the degrees of all slices.

Write a degree-e map to $\mathbf{P}^{n-1}$ as $x(t) = (x_1(t) : \cdots : x_n(t))$. Expanding $F_i(x(t))$ yields $d_i e + 1$ equations of degree d_i in its coefficients. The morphism space is an open subset of their projective zero locus (the open conditions impose exact degree and no common factor). Passing to affine cones and applying Bezout gives Proposition 3.1:

$$\#Z(\mathbf{F}_q) \leq 2 d_1 \cdots d_R \, C^e q^{\dim Z}$$

for any union Z of components of the morphism space. Taking the closure of $\mathrm{Mor}_{e,U}$ proves the general theorem. As a toy model, one degree-d equation contributes $de + 1$ coefficient equations, so its degree cost is at most d^{de+1} while its dimension determines the power of q.

A parallel application: del Pezzo surfaces

For a smooth del Pezzo surface of degree $1 \le d \le 5$, use anticanonical height and remove all rational curves C with $-K_X \cdot C \le 1$. Theorem 1.1 concerns the cumulative count

$$N_U^{\le}(B) = \#\{x \in U(\mathbf{F}_q(t)) : H_{-K_X}(x) \le B\}$$

and gives

$$N_U^{\le}(q^e) = O(C_d^e q^e), \qquad (C_5, C_4, C_3, C_2, C_1) = (1024, 16, 27, 4^4, 6^6).$$

The imported theorem of Beheshti–Lehmann–Riedl–Tanimoto says that the only non-expected components parametrize multiple covers of precisely those low-degree curves. The counting proposition then applies to the remaining components. Anticanonical embeddings treat degrees 3–5; weighted projective models $\mathbf{P}(2,1,1,1)$ and $\mathbf{P}(3,2,1,1)$ supply the explicit Bezout constants for degrees 2 and 1.

Ingredient 2: the line case and its exceptional points

To prove expected dimension for all e, the paper first calculates fibers of the space of lines through a point. For an intersection of two quadrics $X \subset \mathbf{P}^n$, every such fiber has dimension $n-5$. For a cubic $X \subset \mathbf{P}^n$, it has dimension $n-4$ off the Eckardt locus and $n-3$ at an Eckardt point, where $X \cap T_x X$ is a cone with vertex x.

Finiteness of the Eckardt locus is the hardest positive-characteristic step. The characteristic-zero shortcut through the differential of the Gauss map is unavailable. Instead, a smooth hyperplane section through any line reduces the number of Eckardt points on that line to the cubic-surface case; in characteristic different from 2 each line contains at most two. A line through an Eckardt point must be of type II, and the type-II locus in the Fano scheme has dimension at most $n-3$ when the characteristic is not 2 or 3. The incidence correspondence between Eckardt points and type-II lines then has fibers of dimension $n-3$, forcing the Eckardt locus to be zero-dimensional (Corollary 4.10).

Ingredient 3: bend-and-break propagates degree one

Work in the proper Kontsevich spaces $\overline{\mathcal{M}}_{0,k}(X, e)$ and let $\mathrm{ev}_e : \overline{\mathcal{M}}_{0,1}(X, e) \to X$ be evaluation. The basic bend-and-break lemma says that no complete curve of irreducible-domain two-pointed maps can lie in one fiber of the two-point evaluation map. Consequently, if a closed substack $Y \subset \mathrm{ev}_e^{-1}(x)$ has dimension at least $\dim X$, then it contains reducible maps and their locus has codimension at most one in Y (Corollary 5.2).

For a complete intersection $X \subset \mathbf{P}^n$ of codimension R and total defining degree d, Proposition 5.3 sets

$$e_0 = \left\lceil \frac{d-R}{n+1-d} \right\rceil.$$

If, through degree e_0, the evaluation fibers have dimension

$$\dim \mathrm{ev}_e^{-1}(x) = e(n+1-d) - 2 + \delta_{x \in S}$$

for a finite exceptional set S, then this equality holds for every e. The proof inducts simultaneously on the degree of the reducible locus. A boundary map splits into lower-degree maps glued over X, so the inductive fiber bounds put that boundary one dimension below the target bound. Any oversized irreducible component is large enough for bend-and-break; its codimension-one boundary then contradicts the just-established estimate.

For two quadrics and cubics of dimension at least three, $e_0 \le 1$. Thus the line calculations provide the base case, with $S = \varnothing$ for two quadrics and S the finite Eckardt set for cubics. Quotienting the irreducible-domain map space by PGL_2 translates the evaluation-fiber bounds into the displayed expected dimensions. The method depends on the varieties being constant over $\mathbf{F}_q$ for the counting application, gives no asymptotic constant, and retains the theorem's dimension and characteristic restrictions.