

Projectivity and Birational Geometry of Bridgeland Moduli spaces

Arend Bayer, Emanuele Macrì

arXiv:1203.4613

<http://arxiv.org/abs/1203.4613v2>

A divisor extracted from stability

Let X be a smooth projective complex variety, let $\sigma = (Z, \mathcal{A})$ be a numerical Bridgeland stability condition on $D^b(X)$, and fix a numerical class v . Suppose a proper algebraic space S carries an S -perfect family $\mathcal{E}$ of σ -semistable objects of class v . For an integral projective curve $C \subset S$, set

$$\ell_{\sigma, \mathcal{E}} \cdot C = \Im \left(-\frac{Z(\Phi_{\mathcal{E}}(\mathcal{O}_C))}{Z(v)} \right), \quad \Phi_{\mathcal{E}}(-) = (p_X)_*(\mathcal{E} \otimes p_S^*(-)). \quad (1)$$

The expression is unchanged by rescaling Z or twisting the family from the base.

Main theorem (Bayer–Macrì, Theorems 1.1 and 1.3). *Formula (1) defines a nef numerical Cartier divisor class on S , and*

$$\ell_{\sigma, \mathcal{E}} \cdot C = 0 \iff \mathcal{E}_c \text{ and } \mathcal{E}_{c'} \text{ are } S\text{-equivalent for general } c, c' \in C.$$

If X is a projective K3 surface, $v \in H_{\text{alg}}^(X, \mathbb{Z})$, and $\sigma \in \text{Stab}^\dagger(X)$ is generic for v , then the coarse moduli space $M_\sigma(v)$ exists as a normal, projective, irreducible, $\mathbb{Q}$ -factorial variety, and the induced class ℓ_σ is ample.*

The first statement is valid far beyond K3 surfaces and uses no GIT presentation. The K3 statement answers both existence and positivity: stability conditions inside a chamber map naturally into the ample cone of the same moduli space.

Ingredient I: categorical positivity on a curve

After rotating the stability condition, assume $Z(v) = -1$. Lemma 2.5 first replaces σ by finitely many nearby *algebraic* stability conditions having the same semistable objects of class v , with Z in the convex hull of their central charges. Their hearts are Noetherian, so the Abramovich–Polishchuk construction extends each heart $\mathcal{A}$ to a Noetherian heart $\mathcal{A}_S \subset D^b(S \times X)$. For a family over a projective curve and every sufficiently positive line bundle L on C , Lemma 3.6 gives

$$\Phi_{\mathcal{E}}(L) \in \mathcal{A}.$$

The imaginary part of each algebraic central charge is nonnegative on its heart, and replacing $\mathcal{O}_C$ by L does not change (1); convexity then gives nefness for the original Z (and zero for Z forces zero for every summand).

The equality criterion is the sharper part of the Positivity Lemma 3.3. If the degree is zero, the exact sequence $0 \rightarrow L(-c) \rightarrow L \rightarrow k(c) \rightarrow 0$ becomes an exact sequence in $\mathcal{A}$; all terms have the same phase, so the stable factors of every general fibre $\mathcal{E}_c$ lie in one fixed finite list. Semicontinuity then forces the fibres to be S-equivalent (Lemma 3.7). Conversely, a family of generically S-equivalent fibres admits, away from finitely many points, a relative Jordan–Hölder filtration with quotients $F_i \boxtimes L_i$ (Lemma 3.9). Each F_i has the phase of v , so its contribution to the imaginary part in (1) vanishes. The resulting curve functional descends to $N_1(S)$ (Theorem 4.1) and agrees with a Donaldson determinant class (Proposition 4.4).

Ingredient II: turning nefness into K3 projectivity

For a generic stability condition on a K3 surface, the proof first moves dense nearby conditions within the same chamber to one admitting a primitive isotropic Mukai vector w of the same phase as v (Lemma 7.3). The moduli space $Y = M_\tau(w)$ is itself a K3 surface, possibly with a Brauer class, and its quasi-universal family gives a Fourier–Mukai equivalence

$$D^b(X) \simeq D^b(Y, \alpha).$$

After applying this equivalence and shifting by $[-1]$, the objects of class v become twisted Gieseker-semistable sheaves. Classical sheaf-moduli theory supplies the coarse projective space and its normality, irreducibility, and $\mathbb{Q}$ -factoriality; chamber constancy transports it back to $M_\sigma(v)$.

For primitive v with $v^2 > 0$, write $Z(u) = (u, \eta(\sigma))$ under the Mukai pairing and set $w_\sigma = \Im \eta(\sigma)$. The Mukai isometry identifies the Beauville–Bogomolov square of ℓ_σ with $w_\sigma^2 > 0$ (Theorem 6.10 and Lemma 7.4). Thus the Positivity Lemma gives a big nef class with positive degree on every curve. Triviality of the canonical bundle and the base-point-free theorem make it semiample, and the absence of contracted curves makes it ample (Corollary 7.5). Mukai-square -2 and 0 cases, including their multiples, are handled separately by Lemmas 7.1–7.2. The remaining nonprimitive positive-square case is treated by comparison with determinant line bundles on the corresponding Gieseker moduli space and density inside the chamber.

Ingredient III: stability walls become birational walls

Let v be primitive, let σ_0 be a generic point of a wall, and take nearby generic conditions $\sigma_\pm$ on its two sides. The wall families on $M_{\sigma_\pm}(v)$ are σ_0 -semistable, so (1) gives nef classes $\ell_{\sigma_0, \pm}$. Their Beauville–Bogomolov squares are positive; hence they are big and, by base-point-freeness, induce contractions

$$\pi_\pm: M_{\sigma_\pm}(v) \longrightarrow Y_\pm.$$

The zero-curve criterion identifies exactly what is contracted: curves whose general members become S-equivalent on the wall.

If σ_0 -stable objects exist and the complements of the σ_0 -stable loci in $M_{\sigma_+}(v)$ and $M_{\sigma_-}(v)$ have codimension at least two, Theorem 1.4 proves a dichotomy. Either both wall classes are ample and the wall-crossing map extends to an isomorphism, or neither is ample and the two small contractions have the same target $Y_+ = Y_-$; crossing the wall is their flop, with the wall divisor classes identified. This is the promised bridge from the wall-and-chamber decomposition of $\text{Stab}^\dagger(X)$ to the minimal-model geometry of $M_\sigma(v)$.

A toy picture

Inside a chamber, a nonconstant curve of stable objects has pairwise nonisomorphic general fibres; stability makes S-equivalence equal to isomorphism, so the theorem reads $\ell_\sigma \cdot C > 0$. On a wall, some objects acquire the same graded Jordan–Hölder object, and precisely those curves have degree zero. The wall divisor therefore behaves like a GIT linearization at a wall even though its construction used only the central charge and the relative derived category.

Applications and limits

The paper uses this mechanism to obtain lattice-controlled ample regions for sheaf moduli, determine the nef cone of $\text{Hilb}^n(X)$ in a Picard-rank-one range, and construct new birational Lagrangian fibrations (Theorem 10.8). The wall-crossing theorem is deliberately partial: it does not cover totally semistable walls or the case where the stable locus has divisorial complement. These are not cosmetic exclusions; the codimension-two identification is what lets divisor classes be compared across the birational map.