

Kawaguchi-Silverman conjecture for endomorphisms on several classes of varieties

Yohsuke Matsuzawa

arXiv:1902.06072

<http://arxiv.org/abs/1902.06072v4>

Height growth and the zero-locus problem

Let X be a normal projective variety over $\overline{\mathbf{Q}}$ and $f: X \rightarrow X$ a surjective endomorphism. Its first dynamical degree δ_f is the spectral radius of $f^*: N^1(X)_{\mathbf{R}} \rightarrow N^1(X)_{\mathbf{R}}$. For an ample height $h_X \geq 1$, the arithmetic degree of x is the exponential growth rate of $h_X(f^n(x))$. The Kawaguchi–Silverman conjecture (KSC) predicts $\alpha_f(x) = \delta_f$ whenever the orbit of x is Zariski dense.

For $\delta_f > 1$, let l_f be the least nonnegative integer for which $h_X(f^n(x))/(n^{l_f} \delta_f^n)$ is bounded above for every $x \in X(\overline{\mathbf{Q}})$, and put

$$\hat{h}_f(x) = \liminf_{n \rightarrow \infty} \frac{h_X(f^n(x))}{n^{l_f} \delta_f^n}.$$

Shibata’s conjecture says that, over each fixed number field K , the zero set of this ample canonical height lies in an f -invariant proper closed subset. This immediately forces KSC for a dense orbit.

Main theorem (Theorem 4.1). *Assume $\delta_f > 1$, that $\text{Pic}(X)_{\mathbf{Q}} \simeq N^1(X)_{\mathbf{Q}}$, and that $\text{Nef}(X)$ is generated by finitely many semi-ample integral divisors. Then Shibata’s conjecture holds for f . In particular, every $x \in X(\overline{\mathbf{Q}})$ with Zariski-dense f -orbit satisfies $\alpha_f(x) = \delta_f$.*

The hypotheses include Mori dream spaces, varieties of Fano type, and projective toric varieties. Their role is quite exact: polyhedrality gives finitely many dynamical directions, semi-ampleness turns each direction into a genuine quotient, and $\text{Pic}_{\mathbf{Q}} = N_{\mathbf{Q}}^1$ upgrades numerical eigenvectors to linear equivalences suitable for heights.

Diagonalizing the nef cone geometrically

After replacing f by an iterate, f^* fixes every extremal ray of the rational polyhedral nef cone (Lemma 4.6). Choose its semi-ample integral generators $D_1, \dots, D_r$ so that

$$f^*D_i \sim \lambda_i D_i, \quad \lambda_1 = \dots = \lambda_s = \delta_f > \lambda_{s+1} \geq \dots \geq \lambda_r > 0.$$

The integer λ_i is not merely a numerical eigenvalue. Semi-ampleness gives a morphism $\pi_i: X \rightarrow Y_i$ and an ample H_i with $D_i \sim \pi_i^* H_i$; Lemma 4.5 descends f to a surjective $g_i: Y_i \rightarrow Y_i$ satisfying $g_i \pi_i = \pi_i f$ and $g_i^* H_i \sim \lambda_i H_i$.

For $\lambda_i > 1$, polarized canonical heights therefore exist and satisfy

$$\hat{h}_{D_i} = \hat{h}_{H_i} \circ \pi_i \geq 0, \quad \hat{h}_{D_i} \circ f = \lambda_i \hat{h}_{D_i}.$$

This descent is the decisive bridge: each extremal numerical direction becomes an ordinary polarized dynamical system, where Northcott finiteness is available.

The zero locus and the proof roadmap

The sum $H = D_1 + \dots + D_r$ is ample. Directions with $\lambda_i < \delta_f$ disappear after normalization by δ_f^n , whereas the top eigendirections survive. Thus $l_f = 0$ and Proposition 4.7 identifies

$$\hat{h}_f(x) = \sum_{i=1}^s \hat{h}_{D_i}(x) = \sum_{i=1}^s \hat{h}_{H_i}(\pi_i(x)).$$

All terms are nonnegative. For a number field K containing the data,

$$Z_f(K) = \bigcap_{i=1}^s \pi_i^{-1} \{y \in Y_i(K) : \hat{h}_{H_i}(y) = 0\}.$$

Each set in braces is finite by Northcott because H_i is ample and $g_i^* H_i \sim \delta_f H_i$; moreover $\dim Y_i > 0$. The intersection is therefore the K -points of a proper closed subset. The elementary relation between the zero loci for f and for an iterate transfers this conclusion back from the chosen power of f .

The proof may be read as four moves: permute and then fix the extremal nef rays; descend each ray through its semi-ample fibration; retain only the δ_f -eigendirections in the normalized ample height; and apply Northcott separately on their bases. The hard point is the second move: numerical spectral data alone do not produce a canonical height, while semi-ampleness and the Picard hypothesis produce both the quotient Y_i and the exact linear relation needed to define one.

A model and the paper's wider scope

On $X = \mathbf{P}^1 \times \mathbf{P}^1$, take $f(u, v) = (u^a, v^b)$ with $a > b > 1$. The two nef rays are the two projections, and $\delta_f = a$. After division by a^n , the second coordinate contributes nothing: $\hat{h}_f(u, v) = \hat{h}_a(u)$. Over a fixed number field the zero-height u -coordinates form a finite set, so the zero locus is a finite union of vertical fibers. The theorem replaces these coordinate projections by the semi-ample maps π_i and makes the same argument on an arbitrary polyhedral nef cone.

The paper also develops a distinct equivariant-MMP route. For a $\mathbf{Q}$ -factorial klt variety admitting an int-amplified endomorphism, Theorem 5.8 proves KSC for automorphisms, for maps whose every iterate is primitive, and when $\kappa(X) \geq 0$; Corollary 5.13 treats a rationally connected threefold when $\kappa(-K_X) > 0$. Finally, Theorem 6.1 proves the dense-orbit assertion for group endomorphisms of connected linear algebraic groups by reducing the dynamical degree to a maximal torus. These results use different hypotheses: the polyhedral-nef theorem is not a blanket KSC statement for every endomorphism of every normal projective variety.