

Arithmetic degrees and dynamical degrees of endomorphisms on surfaces

Yohsuke Matsuzawa, Kaoru Sano, Takahiro Shibata

arXiv:1701.04369

<http://arxiv.org/abs/1701.04369v2>

Two growth rates on a surface

Let k be a number field, let X be a smooth projective surface over $\bar{k}$, and let $f: X \rightarrow X$ be a surjective endomorphism. For an ample divisor H , the first dynamical degree is

$$\delta_f = \lim_{m \rightarrow \infty} ((f^m)^* H \cdot H)^{1/m},$$

while the arithmetic degree of $P \in X(\bar{k})$, when the limit exists, is

$$\alpha_f(P) = \lim_{m \rightarrow \infty} \max\{h_H(f^m(P)), 1\}^{1/m}.$$

The first measures numerical pullback growth; the second measures height growth along one orbit. Kawaguchi–Silverman conjectured existence of the second limit and the equality $\alpha_f(P) = \delta_f$ whenever $\mathcal{O}_f(P)$ is Zariski dense.

Main theorem (Matsuzawa–Sano–Shibata, Theorem 1.3). *For every surjective endomorphism f of a smooth projective surface as above, the arithmetic degree exists at every algebraic point. If $\mathcal{O}_f(P)$ is Zariski dense in X , then*

$$\alpha_f(P) = \delta_f. \tag{1}$$

Existence for morphisms and the inequality $\alpha_f(P) \leq \delta_f$ were already known. Thus the substantive task is to force the reverse inequality for dense orbits, or to show that the geometry forbids such an orbit.

Ingredient I: make the assertion invariant under reduction

Two functorial devices let the proof replace X without losing (1). First, passing from f to a positive iterate changes both degrees by the same power. Second, if a finite surjective map $\psi: X \rightarrow Y$ intertwines f_X and f_Y , then

$$\delta_{f_X} = \delta_{f_Y}, \quad \alpha_{f_X}(P) = \alpha_{f_Y}(\psi(P)). \tag{2}$$

The first equality follows from intersection theory after pulling back an ample divisor; the second is functoriality of heights.

The paper also proves birational invariance along a dense orbit. If $\mu: X \dashrightarrow Y$ conjugates f to g , ample heights are comparable on an open set where μ is an isomorphism. A dense orbit has a subsequence there, so, provided the relevant limits exist,

$$\alpha_f(P) = \alpha_g(\mu(P)). \tag{3}$$

This allows contraction of periodic (-1) -curves after replacing f by an iterate. It is the bridge from arbitrary surfaces to the relatively minimal classification.

Ingredient II: split by Kodaira dimension

Automorphisms were already known, so assume that f is finite of degree greater than one.

If $\kappa(X) = -\infty$, equivariant contraction reduces to a rational minimal surface or a $\mathbb{P}^1$ -bundle $\pi: X \rightarrow C$. The rational minimal surfaces are $\mathbb{P}^2$ or Hirzebruch surfaces: $\mathbb{P}^2$ and $\mathbb{F}_0 \simeq \mathbb{P}^1 \times \mathbb{P}^1$ are known cases, while the remaining $\mathbb{F}_e$ are handled by the ruled argument that follows. For a ruled surface, an iterate preserves the fibers. Writing F for a fiber and C_0 for a normalized section, one obtains

$$f^*F \equiv aF, \quad f^*C_0 \equiv cF + dC_0, \quad \delta_f = \max\{a, d\}, \quad (4)$$

where a and d are respectively the degrees on the base and a general fiber. When the normalized bundle has invariant $e > 0$, preservation of the nef cone and the intersection form force $c = 0$ and $a = d$. A dense orbit projects densely to C , and the fiber-divisor height then gives

$$\alpha_f(P) \geq \lim_{m \rightarrow \infty} h_F(f^m(P))^{1/m} = \alpha_{f_C}(\pi(P)) = a = \delta_f. \quad (5)$$

The genus-one, degree-zero bundle case is the delicate exception. Fiber degree one is handled directly by the base-height argument. When the fiber degree is greater than one—in particular in the potentially problematic case $d > a$ —a finite étale base change makes the bundle $\mathbb{P}(\mathcal{O}_C \oplus L)$. Its two disjoint zero-self-intersection sections constrain their images under f . If L is torsion, another finite cover trivializes the bundle. If L is non-torsion, the section calculation and the induced action on $\text{Pic}^0(C)$ rule out $d > a$; hence again $\delta_f = a$, and (5) applies. For $g(C) \geq 2$, the induced base automorphism has finite order, so no orbit on X can be dense.

Ingredient III: the nonnegative Kodaira cases

If $\kappa(X) \geq 0$ and f is nontrivial, classification results used in the paper imply that X is minimal and f is étale.

For $\kappa(X) = 0$, the identity $\chi(\mathcal{O}_X) = \deg(f)\chi(\mathcal{O}_X)$ leaves only abelian and hyperelliptic surfaces. The abelian case was known. A hyperelliptic surface becomes a product of elliptic curves after a finite étale base change; after a further cover the endomorphism lifts. Formula (2) therefore descends the abelian result.

For $\kappa(X) = 1$, the Iitaka fibration $\pi: X \rightarrow C$ is elliptic and is respected by an automorphism f_C of the base. If $m_i F_i$ are its multiple fibers, the canonical bundle formula and $\chi(\mathcal{O}_X) = 0$ give

$$2(g(C) - 1) + \sum_i \frac{m_i - 1}{m_i} > 0. \quad (6)$$

The map f_C permutes their base points. If $g(C) \geq 2$, its automorphism group is finite; if $g(C) = 1$, (6) supplies a fixed point after iteration and the remaining group automorphism has finite order; if $C \simeq \mathbb{P}^1$, (6) supplies at least three fixed points. Thus some iterate of f_C is the identity, and an orbit is trapped over finitely many fibers. There is no Zariski-dense orbit. Finally, general-type surfaces have only finite-order automorphisms, so they contribute no nontrivial case.

A product model

On $X = \mathbb{P}^1 \times \mathbb{P}^1$, consider $f(x, y) = (x^a, y^d)$ with $a, d \geq 2$. The two ruling classes are eigenvectors of f^* with eigenvalues a, d , so $\delta_f = \max\{a, d\}$. For a point whose two coordinates are nonpreperiodic and whose orbit is Zariski dense, the two coordinate heights grow as a^m and d^m ; hence $\alpha_f(P) = \max\{a, d\}$. This is exactly the ruled-surface mechanism without twisting. The full proof must show that the line bundle defining a nontrivial bundle cannot hide a faster fiber eigenvalue from the base-height argument.

Hard point and scope

The hard work is not an estimate for a single height but the exhaustive geometric reduction. In particular, the genus-one ruled case couples intersection theory, finite étale descent, two distinguished sections, and the isogeny action on $\text{Pic}^0(C)$ to recover the dominant eigenvalue. Once that is done, a single divisor height supplies the missing lower bound.

The theorem concerns everywhere-defined surjective endomorphisms of smooth projective surfaces; it does not settle arbitrary rational self-maps. Its proof imports the already-known automorphism, abelian-variety, product, and projective-space cases. The paper also proves that every surjective endomorphism in any dimension has some point with $\alpha_f(P) = \delta_f$, and obtains a Zariski-dense family of pairwise disjoint full-growth orbits for automorphisms; those separate results are not needed for Theorem 1.3.