

Geometry of the moduli space of n -pointed K3 surfaces of genus 11

Ignacio Barros

arXiv:1705.05290

<http://arxiv.org/abs/1705.05290v4>

The birational question

Let $\mathcal{F}_{g,n}$ parameterize primitively polarized complex K3 surfaces (S, H) of genus g (so $H^2 = 2g - 2$) together with n ordered points of S . Although Mukai's models make $\mathcal{F}_{11}$ unirational, adding points introduces powers of a K3 surface in the fibers, so unirationality is not inherited formally. Barros obtains a replacement parameterization from nodal curves in $|H|$.

Main theorem (Theorem 0.1). *The moduli space $\mathcal{F}_{11,n}$ is unirational for $n \leq 6$ and uniruled for $n \leq 7$. Consequently $\mathcal{M}_{11,n}$ is unirational for $n \leq 6$.*

The last assertion uses Mukai's rational map $\mathcal{M}_{11,n} \dashrightarrow \mathcal{F}_{11,n}$, birationally a $\mathbf{P}^{11-n}$ -bundle in the relevant range.

The correspondence space

Let $\mathcal{V}_{g,\delta,l}$ parameterize tuples

$$(S, H, X, x_1, \dots, x_\delta, y_1, \dots, y_l),$$

where $X \in |H|$ is irreducible with marked nodes x_i , and the y_j are marked smooth points of X . If $\nu : C \rightarrow X$ is the normalization and $\nu^{-1}(x_i) = \{p_i, q_i\}$, there are two maps

$$\mathcal{F}_{g,\delta+l} \xleftarrow{\pi} \mathcal{V}_{g,\delta,l} \xrightarrow{c_{g,\delta,l}} \mathcal{M}_{g-\delta, [2\delta]+l}.$$

The left map forgets X . The right map remembers $[C, p_1 + q_1, \dots, p_\delta + q_\delta, y_1, \dots, y_l]$; the notation $[2\delta]$ means quotienting by $(\mathbf{Z}/2)^\delta$, which exchanges the two branches over each node. The stronger structural result is that $c_{g,\delta,l}$ is dominant for $3 \leq g \leq 11$, $0 \leq \delta \leq g - 2$, $g \neq 10$, and is birational when $g = 11$.

Three proof mechanisms

- Deformations make normalization dominant.** Germs of vector fields on S tangent to the pointed nodal curve form a sheaf $T_S(X | y_1, \dots, y_l)$. Its H^1 is the tangent space to $\mathcal{V}_{g,\delta,l}$, while the differential of $c_{g,\delta,l}$ is induced by restriction to $T_X(-y_1 - \dots - y_l)$. The cokernel is $H^2(S, T_S(-X) \otimes I_y)$, dual to $H^0(S, \Omega_S^1(H))$. The latter vanishes for a general polarized K3 in the stated range, proving dominance (Corollary 2.4).
- Nodes and markings impose the expected conditions.** For the composite $f : C \xrightarrow{\nu} X \hookrightarrow S$, the relevant normal sheaf satisfies $N_f \simeq \omega_C$; blowing up the marked points compares the deformation spaces. For general branch and marked points, the differential of π is surjective once

$$h^0(C, \mathcal{O}_C((p_1 + q_1) + \dots + (p_\delta + q_\delta) + y_1 + \dots + y_l)) \leq 1.$$

This holds when $2\delta + l \leq g - \delta$, equivalently $3\delta + l \leq g$, giving dominance of π (Corollary 2.7).

- A cone degeneration makes the generic normalization fiber irreducible.** Dimension count makes $c_{11,\delta,l}$ generically finite. To prove degree one, the paper degenerates every K3 extension of a nodal canonical curve X to the projective cone S_X . That cone point lies in every component of the fiber. Smoothness there is reduced to

$$H^0(S_X, N_{S_X/\mathbf{P}^g}(-1)) \simeq \bigoplus_{k \geq 1} H^0(X, N_{X/\mathbf{P}^{g-1}}(-k)).$$

After specializing X to a trivalent canonical graph curve, this space has dimension $23 - g$, exactly the lower bound for the fiber. Upper semicontinuity therefore gives a smooth common point; the general fiber is irreducible, hence a single point in the generically finite case.

From curves back to K3 surfaces

Take $(\delta, l) = (2, 4)$. Then $\mathcal{V}_{11,2,4}$ is birational to a $(\mathbf{Z}/2)^2$ -quotient of $\mathcal{M}_{9,8}$, which is unirational, and it dominates $\mathcal{F}_{11,6}$. Thus $\mathcal{F}_{11,6}$, and hence every $\mathcal{F}_{11,n}$ with $n \leq 6$, is unirational. With $(\delta, l) = (2, 5)$, the analogous quotient of $\mathcal{M}_{9,9}$ is uniruled and maps dominantly and generically finitely to $\mathcal{F}_{11,7}$, proving uniruledness.

In the useful dimension count, a node costs three conditions and a marked point one. At $3\delta + l = g$ the residual system is expected to contain one curve; Lemma 4.2 proves this and makes π birational. The hard point is instead the normalization fiber's irreducibility, where the cone and graph-curve degenerations are essential.

Limits of the method

The paper also proves $\text{Kod}(\mathcal{F}_{11,n}) \geq 0$ for $n \geq 9$ via $\mathcal{M}_{10,10}/(\mathbf{Z}/2)$ and the universal Jacobian. Its method does not settle $n = 8$, because the available covering rational curves are contracted; genus 10 is excluded from the vanishing and graph-curve range.

Codex mathematical paper digest, 2026-10-04. Source paper should be consulted for full statements and proofs.