

Moduli Spaces of Rational Graphically Stable Curves

Andy Fry

arXiv:2212.08157

<http://arxiv.org/abs/2212.08157v1>

A graph-controlled boundary for rational curves

Fix a simple connected graph Γ on the labels $2, \dots, n$ and single out the marking p_1 . The component containing p_1 is the root. A stable rational n -pointed curve is Γ -stable if every non-root component meeting the rest of the curve in one node contains two markings p_i, p_j with $e_{ij} \in E(\Gamma)$. Equivalently, one contracts precisely the tails not containing p_1 whose labels form an independent set in Γ . Smyth's extremal-assignment formalism therefore gives a modular compactification $\overline{\mathcal{M}}_{0,\Gamma}$, with smooth locus $\mathcal{M}_{0,\Gamma}$; Lemma 3.2 shows that its boundary is a simple normal crossings divisor.

Let N be the number of edges deleted from the complete graph K_{n-1} to obtain Γ (after relabelling, take $e_{23} \in E(\Gamma)$). Pluecker coordinates give a torus embedding

$$\mathcal{M}_{0,\Gamma} \hookrightarrow T_\Gamma = T^{\binom{n}{2} - n - N}. \quad (1)$$

For a boundary divisor D_I , the orders of the coordinate ratios x_{ij}/x_{23} define a divisorial valuation vector $\mathbf{v}_{D_I}$ and hence a map

$$\pi_\Gamma : \Delta(\partial \overline{\mathcal{M}}_{0,\Gamma}) \longrightarrow N_{T_\Gamma, \mathbb{R}}. \quad (2)$$

Main theorem (Fry, Theorem 3.13). *The map π_Γ embeds the cone complex $\mathcal{M}_{0,\Gamma}^{\text{trop}}$ as a balanced fan if and only if Γ is complete multipartite. For such Γ , (1) has*

$$\text{trop}(\mathcal{M}_{0,\Gamma}) \cong \mathcal{M}_{0,\Gamma}^{\text{trop}}, \quad (3)$$

and the closure of $\mathcal{M}_{0,\Gamma}$ in the toric variety $X(\mathcal{M}_{0,\Gamma}^{\text{trop}})$ is exactly $\overline{\mathcal{M}}_{0,\Gamma}$.

A graph is complete multipartite when its vertices split into independent parts and every pair from different parts is joined. Equivalently, its complement is a disjoint union of cliques, or no three vertices span exactly one edge (Lemma 3.9).

Pluecker projection and valuation injectivity

The open moduli space is the quotient of an open part of $G(2, n)$: a matrix of homogeneous coordinates for the markings has minor $x_{ij} = x_i y_j - x_j y_i$, nonzero whenever $i = 1$ or $e_{ij} \in E(\Gamma)$. Dropping the minors belonging to missing edges and then quotienting by column rescaling yields (1). Lemmas 3.4–3.5 identify this algebraic projection with the tropical projection pr_Γ . Since the boundary is snc, geometric tropicalization gives

$$\text{trop}(\mathcal{M}_{0,\Gamma}) = \text{pr}_\Gamma(\mathcal{M}_{0,n}^{\text{trop}}) \quad (4)$$

with weight one on maximal cones (Proposition 3.6).

The central combinatorial issue is whether (2) remembers every boundary stratum. In the chosen coordinates, for $|I| \geq 3$ one has

$$\mathbf{v}_{D_I} = \sum_{\substack{\{i,j\} \subset I \\ e_{ij} \in E(\Gamma)}} \mathbf{v}_{D_{\{i,j\}}}, \quad (5)$$

with the analogous complementary formula when $\{2, 3\} \subset I$. If Γ is not complete multipartite, Lemma 3.9 supplies vertices i, j, k with e_{ij} present and e_{ik}, e_{jk} absent. Equation (5) then collapses two distinct divisors:

$$\pi_\Gamma(D_{\{i,j,k\}}) = \pi_\Gamma(D_{\{i,j\}}). \quad (6)$$

Conversely, for complete multipartite Γ , equality of two valuation vectors forces the induced subgraphs on their index sets to have the same edges. Any label occurring in only one index set would be isolated in a complete multipartite stable subgraph, which is impossible. Thus the index sets coincide. This proves Lemma 3.10 and the fan-level equivalence.

Cross ratios promote the fan to the compactification

Injectivity alone does not yet identify the toric closure. The geometric tropicalization criterion used in Theorem 3.13 asks, for every boundary stratum S , that

1. each boundary divisor through S be isolated by a unit having valuation 1 there and 0 on the other such divisors; and
2. S be very affine, with the relevant restriction map on unit lattices surjective.

Lemma 3.12 says that all units are products of cross ratios, equivalently forgetful maps to $\mathcal{M}_{0,4}$. If the forbidden one-edge triple from (6) exists, every cross ratio detecting $D_{\{i,j,k\}}$ also detects $D_{\{i,j\}}$, so the first condition fails. When Γ is complete multipartite, one can choose four markings $\{1, a, b, c\}$ with $e_{ab} \in E(\Gamma)$ so that the associated cross ratio keeps precisely the chosen divisor nodal and smooths the others through S . This verifies the first condition. Each stratum is a product of smaller graphically stable moduli spaces, hence is very affine, and the local coordinate-hyperplane description of the snc boundary makes restriction of units surjective. The criterion then identifies the closure in the toric variety with the modular compactification.

Toy obstruction and scope

For the complete bipartite graph $K_{2,2}$, the complement is two disjoint edges, so no forbidden triple occurs and all boundary rays remain distinct. By contrast, remove e_{35} and e_{45} from K_4 . The triple $\{3, 4, 5\}$ spans only e_{34} , and (6) identifies $D_{\{3,4,5\}}$ with $D_{\{3,4\}}$. In the paper's Example 3.14 the modular boundary has eight rays and nine two-dimensional cones, whereas the tropicalization has only seven rays and eight such cones. The toric closure replaces a limiting $\mathbb{P}^1$ of curves by a single point. This small example isolates the full theorem's mechanism: a missing graph edge becomes a forgotten Pluecker coordinate, and enough forgotten coordinates make distinct degenerations indistinguishable.

For non-complete-multipartite graphs, (4) and the geometric tropicalization still exist; what fails is faithful realization of the modular boundary cone complex by π_Γ , and consequently the stated toric closure need not be $\overline{\mathcal{M}}_{0,\Gamma}$. The theorem concerns genus zero and this specific graphical stability construction; it does not classify arbitrary compactifications or arbitrary torus embeddings.