

On the K-stability of Fano varieties and anticanonical divisors

Kento Fujita, Yuji Odaka

arXiv:1602.01305

<http://arxiv.org/abs/1602.01305v2>

Testing K-stability with anticanonical bases

Work over an algebraically closed field of characteristic zero. Let X be an n -dimensional $\mathbb{Q}$ -Fano variety: X is projective and klt, and $-K_X$ is ample and $\mathbb{Q}$ -Cartier. For a basis $s_1, \dots, s_{N_k}$ of $H^0(X, -kK_X)$, with $D_i = \text{div}(s_i)$, form the basis-type divisor

$$D = \frac{D_1 + \dots + D_{N_k}}{kN_k} \sim_{\mathbb{Q}} -K_X. \quad (1)$$

Define

$$\delta_k(X) = \inf_{\substack{D \text{ of} \\ k\text{-basis type}}} \text{lct}(X; D), \quad \delta(X) = \limsup_{k \rightarrow \infty} \delta_k(X). \quad (2)$$

Main theorem (Fujita–Odaka, Theorems 0.3 and 2.1). *If $\delta(X) > 1$, then $(X, -K_X)$ is uniformly K-stable. If $\delta(X) \geq 1$, then $(X, -K_X)$ is K-semistable.*

This turns a condition involving all test configurations into an asymptotic singularity test for concrete anticanonical divisors. The paper conjectures the converse, but proves only the sufficient direction stated above.

The valuative target

For a prime divisor F over X , write $A_X(F)$ for its log discrepancy, $\tau(F)$ for the pseudo-effective threshold of $-K_X$ along F , and $V = (-K_X)^n$. Set

$$\beta(F) = A_X(F)V - \int_0^\infty \text{vol}(-K_X - xF) dx, \quad (3)$$

$$j(F) = \int_0^{\tau(F)} (V - \text{vol}(-K_X - xF)) dx. \quad (4)$$

The valuative criterion recalled as Theorem 1.1 says that K-semistability is equivalent to $\beta(F) \geq 0$ for every F , while uniform K-stability is equivalent to $\beta(F) \geq c \cdot j(F)$ for one $c > 0$ and every F .

Concavity of volume gives

$$V\tau(F) \geq \int_0^\infty \text{vol}(-K_X - xF) dx \geq \frac{V\tau(F)}{n+1}. \quad (5)$$

Consequently, a uniform strict inequality

$$(1 - \varepsilon)A_X(F)V \geq \int_0^\infty \text{vol}(-K_X - xF) dx \quad (6)$$

implies the $\beta \geq cj$ criterion (Theorem 1.3). Thus the problem is to extract (6) from the singularities of basis-type divisors.

Ingredient I: a basis adapted to a valuation

Filter $H^0(-kK_X)$ by vanishing along F :

$$\mathcal{F}^a H^0(-kK_X) = H^0(-kK_X - aF).$$

Choose a basis compatible with this decreasing filtration. If $i_a = h^0(-kK_X - aF)$, summation by parts shows that the largest possible vanishing of (1) along F is

$$\sup_D v_F(D) = \frac{\sum_{a \geq 1} h^0(-kK_X - aF)}{k \cdot h^0(-kK_X)}. \quad (7)$$

Therefore Lemma 2.2 gives the exact discrepancy formula

$$\min_D A_{(X, cD)}(F) = A_X(F) - c \frac{\sum_{a \geq 1} h^0(-kK_X - aF)}{k \cdot h^0(-kK_X)}. \quad (8)$$

The minimum is important: the global infimum in $\delta_k(X)$ controls the worst basis simultaneously for every divisorial valuation.

Ingredient II: discrete vanishing becomes volume

By definition of $\delta_k(X)$, equation (8) implies

$$A_X(F) \geq \delta_k(X) \frac{\sum_{a \geq 1} h^0(-kK_X - aF)}{k \cdot h^0(-kK_X)}. \quad (9)$$

As $k \rightarrow \infty$, the quotient is a Riemann sum for the normalized volume integral:

$$\frac{\sum_{a \geq 1} h^0(-kK_X - aF)}{k \cdot h^0(-kK_X)} \longrightarrow \frac{1}{V} \int_0^{\tau(F)} \text{vol}(-K_X - xF) dx. \quad (10)$$

If $\delta(X) \geq 1$, (9)–(10) give $\beta(F) \geq 0$ for every F . If $\delta(X) > 1$, they leave a uniform positive margin, yielding (6) and hence uniform K-stability. This is the whole proof's central conversion: singularities of many explicit divisors encode the expected vanishing of every anticanonical linear series.

A one-dimensional model

For $X = \mathbb{P}^1$, take the monomial basis $x^i y^{2k-i}$ of $H^0(\mathcal{O}_{\mathbb{P}^1}(2k))$. At $F = \{x = 0\}$ the vanishing orders are $0, 1, \dots, 2k$, so their sum is $k(2k + 1)$. Dividing by $kN_k = k(2k + 1)$ in (1) gives $v_F(D) = 1$; in fact this basis produces $D = [0] + [\infty]$. Thus its log canonical threshold is exactly one. The toy case shows why the boundary value $\delta = 1$ naturally corresponds to semistability, whereas the theorem needs a strict gap above one for uniform K-stability.

Two further consequences

The Berman–Gibbs invariant $\gamma_k(X)$ is defined from the determinantal divisor on X^{N_k} along the diagonal. Blowing up the product center F^{N_k} compares its multiplicity with the same sum $\sum_{a \geq 1} h^0(-kK_X - aF)$ appearing in (7). Theorem 2.5 obtains

$$\delta_k(X) \geq \gamma_k(X),$$

so Berman–Gibbs stability implies uniform K-stability, and its semistable version implies K-semistability.

The same valuative language also gives a new algebraic proof of Tian's criterion. The alpha invariant satisfies

$$\alpha(X)\tau(F) \leq A_X(F). \quad (11)$$

For a dreamy F , finite generation produces a birational model on which an Okounkov body Δ_ε records the volume function. Its first-coordinate barycenter is the corresponding ε -truncated ratio; as $\varepsilon \rightarrow 0$, this ratio tends to the normalized integral in (10). The centroid bound for a convex body yields

$$\int_0^{\tau(F)} \text{vol}(-K_X - xF) dx \leq \frac{n}{n+1} \tau(F)V. \quad (12)$$

Combining (11)–(12) with the valuative criterion proves that $\alpha(X) > n/(n+1)$ implies uniform K-stability, while equality or a larger value implies K-semistability (Theorem 3.2).

Proof roadmap and the hard point

The main theorem follows in four moves: fix an arbitrary divisor F over X ; choose an anticanonical basis adapted to its vanishing filtration; identify the extremal discrepancy through (7)–(8); and pass from the discrete dimensions to the volume integral (10), where the valuative K-stability criterion applies.

The delicate point is uniformity. An arbitrary basis need not see the worst vanishing along F , and a basis optimized for one valuation need not be optimized for another. Taking the infimum over all basis-type divisors before passing to the limit makes (9) valid for every F with the same $\delta_k(X)$. The filtration-compatible basis then computes the exact worst case.

The result is a sufficient criterion, not the conjectured equivalence. It assumes characteristic zero and the $\mathbb{Q}$ -Fano (hence klt) hypotheses. Its alpha-invariant proof initially reduces to dreamy divisors through the cited valuative theory; the finite-generation model and Okounkov-body argument are not asserted for an arbitrary non-dreamy valuation.