Birational Geometry of the Space of Complete Quadrics

César Lozano Huerta

arXiv:1401.8050

http://arxiv.org/abs/1401.8050v2

Complete quadrics as a blow-up compactification

Work over $\mathbf{C}$. Let $N = \binom{n+2}{2} - 1$, so $\mathbf{P}^N$ parametrizes quadric hypersurfaces in $\mathbf{P}^n$. Write $\Phi_i \subset \mathbf{P}^N$ for the rank-$\leq i$ locus. The space X_n of complete $(n-1)$-quadrics is obtained by successively blowing up the strict transforms of

$$\Phi_1, \Phi_2, \ldots, \Phi_{n-1}.$$

Its boundary divisors $E_1, \ldots, E_n$ record a singular quadric together with the successive quadric markings on its singular loci; E_n is the strict transform of Φ_n. For $1 \leq i \leq n$, let H_i be the closure of the locus of smooth quadrics tangent to a fixed $(i-1)$-plane.

Main theorem (Theorems B and C)**.** *The divisor cones are*

$$\mathrm{Eff}(X_n) = \langle E_1, \ldots, E_n \rangle, \qquad \mathrm{Nef}(X_n) = \langle H_1, \ldots, H_n \rangle.$$

Moreover X_n is Fano, hence a Mori dream space. For each k, the model

$$X_n(H_k) = \mathrm{Proj} \bigoplus_{m \geq 0} H^0(X_n, mH_k)$$

is the normalization of the second-order Chow variety $\mathrm{Chow}_2(k-1, X_n)$ *parametrizing tangent $(k-1)$-planes to complete quadrics.*

For quadric surfaces ($n = 3$), Theorem A further says that $\mathrm{Eff}(X_3)$ has eight Mori chambers. The paper identifies the nef contractions, the small contraction across which X_3 flips, the flipped model, and an additional Grassmannian quotient model.

How the two cones are found

1. Tangency divisors and their dual flag curves. The H_i are basepoint-free. At a singular complete quadric one tests tangency on its marking quadric; if that marking is singular, the same test is repeated on the next marking. A general $(i-1)$-plane can therefore be chosen non-tangent at every stage.

At the maximally singular point, a complete quadric is a full flag

$$\mathrm{pt} = F_1 \subset F_2 \subset \cdots \subset F_{n+1} = \mathbf{P}^n.$$

Varying F_i between its neighbors gives an effective curve Fl_i with

$$\mathrm{Fl}_j \cdot H_i = \delta_{ij}.$$

Thus these curves span the cone dual to $\langle H_1, \ldots, H_n \rangle$, proving that no further nef direction exists.

2. Boundary pencils force effective coefficients to be positive. For $k < n$, fix a rank-k quadric and vary its marking in a general pencil; the resulting curve B_k sweeps E_k and satisfies

$$B_k \cdot E_k = -(n-k), \qquad B_k \cdot E_{k+1} = n-k+1,$$

with all other boundary intersections zero. If $D = \sum a_i E_i$ is a general effective divisor, it does not contain a general B_k, hence $(n-k)a_k \leq (n-k+1)a_{k+1}$. Intersecting with the pullback of a general pencil in the dual projective space gives $a_1 \geq 0$; the inequalities then give every $a_i \geq 0$. This proves the effective-cone statement.

The blow-up canonical-class calculation becomes

$$K_{X_n} = -2H_1 - H_2 - \cdots - H_{n-1} - 2H_n.$$

Since $-K_{X_n}$ lies in the interior of the computed nef cone, it is ample. The cited BCHM theorem then makes X_n a Mori dream space, so its effective cone admits a finite chamber decomposition.

Why the nef models are Chow models

A smooth quadric is a symmetric map $Q : V \to V^*$. Its kth wedge $\bigwedge^k Q$ is a quadratic form on $\bigwedge^k V$. If $L \in G(k-1, n)$, then

$$L^t(\textstyle\bigwedge^k Q)L = \bigwedge^k (L^t Q L) = \det(Q|_L),$$

so its zero divisor is exactly the locus of $(k-1)$-planes tangent to Q. This identifies $\bigwedge^k Q$ with the second-order Chow form and extends to a morphism $\rho_k : X_n \to \mathrm{Chow}_2(k-1, X_n)$.

The key comparison is numerical. The H_k-model contracts precisely the extremal flag curves Fl_j with $j \neq k$. In local triangular coordinates for a maximally singular marked quadric, an explicit wedge-power calculation shows that ρ_k contracts exactly the same curve classes. Two birational contractions with the same extremal rays have the same normal target; hence $X_n(H_k)$ is the normalization of the Chow image. The normalization is essential in the general statement.

For $n = 2$ this is the classical toy model: a complete conic remembers a conic and its dual, and the two nef generators give the projections to $\mathbf{P}^5$ and $\mathbf{P}^{5*}$. Higher k replace ordinary duality by the family of tangent $(k-1)$-planes.

The surface chamber calculation and its models

For X_3, the divisor identities

$$H_2 = 2H_1 - E_1, \qquad H_3 = 3H_1 - 2E_1 - E_2$$

and a new movable ray, for which we choose the positive representative

$$P = 2H_1 - H_2 + 2H_3 = E_3 + 2H_1 = E_1 + 2H_3,$$

put all relevant walls in one three-dimensional picture. The paper builds eight covering curve classes for X_3, the boundary divisors, and $E_1 \cap E_3$. Negative intersection forces the covered locus into the stable base locus. The involution $Q \mapsto \bigwedge^3 Q$ exchanges $E_1 \leftrightarrow E_3$ and $H_1 \leftrightarrow H_3$, cutting the work in half. Finally the displayed decompositions of P and basepoint-freeness of the H_i prove that no larger base locus occurs. This yields the eight chambers and

$$\mathrm{Mov}(X_3) = \langle H_1, H_2, H_3, P \rangle.$$

More explicitly, the eight stable base loci are

$$\varnothing, \quad E_1 \cap E_3, \quad E_3, \quad E_1, \quad E_1 \cup E_3, \quad E_2 \cup E_3, \quad E_1 \cup E_2, \quad E_2.$$

The wall tests are not merely necessary: expressions such as $P = E_3 + 2H_1 = E_1 + 2H_3$ bound the base locus from above, so the covering-curve calculation gives equality in every region.

The chamber geometry has concrete targets. The H_1- and H_3-models are $\mathbf{P}^9$ and its dual; the H_2-model is the second-order Chow variety of tangent lines. A divisor between H_1 and H_3 gives a small contraction to the $\mathbf{Z}/2$-quotient of the Chow variety of conics in $G(1,3)$. Its flip is

$$X_3^+ = \mathrm{Hilb}_{2x+1}(G(1,3))/(\mathbf{Z}/2),$$

where the involution exchanges the two rulings of a smooth quadric; the P-model is $G(2,5)/(\mathbf{Z}/2)$. The hardest point is globalizing this involution over degenerations and checking relative ampleness on both sides of the flip; a pencil of planes through the doubled supporting line supplies the decisive fiber curve.

Here the Hilbert scheme itself has a useful concrete construction:

$$\mathrm{Hilb}_{2x+1}(G(1,3)) \cong \mathrm{Bl}_{\mathrm{OG}}\, G(2,5),$$

where OG is the two-component orthogonal Grassmannian of planes contained in the Plücker quadric. Away from OG a plane cuts $G(1,3)$ in a conic; blowing up records the limiting conic when the plane is contained in the Grassmannian. A smooth quadric surface has two conics of lines, one for each ruling. Exchanging them defines the involution; in ranks one and three the two planes coincide, so the action preserves the blow-up center and extends globally.

The small contraction forgets scheme structure and retains the conic cycle. On the flipped side, the fiber over a doubled line $2L$ is the pencil $\mathbf{P}^1$ of planes containing L. If $D = aH_1 + bH_3 + cP$ with $a, b, c > 0$, this fiber has D-degree $c(\gamma \cdot P) > 0$. On the original side, the covering curve $C_{1,2} \subset E_1 \cap E_3$ has $C_{1,2} \cdot D = c(C_{1,2} \cdot P) < 0$. These opposite signs prove that D is relatively ample after the flip while $-D$ is relatively ample before it. The middle Chow model also has an enumerative shadow: its degree is 92, the number of smooth quadric surfaces tangent to nine general lines.

The paper works over $\mathbf{C}$ and computes the indicated divisor models, not every possible modular compactification of quadrics. Some identifications use earlier results on complete quadrics, normality of a Chow image, and the birational geometry of stable maps. Its principal contribution is the explicit bridge from rank-stratum blowups and intersection theory to a finite MMP whose main targets retain moduli meanings.

Codex mathematical paper digest, 2026-11-02. Source paper should be consulted for full statements and proofs.