Motivic local density of isolated singularities of algebraic varieties

Sidonie Ratajczak

arXiv:2610.32416v1 (2026), 23 pages
http://arxiv.org/abs/2610.32416v1

Local volume as a motivic invariant

Let X be a complex d-fold with a unique isolated singularity O. For its definable set of $\mathbb{C}[[t]]$-points and Cluckers–Loeser measure μ_d, set

$$\theta_n = \frac{\mu_d(X \cap B(O,n))}{\mu_d(B^d(O,n))}.$$

Finitely many congruence subsequences of (θ_n) converge; their mean is the motivic local density $\Theta_d^{\mathrm{mot}}(X,O) \in \mathbb{Q}\otimes\mathcal{M}_{\mathbb{C}}$.

Choose a good resolution $\pi\colon Y \to X$, factoring through the maximal- ideal blowup and Nash modification, with simple-normal-crossing exceptional divisor $D = \sum_{i\in J} m_i E_i$. Put

$$E_I^\circ = \bigcap_{i\in I} E_i \setminus \bigcup_{j\notin I} E_j, \qquad \alpha_i = \widehat{k}_i + 1 - dm_i,$$

where $\widehat{k}_i$ is the Mather discrepancy along E_i and $\mathbb{L} = [\mathbb{A}^1]$.

Main theorem (Theorems 1.1 and 1.2)**.** *For a d-dimensional definable set A, the generating series of normalized local volumes*

$$Z_A(T) = \sum_{n\geq 0} \frac{\mu_d(A\cap B(x,n))}{\mu_d(B^d(x,n))} T^n$$

is rational with at most a simple pole at $T=1$, and $\Theta_d^{\mathrm{mot}}(A,x) = \mathrm{Res}_{T=1} Z_A(T)$. For the isolated singularity above this becomes

$$\Theta_d^{\mathrm{mot}}(X,O) = \sum_{\substack{i\in J\\ \alpha_i=0}} \frac{1}{m_i} \sum_{\substack{I\subseteq J\\ i\in I}} \frac{(\mathbb{L}-1)^{|I|}[E_I^\circ]}{\mathbb{L}^d-1} \prod_{j\in I\setminus\{i\}} \frac{\mathbb{L}^{-\alpha_j}}{1-\mathbb{L}^{-\alpha_j}}.$$

Thus an invariant defined by all small balls is recovered from finite resolution data.

How the proof produces the formula

1. Residue extracts cyclic limits. Presburger cell decomposition expresses the coefficients of Z_A by polynomials in n times powers of $\mathbb{L}$. Negative-degree terms have zero residue; a degree-zero factor $(1-T^e)^{-1}$ contributes the average of the e congruence limits.

2. Resolve the valuative spheres. Proposition 2.10 permits replacing balls by valuative spheres $S(O,n)$ (exact distance n). Lemma 5.1 identifies the inverse image of $X\cap S(O,n)$ near E_I° with arcs whose contact orders satisfy $\sum_{i\in I} m_i\ell_i = n$. This converts a geometric shrinking condition at O into a linear constraint on exceptional contacts.

3. The Jacobian becomes Mather discrepancy. Change of variables pulls the volume to Y. For singular X, the image of $\pi^*\bigwedge^d \Omega_X$ in $\bigwedge^d \Omega_Y$ measures the Jacobian by the Mather discrepancy. Normalization makes contact order ℓ_i contribute $\mathbb{L}^{-\alpha_i\ell_i}$.

4. Factor and take the pole. Summing independently over positive contact orders on a fixed stratum gives

$$Z_I(T) = \prod_{i\in I} \frac{\mathbb{L}^{-\alpha_i} T^{m_i}}{1-\mathbb{L}^{-\alpha_i}T^{m_i}}.$$

At most one α_i vanishes on a nonempty stratum. Other factors are regular at $T=1$, while a zero exponent contributes $1/m_i$. Restoring $(\mathbb{L}-1)^{|I|}[E_I^\circ]/(\mathbb{L}^d-1)$ gives the theorem.

Toy case, hard point, and scope

If the exceptional divisor has a single smooth component E of multiplicity m, the formula reads

$$\Theta_d^{\mathrm{mot}}(X,O) = \begin{cases} \dfrac{(\mathbb{L}-1)[E]}{m(\mathbb{L}^d-1)}, & \widehat{k}_E + 1 = dm,\\ 0, & \widehat{k}_E + 1 \neq dm. \end{cases}$$

This isolates pole selection from stratum combinatorics. The hard step is controlling the singular Jacobian by Mather discrepancy while retaining only a simple pole. The answer lies in a localized Grothendieck ring, not in $\mathbb{Q}$. The theorem requires an isolated complex singularity and the stated resolution. For surfaces, $\alpha_i = m_i(q_i-1)$, so precisely the inner-rate-1 components contribute. This digest omits Section 6's broader ideal-sheaf Zeta function and its comparisons with existing motivic Zeta functions.

Codex mathematical paper digest, 2026-10-27. Source paper should be consulted for full statements and proofs.