

Rational curves on elliptic K3 surfaces

Salim Tayou

arXiv:1805.07975

<http://arxiv.org/abs/1805.07975v4>

Rational curves from a genus-one fibration

Let k be algebraically closed and let $\pi : X \rightarrow \mathbf{P}_k^1$ be an elliptic K3 surface: the generic fiber is a smooth integral genus-one curve, but a section is not assumed. The fibration is non-isotrivial when its smooth fibers are not all isomorphic. A rational curve here means an integral curve of geometric genus zero.

Main theorem (Theorem 1.1). *The surface X contains infinitely many rational curves in either of the following cases:*

1. *the elliptic fibration is non-isotrivial, in arbitrary characteristic;*
2. *$\text{char}(k) \neq 2, 3$, with no non-isotriviality assumption.*

The characteristic-zero cases were known; the paper's new work treats positive characteristic. The strategy is to manufacture rational multisections of unbounded degree. The subtlety is that a natural fiberwise multiplication map can identify points of a multisection, so one must either rule out that collapse or extract strong geometry from it.

Period, index, and divisible Brauer classes

Let $J(X) \rightarrow \mathbf{P}_k^1$ be the Jacobian fibration. Over the smooth locus, X is a torsor under $J(X)$, with class $[X]$ in its Tate–Shafarevich group, which is isomorphic to $\text{Br}(J(X))$. Define d_X as the positive generator of the degrees of multisections, equivalently the index of the generic genus-one curve. Lemma 2.4 says that

$$\text{ord}([X]) = d_X,$$

and supplies both a rational multisection of degree d_X and a generically étale multisection of that degree. Rationality follows from Bogomolov–Mumford: a minimal-degree effective divisor is a sum of rational curves, and exactly one component is horizontal.

After disposing of Picard rank at least 20 by known automorphism results, assume $\rho(X) \leq 19$ and work in positive characteristic p . Split $[X] = \alpha_p + \alpha$ into its p -primary and prime-to- p parts, with $\text{ord}(\alpha_p) = p^a$. For arbitrarily large primes $\ell \equiv 1 \pmod{p^a}$, divisibility and nontriviality of the prime-to- p part of $\text{Br}(J(X))$ yield a torsor X_ℓ with

$$J(X_\ell) \simeq J(X), \quad \ell[X_\ell] = [X], \quad d_{X_\ell} = \ell d_X.$$

Hence $J^\ell(X_\ell) \simeq X$, giving a rational fiberwise multiplication map $\eta_\ell : X_\ell \dashrightarrow X$. Lemma 2.4 provides a rational multisection $M_\ell \subset X_\ell$ of degree ℓd_X .

The monodromy contradiction

If η_ℓ is generically injective on M_ℓ , its image is a rational curve on X whose degree is divisible by ℓ ; letting ℓ vary gives infinitely many curves. Otherwise, on infinitely many smooth fibers there are distinct $P_1, P_2 \in M_\ell$ with $\ell(P_1 - P_2) = 0$. Thus M_ℓ becomes an irreducible component of the torsion correspondence

$$J(X)_{U,\text{prim}}[\ell] \times_U M_{\ell,U} \longrightarrow X_{\ell,U},$$

where the first factor parametrizes nonzero ℓ -torsion on the smooth fibers.

For a non-isotrivial fibration, Proposition 2.5 says that, for large ℓ , the mod- ℓ monodromy is all of $\text{SL}_2(\mathbf{F}_\ell)$. Its transitive action on nonzero torsion makes $J(X)_{U,\text{prim}}[\ell]$ irreducible of degree $\ell^2 - 1$ over U . Projection of the component above is then surjective, forcing

$$\deg(M_\ell) = \ell d_X \geq \ell^2 - 1,$$

contrary to the choice $\ell > d_X$ (and sufficiently large). Collapse is impossible, so the rational images have unbounded prime divisibility. This proves the non-isotrivial case in every characteristic.

The isotrivial alternative

Assume now that $\text{char}(k) \neq 2, 3$ and that the fibration is isotrivial. Monodromy then factors through the automorphism group of the geometric generic elliptic fiber, cyclic of order 2, 4, or 6. If η_ℓ collapses M_ℓ , the same torsion correspondence and a Riemann–Hurwitz computation give, for some constant $C > 0$,

$$g(M_\ell) \geq C c(J), \quad c(J) = \frac{1}{2}n_0 + n'_1 + n''_1 + \frac{5}{6}n_2 + \frac{3}{4}n_3 + \frac{2}{3}n_4 - 2,$$

where n_0 counts fibers of type I_0^* ; n'_1 and n''_1 count types I_a and I_a^* ($a > 0$); and n_2, n_3, n_4 count types II or II^* , III or III^* , and IV or IV^* , respectively. Since M_ℓ is rational, one gets $c(J) \leq 0$.

For $\text{char}(k) \neq 2, 3$, wild conductors vanish. Combining the Shioda–Tate formula with the ℓ -adic Euler formula for the singular fibers turns $c(J) \leq 0$ into

$$18 + r(X) + 3n'_1 + 2n''_1 + \frac{4}{3}n_2 + n_3 + \frac{2}{3}n_4 \leq \rho(X).$$

The resulting finite classification leaves either Picard rank 22 or a surface on Hassett’s list; in every case X is Kummer or has infinite automorphism group, and known results then give infinitely many rational curves. In characteristics 2 and 3, wild ramification spoils this tame numerical reduction, which is precisely why the theorem retains non-isotriviality there.

Model, significance, and scope

The smooth-fiber picture is the clearest toy model: multiplication by ℓ on an elliptic curve identifies two points exactly when their difference is ℓ -torsion. Full monodromy moves all $\ell^2 - 1$ nonzero torsion points in one orbit. A degree- ℓd_X rational multisection cannot carry that degree- $\ell^2 - 1$ orbit once ℓ is large. The global proof turns this elementary count into an irreducibility statement for a relative torsion divisor.

The decisive technical point is the dichotomy between injectivity, which immediately produces a new rational curve, and noninjectivity, which forces either an impossible monodromy degree or a rigid singular-fiber classification. The theorem does not settle isotrivial elliptic K3 surfaces in characteristics 2 and 3: wild conductor terms can be large, and the paper explicitly leaves the remaining small-rank configurations open.