# Canonical heights for correspondences

Patrick Ingram

arXiv:1411.1041

http://arxiv.org/abs/1411.1041v2

## From orbits to paths

Let $X$ be a projective variety over a number field $K$, and let $C \subset X \times X$ be a correspondence whose coordinate projections $x, y : C \to X$ are finite and surjective. Unlike a morphism, $C$ does not choose a unique successor. The paper therefore replaces $X$ by the inverse limit

$$\mathcal{P} = \varprojlim \mathcal{P}_n,$$

where $\mathcal{P}_n$ parametrizes paths of length $n$. It carries maps $\pi : \mathcal{P} \to X$ (initial vertex), $\varepsilon : \mathcal{P} \to C$ (first edge), and a finite surjective shift $\sigma : \mathcal{P} \to \mathcal{P}$, with $\pi = x\varepsilon$ and $\pi\sigma = y\varepsilon$. In general $\mathcal{P}$ is not of finite type, which is the source of both the novelty and the technical difficulty.

Assume that $C$ is polarized: for an ample Cartier $L \in \mathrm{Div}(X) \otimes \mathbb{R}$ and $\alpha > 1$, $y^*L \sim \alpha x^*L$.

**Main theorem** (Theorem 1.1)**.** *There is a function $\widehat{h}_{X,C,L} : \mathcal{P}(\overline{K}) \to \mathbb{R}$ such that*

$$\widehat{h}_{X,C,L}(\sigma P) = \alpha \widehat{h}_{X,C,L}(P), \qquad \widehat{h}_{X,C,L}(P) = h_{X,L}(\pi P) + O(1).$$

*It is explicitly*

$$\widehat{h}_{X,C,L}(P) = \lim_{n\to\infty} \alpha^{-n} h_{X,L}(\pi\sigma^n P).$$

*If $C$ is the graph of a polarized endomorphism, $\pi$ is an isomorphism and this is the Call–Silverman canonical height.*

The path, rather than merely its initial point, is essential: distinct choices of edges can have different heights. A finitely supported path has height zero, but a path that merely revisits one vertex need not. Corollary 1.3 nevertheless shows that initial vertices in $X(\overline{K})$ admitting a repetitive path have bounded $L$-height; hence only finitely many such initial vertices lie in $X(K)$.

## Ingredient 1: the height-machine error telescopes

The commutative path diagram and functoriality of Weil heights give, uniformly in $P$,

$$\left|\alpha^{-1} h_{X,L}(\pi\sigma P) - h_{X,L}(\pi P)\right| \le \kappa.$$

Applying this after $m$ shifts and summing the resulting geometric series shows that $\alpha^{-n} h_{X,L}(\pi\sigma^n P)$ is Cauchy. The same estimate bounds its limit from $h_{X,L}(\pi P)$; shifting the limit proves the functional equation. This is Tate's construction, but the finite-type height machine is used only on $\pi(P) \in X$ and $\varepsilon(P) \in C$, never directly on the provariety $\mathcal{P}$.

Compactness of each inverse-limit fibre $\pi^{-1}(a)$ in the tree topology, together with the exponentially decaying tail estimate, also proves continuity (Proposition 3.3). Consequently the maximum and minimum path heights above a fixed vertex are attained.

## Ingredient 2: specialization is made uniform by shifting

For a family over a curve $U/K$, polarization means that $y^*L - \alpha x^*L$ is fibral. Theorem 4.1 treats paths over the inverse limit of finite covers of $U$, allowing all vertices to be algebraic over $K(U)$ while requiring the initial vertex to descend to a finite cover. If $h_U$ has degree one, then

$$\widehat{h}_t(P_t) = \big(\widehat{h}_\eta(P_\eta) + o(1)\big)h_U(t) \quad (h_U(t) \to \infty),$$

with the paper's more general notation allowing a dominant base change. The proof first obtains only an $O(h_U(t))$ discrepancy: a fibral divisor controls the canonical-versus-naive height, while a degree computation on the finite cover controls the initial vertex up to $O(\sqrt{h_U(t)})$. Replacing $P$ by $\sigma^n P$ multiplies the uniform error by $\alpha^{-n}$; letting $n \to \infty$ forces the normalized error to vanish. Thus positive generic height makes finitely supported specializations a bounded-height set (and, in the stated $K$-rational corollary, a finite set).

## Ingredient 3: local heights over all algebraic places

A path in $\mathcal{P}(\overline{K})$ need not be defined over any number field, so an ordinary finite sum over places is unavailable. Gubler's measured space $M_{\overline{K}}$ replaces it. The local height error is bounded, continuous in the path, measurable in the place, and supported over a finite-measure set. Lemma 6.2 corrects it by the fixed point of the contraction

$$S(\delta)(P, v) = \alpha^{-1}\big(\delta(\sigma P, v) - \gamma(P, v)\big).$$

The corrected local height is integrable, satisfies the local scaling equation, and the product formula shows

$$\widehat{h}_{X,C,L}(P) = \int_{M_{\overline{K}}} \widehat{\lambda}_{X,C,L}(P, v)\, d\mu(v).$$

The hardest point is not the contraction itself, but extending its error across divisor supports on a non-finite-type path space while retaining measurability and integrability.

## A toy correspondence

For $C : y^2 = x^3 + 1$ on $\mathbb{P}^1$, the polarization factor is $\alpha = 3/2$. The paper checks that a path beginning

$$P : 0 \longrightarrow 1 \longrightarrow 2^{1/2} \longrightarrow (2^{3/2} + 1)^{1/2} \longrightarrow \cdots$$

has positive canonical height. Prepending the edges $0 \to -1 \to 0$ gives a repetitive path $Q$ with $\sigma^2 Q = P$, hence $\widehat{h}(Q) = \alpha^{-2}\widehat{h}(P) = \frac{4}{9}\widehat{h}(P) > 0$. This cleanly separates revisiting a vertex from finite support. Conversely, $y^2 = x^3$ has nonrepeating paths through roots of unity with height zero: branching breaks both deterministic implications.

## Scope and a source-level caveat

The construction does not characterize every height-zero path: unbounded algebraic degree permits nonrepeating examples, so the deterministic equivalence *height zero if and only if preperiodic* fails. Also, the displayed sign in the introduction's Theorem 1.8 differs from the detailed Theorem 6.1 and its proof. For $\operatorname{div}(f) = y^*L - \alpha x^*L$, pages 21–24 use $\widehat{\lambda}(\sigma P, v) = \alpha\widehat{\lambda}(P, v) - \log|f\varepsilon(P)|_v$; the digest follows that proved convention.