A note on the Bloch-Beilinson conjecture for K3 surfaces and spherical objects

Daniel Huybrechts

arXiv:1009.4371v1 (2010), 8 pages
http://arxiv.org/abs/1009.4371v1

The question

Let X be a projective K3 surface over an algebraically closed field k, and write $D^b(X) = D^b(\operatorname{Coh} X)$. An object A is *spherical* if

$$\operatorname{Ext}^i(A,A) \simeq \begin{cases} k, & i = 0, 2, \\ 0, & \text{otherwise.} \end{cases}$$

Let $\mathcal{S}^* \subset D^b(X)$ be the smallest strictly full triangulated subcategory containing all spherical objects. Crucially, its definition does not close under direct summands, so density need not imply equality with $D^b(X)$.

Main theorem (Theorems 0.1 and 3.1)**.** *Let X be a projective K3 surface over $\overline{\mathbb{Q}}$ with Picard number $\rho(X) \geq 2$. The following are equivalent:*

(i) $\mathrm{CH}^2(X) \simeq \mathbb{Z}$*;*
(ii) $D^b(X) = \mathcal{S}^*$*, so spherical objects generate without taking direct summands;*
(iii) $\mathcal{S}^*$ *admits a bounded t-structure.*

Condition (i) is the Bloch–Beilinson prediction for zero-cycles on a K3 surface over $\overline{\mathbb{Q}}$. The theorem turns that cycle-theoretic statement into both a categorical generation criterion and a structural criterion internal to $\mathcal{S}^*$.

Proof mechanism

1. Spherical objects give a dense subcategory. Every line bundle on a K3 surface is spherical. A finite portion of a resolution by sufficiently negative line bundles gives, for each coherent sheaf E, a complex $K^\bullet$ of line bundles with $K^\bullet \simeq E \oplus F[2]$ for some F. Passing from sheaves to bounded complexes shows that $\mathcal{S}^*$ is dense: every object becomes an object of $\mathcal{S}^*$ after adding a suitable summand.

2. Thomason detects membership by K-theory. For an essentially small triangulated category $\mathcal{D}$, Thomason's theorem classifies strictly full dense triangulated subcategories by subgroups of $K(\mathcal{D})$. Applied here, it says

$$\mathcal{S}^* = \{E \in D^b(X) : [E] \in K(\mathcal{S}^*)\}.$$

Thus the apparent problem of constructing every object by cones is replaced by the computation of one subgroup of the Grothendieck group.

3. The subgroup is the Beauville–Voisin ring. The main imported input (Theorem 1.8, from Huybrechts's earlier work) is

$$K(\mathcal{S}^*) = R(X) \qquad (\rho(X) \geq 2),$$

where $R(X) \subset \mathrm{CH}^*(X)$ is generated by Chern characters of line bundles and the distinguished zero-cycle c_X with $24c_X = c_2(X)$. The Beauville–Voisin cycle map identifies $R(X) \simeq \mathbb{Z} \oplus \mathrm{NS}(X) \oplus \mathbb{Z}$.

If $D^b(X) = \mathcal{S}^*$, equality of Grothendieck groups gives $\mathrm{CH}^*(X) = R(X)$ and hence $\mathrm{CH}^2(X) \simeq \mathbb{Z}$. Conversely, if $\mathrm{CH}^2(X) \simeq \mathbb{Z}$, then $\mathrm{CH}^*(X) = R(X) = K(\mathcal{S}^*)$. Every $E \in D^b(X)$ therefore has $[E] \in K(\mathcal{S}^*)$, so Thomason's criterion puts E itself in $\mathcal{S}^*$.

4. A bounded t-structure forces equality. A triangulated category with a bounded t-structure is Karoubian. Since $D^b(X)$ is precisely the idempotent completion of the dense category $\mathcal{S}^*$, the latter is Karoubian exactly when $\mathcal{S}^* = D^b(X)$. Hence (iii) implies (ii), while equality with the usual derived category supplies the converse.

A diagnostic example and caveats

Thomason's description immediately places $E \oplus E[1]$ in $\mathcal{S}^*$ for every E, because its K-class is zero. This is a useful toy model for the distinction between density and actual generation: it says nothing by itself about either summand. The hard mathematical input is the exact identification $K(\mathcal{S}^*) = R(X)$, not the final two-line equivalence. For $\rho(X) = 1$, the proof still gives $\mathrm{CH}^2(X) \simeq \mathbb{Z} \Rightarrow D^b(X) = \mathcal{S}^*$, but the reverse implication is not obtained. Over an algebraically closed extension of $\overline{\mathbb{Q}}$ of positive transcendence degree, Mumford's theorem forces $\mathrm{CH}^2(X) \neq \mathbb{Z}$, so none of the three conditions holds.

Codex mathematical paper digest, 2026-10-26. Source paper should be consulted for full statements and proofs.