The pseudo-index of horospherical Fano varieties

Boris Pasquier

Boris Pasquier, arXiv:0802.0612v1 (2008), 9 pages

http://arxiv.org/abs/0802.0612v1

A sharp Mukai inequality beyond the toric case

For a normal complex projective Fano variety X, so that $-K_X$ is Cartier and ample, its pseudo-index is

$$\iota_X = \min\{-K_X \cdot C : C \subset X \text{ a rational curve}\}.$$

A horospherical variety has an action of a connected reductive group G with open orbit G/H, where G/H is a torus bundle over a flag variety. This class contains both toric and flag varieties.

Main theorem (Pasquier, Theorem 1)**.** *Let X be a $\mathbb{Q}$-factorial horospherical Fano variety of dimension d, Picard number ρ_X, and pseudo-index ι_X. Then*

$$(\iota_X - 1)\rho_X \leq d.$$

Equality holds if and only if

$$X \simeq \left(\mathbf{P}^{\iota_X - 1}\right)^{\rho_X}.$$

The theorem proves the generalized Mukai inequality in this possibly singular $\mathbb{Q}$-factorial class and shows that equality itself forces the familiar smooth product.

Ingredient 1: encode curves by a weighted reflexive polytope

Fix the open orbit G/H. Its character lattice M and dual N have rank n, and the embedding X corresponds to a G/H-reflexive polytope $Q \subset N_{\mathbb{R}}$. The root data contribute distinguished "color" points and integers $a_\alpha = \langle 2\rho^P, \check{\alpha}\rangle \geq 2$; attach to each vertex u a weight a_u (equal to a_α at a color vertex and to 1 otherwise). If Q has $n + r$ vertices, then

$$\rho_X = r + |S \setminus I| - |D_X|.$$

For $v \in V(Q^*)$, let $F_v = \{u \in Q : \langle v, u\rangle = -1\}$ and define the global minimum

$$\varepsilon_Q = \min_{\substack{u \in V(Q),\ v \in V(Q^*) \\ u \notin F_v}} a_u(1 + \langle v, u\rangle).$$

The crucial bridge is Proposition 5:

$$\iota_X \leq \varepsilon_Q, \qquad \iota_X \leq a_\alpha \quad (\alpha \in (S \setminus I) \setminus D_X).$$

Lemma 6 says that for a minimizing pair (u, v), the vertex u is adjacent to F_v. The corresponding wall curve C_μ, together with Schubert curves $C_{\alpha,v}$ in closed G-orbits, generates the effective curve cone. The intersection formulas of Lemma 7 give

$$-K_X \cdot C_\mu = \frac{a_u(1 + \langle v, u\rangle)}{\langle \chi_\mu, a_u u\rangle}, \qquad -K_X \cdot C_{\alpha,v} = a_\alpha + \langle v, \check{\alpha}_M\rangle.$$

The first denominator is a positive integer and the second correction is nonpositive for a suitable v, yielding the two bounds above. This is where geometric positivity becomes finite polytope arithmetic.

Ingredient 2: count incidences with a positive relation

Because 0 lies inside Q^*, choose vertices v_i and positive integers m_i with $\sum_i m_i v_i = 0$. For each vertex u of Q, let $A(u) = \{i : \langle v_i, u\rangle = -1\}$. Comparing the terms inside and outside $A(u)$ with the definition of ε_Q, then summing over u, gives

$$\varepsilon_Q r \le \sum_{u \in V(Q)} a_u.$$

The $\mathbb{Q}$-factoriality hypothesis implies that Q is simplicial, which makes every facet contribute exactly n incidences. Substituting the Picard-number formula and the two bounds from Proposition 5 yields

$$\rho_X(\iota_X - 1) \le \sum_{\alpha \in S \setminus I} (a_\alpha - 1) + n \le d.$$

The final root-theoretic estimate is the cited Lemma 4.13 from the author's earlier work.

For the color-free toric toy case, $n = d$, every $a_u = 1$, and $\rho_X = r$. The central count reduces to $\iota_X r \le d + r$, or immediately $\rho_X(\iota_X - 1) \le d$. Horospherical colors are precisely the extra terms whose root weights must be absorbed by the last inequality.

Ingredient 3: rigidity when every inequality is sharp

Equality forces four simultaneous conditions: every relevant weighted pairing is either 0 or ε_Q; $\iota_X = \varepsilon_Q$; every non-color root attains $\iota_X = a_\alpha$; and the root-theoretic dimension bound is sharp. Lemma 9 shows that $\iota_X = \varepsilon_Q$ upgrades $\mathbb{Q}$-factoriality to local factoriality. The remaining equalities restrict each active Dynkin component to type A or C with its marked root at a simple end; the horospherical smoothness criterion then makes X smooth.

The equality proof first splits off the roots whose restricted coroot is zero, producing factors $\mathbf{P}^{\iota_X - 1}$. For the remaining polytope, sharpness partitions the character lattice and Q into components of Picard number one. The cited classification of those smooth two-orbit components identifies every component as another $\mathbf{P}^{\iota_X - 1}$. This equality analysis, rather than the numerical count, is the technically deepest part.

The argument relies on the Luna–Vust/polytope classification and two results from the author's earlier work; it is not a self-contained proof for arbitrary Fano varieties. Its scope is complex, horospherical, Fano, and $\mathbb{Q}$-factorial, with $-K_X$ Cartier as stipulated in the paper's definition of Fano.