

Families of Calabi-Yau manifolds and canonical singularities

Valentino Tosatti

arXiv:1311.4845

<http://arxiv.org/abs/1311.4845v3>

Finite distance detects a canonical filling

Let $\pi: \mathcal{X} \rightarrow \Delta$ be a flat projective family of n -folds, smooth over the punctured disc $\Delta^\times$, with $K_{X_t} \cong \mathcal{O}_{X_t}$ for $t \neq 0$, and assume all fibers are connected. Fix a line bundle L on $\mathcal{X}$ whose restrictions L_t are ample, and let ω_t be the unique Ricci-flat Kähler metric in $c_1(L_t)$. A local holomorphic volume form Ω_t determines the Weil–Petersson form by

$$\omega_{\text{WP}} = -\sqrt{-1} \partial \bar{\partial} \log \left((-1)^{n^2/2} \int_{X_t} \Omega_t \wedge \bar{\Omega}_t \right). \quad (1)$$

Main theorem (Tosatti, Theorem 1.2). *The puncture 0 is at finite Weil–Petersson distance if and only if, after a finite base change, the family is birational to a family whose central fiber is a Calabi–Yau variety with at worst canonical singularities: it is normal and projective, and its canonical divisor is Cartier and trivial.*

Thus an analytic condition on the variation of smooth Ricci-flat fibers is exactly the condition that the degeneration admit the natural algebro-geometric Calabi–Yau filling.

The bridge theorem on a minimal model

Suppose now that $\mathcal{X}$ is normal, $K_{\mathcal{X}/\Delta} \cong \mathcal{O}_{\mathcal{X}}$, and $(\mathcal{X}, X_0)$ is dlt; the paper calls such a family relatively minimal. Choose a nowhere-vanishing section $\Omega \in H^0(\mathcal{X}, K_{\mathcal{X}/\Delta})$ and set $\Omega_t = \Omega|_{X_t}$, including Ω_0 on $X_{0,\text{reg}}$. Theorem 1.1 proves that the following are equivalent:

1. X_0 has canonical singularities;
2. some irreducible component $D \subset X_0$ has a resolution $\tilde{D}$ with $H^0(\tilde{D}, K_{\tilde{D}}) \neq 0$;
3. the L^2 masses $(-1)^{n^2/2} \int_{X_t} \Omega_t \wedge \bar{\Omega}_t$ are uniformly bounded above.

These conditions also imply uniform diameter bounds for the Ricci-flat metrics $\omega_t \in c_1(L|_{X_t})$, equivalently the uniform non-collapsing estimate

$$\text{Vol}_{\omega_t} B_{\omega_t}(x, r) \geq C^{-1} r^{2n} \quad (0 < r \leq \text{diam}(X_t, \omega_t)). \quad (2)$$

Ingredient I: an integrability test for discrepancies

The subtle implication is from bounded mass to canonical singularities. The bounded-mass condition above gives the finite central integral

$$(-1)^{n^2/2} \int_{X_{0,\text{reg}}} \Omega_0 \wedge \bar{\Omega}_0 < \infty. \quad (3)$$

The dlt central fiber is reduced with normal components. Because X_0 is connected, reducibility would force two distinct components to meet. Adjunction then makes Ω_0 have at least a simple pole along their intersection, contradicting (3). Hence X_0 is irreducible and normal. Finite mass is then the standard analytic criterion for log-terminal singularities. On a resolution

$$K_{\tilde{X}_0} \sim_{\mathbb{Q}} f^* K_{X_0} + \sum_E a_E E,$$

one has $a_E > -1$. Because K_{X_0} is Cartier, every a_E is an integer, so $a_E \geq 0$: log terminal upgrades to canonical.

The component criterion gives a second route. If a component carrying a top form met another component, adjunction would make its canonical class a nontrivial negative boundary modulo exceptional divisors, so it could not be effective on a resolution. The fiber is therefore irreducible, and effectivity of the resolved canonical class again gives nonnegative discrepancies.

Ingredient II: the metric consequences

Ricci-flatness is the Monge–Ampère identity

$$\omega_t^n = e^{c_t} (-1)^{n^2/2} \Omega_t \wedge \overline{\Omega}_t. \quad (4)$$

The polarized volume $\int_{X_t} \omega_t^n$ is constant, so the mass bound is equivalent to a uniform lower bound in (3). A theorem of Rong–Zhang then gives the diameter bound. Bishop–Gromov comparison turns bounded diameter and constant total volume into (2); conversely, many disjoint unit balls along a very long minimizing geodesic would violate constant total volume, so (2) implies bounded diameter.

Ingredient III: reach the minimal model

Finite Weil–Petersson distance is unchanged by finite base change and by a birational modification that is an isomorphism over $\Delta^\times$: the two volume forms differ by a nowhere-vanishing function of t , whose logarithm contributes no curvature to (1). Semistable reduction makes X_0 a reduced simple-normal-crossings divisor. The relative minimal model program then produces

$$\pi': \mathcal{X}' \rightarrow \Delta, \quad (\mathcal{X}', X'_0) \text{ dlt}, \quad K_{\mathcal{X}'/\Delta} \sim_{\mathbb{Q}} 0.$$

One must strengthen the last relation. Pushing forward $\mathcal{O}_{\mathcal{X}'}(K_{\mathcal{X}'/\Delta})$ gives a rank-one torsion-free sheaf on the disc, hence a line bundle M . The natural map $\pi'^* M \rightarrow \mathcal{O}_{\mathcal{X}'}(K_{\mathcal{X}'/\Delta})$ differs by a $\mathbb{Q}$ -Cartier divisor D supported on X'_0 . Relative nefness and the negativity lemma force $D = rX'_0$; reducedness then upgrades the relative canonical divisor to Cartier and relatively trivial. The bridge theorem now applies.

Ingredient IV: a top form survives the MMP

For the semistable model, Wang’s criterion says that finite Weil–Petersson distance is equivalent to the existence of a component $D_1 \subset X_0$ with $H^0(D_1, K_{D_1}) \neq 0$. Its transform cannot be contracted during the minimal model program: a contracted component would be uniruled, incompatible with a nonzero top form. Thus a component $D'_1 \subset X'_0$ remains birational to D_1 . Birational invariance of $h^{n,0}$ transfers the top form to a resolution of D'_1 , which is condition (2) of Theorem 1.1 and forces X'_0 to be canonical.

Local model and scope

The local picture behind the irreducibility step is a crossing $X_0 = D_1 \cup D_2$: restriction of the relative volume form to D_1 has a logarithmic pole along $D_1 \cap D_2$, and the square norm of dz/z has a logarithmically divergent integral. This explains why bounded Weil–Petersson mass rules out a reducible dlt central fiber.

The theorem asserts existence only after finite base change and birational modification. The paper proves that canonical central fiber implies non-collapsing, but does not prove the converse inside Theorem 1.1. Its technical bottleneck is transporting Wang’s top-form component through the relative MMP while simultaneously upgrading a merely $\mathbb{Q}$ -trivial relative canonical divisor to a genuinely trivial one.