

Multidegrees of monomial rational maps

Paolo Aluffi

arXiv:1308.4152

<http://arxiv.org/abs/1308.4152v1>

A convex formula for a rational map

Let $V \subset \mathbb{P}^r$ be a projective variety. For a rational map $\varphi: V \dashrightarrow \mathbb{P}^N$, let Γ be the closure of its graph, and write h, H for the hyperplane classes pulled back from the two factors. Its multidegrees are

$$\gamma_\ell = h^{\dim V - \ell} H^\ell \cdot [\Gamma], \quad \gamma_\varphi(t) = \sum_{\ell \geq 0} \gamma_\ell t^\ell.$$

They record, among other things, the dimension and degree of the image and the generic degree of φ .

Assume that φ is defined by monomials $\mu_i = s_1^{m_{i1}} \cdots s_n^{m_{in}}$ in sections of line bundles $\mathcal{L}_j$, all lying in one line bundle $\mathcal{L}$. Put $X_j = V(s_j)$. The paper requires that monomial schemes in the X_j can be principalized by codimension-two monomial blowups; it is enough that V is smooth and $\sum X_j$ has simple normal crossings. After subtracting one exponent vector from all the others, let $N_\varphi \subset \mathbb{R}^n$ be the convex hull of their translates by the positive orthant.

Main theorem (Aluffi, Theorems 1.1 and 2.3). *The multidegree class G on V is*

$$G = \int_{N_\varphi} \frac{n! X_1 \cdots X_n da_1 \cdots da_n}{(1 + a_1 X_1 + \cdots + a_n X_n)^{n+1}}. \quad (1)$$

Equivalently, after treating h, t, X_j as parameters and then evaluating the X_j as divisor classes,

$$\gamma_\varphi(t) = \int_{N_\varphi} \frac{n! X_1 \cdots X_n t^n h^{\dim V + 1} da_1 \cdots da_n}{(h + (a_1 X_1 + \cdots + a_n X_n)t)^{n+1}}. \quad (2)$$

The result is independent of the exponent vector chosen as pivot.

This extends familiar toric volume formulas to monomials in sections on a possibly non-toric projective variety.

Ingredient I: the graph is controlled by the base scheme

Let $S \hookrightarrow V$ be the base scheme. If $\pi: \Gamma \rightarrow V$, the “shadow”

$$G = \pi_*((1 + H + H^2 + \cdots) \cap [\Gamma])$$

has codimension- ℓ part whose degree is γ_ℓ . Lemma 2.1 rewrites it entirely in terms of the Segre class:

$$G = c(\mathcal{L}^*)^{-1} \cap ([V] - i_* s(S, V)) \otimes_V \mathcal{L}^*. \quad (3)$$

Thus the indeterminacy scheme, rather than a resolution of the graph, is the bridge from intersection theory to convex geometry.

Ingredient II: translate the Segre integral

Let N be the complement, in the positive orthant, of the convex hull of the untranslated exponent orthants. The monomial Segre-class formula used in Lemma 2.2 says that $[V] - i_*s(S, V)$ is the integral in (1) over the complement N^c . Applying $\otimes_V \mathcal{L}^*$ to the integrand subtracts the pivot exponents, because $\sum_j m_j X_j = c_1(\mathcal{L})$. It therefore translates N^c exactly to N_φ and proves (1). Inserting a bookkeeping variable for codimension and intersecting with powers of h gives (2).

Ingredient III: integrals become simplex volumes

Triangulate N_φ by generalized simplices whose vertices may include positive coordinate directions at infinity. For a rank- s simplex T , Lemma 3.1 evaluates its integral explicitly. Every finite vertex v is a difference of two exponent vectors, hence $\sum_j v_j X_j = 0$ in the Chow group. All denominators contributed by finite vertices disappear, leaving

$$G = \sum_T \widehat{\text{Vol}}(T) X_T, \quad \gamma_\varphi(t) = \sum_T \widehat{\text{Vol}}(T) \deg(T) t^{\text{rk } T}. \quad (4)$$

Here X_T is the product of those X_j whose coordinate direction is not an infinite vertex, and $\deg(T) = h^{\dim V - \text{rk } T} X_T$. This is Theorem 1.4: every multidegree is a sum of normalized projected volumes weighted by intersection numbers on V .

Ingredient IV: a characteristic polynomial shortcut

For a monomial morphism of the $(n-1)$ -torus with exponent matrix $A \in M_{n-1}(\mathbb{Z})$, homogenization gives $\varphi: \mathbb{P}^{n-1} \dashrightarrow \mathbb{P}^{n-1}$. The map is called *well-presented* when every coordinate projection of its Newton outer region is generated by the complementary projected rows and the corresponding complementary minors have the prescribed alternating sign (or vanish). These conditions decompose the complementary Newton region into one generalized simplex for each proper coordinate subset. Their volumes are precisely the signed complementary minors.

The determinant expansion packages those minors. After removing the redundant homogeneous direction, Lemma 4.6 identifies the remaining factor with the characteristic polynomial of A . Theorem 4.10 gives

$$\boxed{\gamma_\varphi(t) = t^{n-1} \det(t^{-1}I - A)}. \quad (5)$$

For the standard Cremona involution, $A = -I$, so $\gamma_\varphi(t) = (1+t)^{n-1}$: the binomial multidegrees arise from one linear-algebra calculation.

Scope and bottleneck

Formula (1) depends on the stated principalization hypothesis, and the characteristic-polynomial shortcut (5) is only proved for well-presented maps; the paper does not characterize all such maps. Nor does it analyze degree growth under iteration. The decisive step is the exact conversion of the graph shadow to a monomial Segre class and then to the translated Newton region; the later volume and determinant formulas work because that translation makes every finite-vertex linear form vanish as a divisor class.