Portraits of preperiodic points for rational maps

Dragos Ghioca, Khoa Nguyen, Thomas J. Tucker

arXiv:1407.1573

http://arxiv.org/abs/1407.1573v1

Problem and setup

Let K/k be a one-variable function field, with k algebraically closed of characteristic 0, and let $\varphi \in K(z)$ have degree $d > 1$. A point has *portrait* (m, n) if its first periodic iterate is $\varphi^m(\alpha)$ and that point has minimum period n. The question is whether a prescribed portrait occurs after reducing (φ, α) at some place of K, with minimum period exactly n, not merely dividing n.

Two genuine obstructions must be removed. Write $X(\varphi)$ for the set of n such that φ is totally ramified at every point of minimum period n, and $Y(\varphi, \alpha)$ for the positive m such that φ is totally ramified over $\varphi^m(\alpha)$, equivalently at $\varphi^{m-1}(\alpha)$. Let $\widehat{h}_\varphi$ be the canonical height.

Main theorem (Theorem 1.3)**.** *Fix $\tau, s > 0$, and a set S of at most s places containing every place of bad reduction. There is an effectively computable finite set $Z(\tau, s) \subset \mathbb{Z}_{\geq 0} \times \mathbb{N}$, depending only on K, φ, τ, s, such that for every $\alpha \in \mathbb{P}^1(K)$, if*

$$\widehat{h}_\varphi(\alpha) > \tau, \qquad (m, n) \notin Z(\tau, s), \qquad m \notin Y(\varphi, \alpha), \qquad n \notin X(\varphi),$$

then the proof produces a good-reduction place $\mathfrak{p} \notin S$ at which α has portrait exactly (m, n).

Thus the exceptional set is uniform in α once its canonical height is bounded below. The ramification exclusions are sharp: total ramification can force a missing inverse branch, and hence prevent the requested tail or cycle.

Why primitive collisions exist

The arithmetic engine is Proposition 4.15. For a squarefree monic polynomial $F \in K[z]$ of degree at least 3, the function-field *abc* theorem gives a lower bound of the shape

$$\#\{\mathfrak{p} : v_\mathfrak{p}(F(\gamma)) > 0\} \;\geq\; h_K(\gamma) - O_F(1).$$

Its proof passes to the splitting field, sends three roots to $0, 1, \infty$, applies Mason–Stothers, and controls the ramified places by Riemann–Hurwitz. Choosing the roots of F among suitable inverse images of an orbit point therefore supplies many primes at which two orbit levels collide.

Write $r_\mathfrak{p}$ for reduction at $\mathfrak{p}$. For large m, n, take the $d^2 - 1 \geq 3$ unramified alternatives $\eta_i \neq \varphi^{m-2}(\alpha)$ satisfying $\varphi^2(\eta_i) = \varphi^m(\alpha)$, and put $F = \prod_i (z - \eta_i)$. After discarding the finite bad-reduction and inverse-root pole set, a prime dividing $F(\varphi^{m+n-2}(\alpha))$ yields

$$r_\mathfrak{p}(\varphi^{m+n}(\alpha)) = r_\mathfrak{p}(\varphi^m(\alpha)),$$

so the reduced period divides n. The *abc* count is of order $\widehat{h}_\varphi(\alpha) d^{m+n-2}$.

Removing false portraits

Two much smaller bad sets turn divisibility into an exact portrait. First, primes for which $\varphi^{m-1}(\alpha)$ is already on the cycle are bounded by heights of the finitely many inverse branches. Second, primes for which the period is n/q for some prime $q \mid n$ contribute at most a sum of heights of $\varphi^{m+n/q}(\alpha) - \varphi^m(\alpha)$, of order $(\log n) d^{m+n/2}$. Both are dominated by the d^{m+n}-scale supply. Outside them, $\varphi^m(\alpha)$ has minimum period n, while $\varphi^{m-1}(\alpha)$ is nonperiodic; Lemma 4.22 then gives preperiod exactly m.

The remaining strips are handled separately. For fixed small $n \notin X(\varphi)$, choose a period-n point with a nonperiodic inverse and count deep preimages. For fixed small $m \notin Y(\varphi, \alpha)$, use several depth-five inverse branches together with Proposition 4.1, which bounds primes where a desired collision is contaminated by an earlier one. These arguments leave only a finite rectangle, producing $Z(\tau, s)$.

Interpretation, application, and scope

A useful toy picture is that a divisor of $\varphi^{m+n}(\alpha) - \varphi^m(\alpha)$ creates a cycle modulo $\mathfrak{p}$; the proof's substance is to manufacture many such divisors and discard those already visible at an earlier tail level or a proper divisor of n. Unlike this affine picture, the actual argument uses chordal distance on $\mathbb{P}^1$, splitting fields, ramification, and canonical heights.

Iterating the theorem over coefficient function fields gives Theorem 2.1: for distinct constants $c_0, \ldots, c_{d-2}$, nested cofinite choices of portraits can be realized simultaneously by a monic degree-d polynomial with zero z^{d-1}-coefficient. The characteristic-zero hypothesis is essential to the proof through function-field *abc*; the paper does not establish the analogous number-field theorem unconditionally.

Codex mathematical paper digest, 2026-11-03. Source paper should be consulted for full statements and proofs.