

The Dynamical Andre-Oort Conjecture: Unicritical Polynomials

Dragos Ghioca, Holly Krieger, Khoa Nguyen, Hexi Ye

arXiv:1505.01447

<http://arxiv.org/abs/1505.01447v1>

Which curves contain many simultaneous PCF parameters?

Fix $d \geq 2$ and write $f_c(z) = z^d + c$. Its only finite critical point is 0, so f_c is postcritically finite (PCF) exactly when 0 is preperiodic. Let $\mathcal{M}_d$ be the generalized Mandelbrot set: the parameters for which the orbit of 0 is bounded.

Main theorem (Ghioca–Krieger–Nguyen–Ye, Theorem 1.1). *Let $C \subset \mathbb{A}_{\mathbb{C}}^2$ be an irreducible plane curve. There are infinitely many $(a, b) \in C$ for which both f_a and f_b are PCF if and only if one of the following holds:*

- (1) $C = \{t_0\} \times \mathbb{A}^1$ for a PCF parameter t_0 ;
- (2) $C = \mathbb{A}^1 \times \{t_0\}$ for a PCF parameter t_0 ;
- (3) C is $y = \zeta x$, where $\zeta^{d-1} = 1$.

Thus the dynamical analogue of a special curve is completely rigid in the two-dimensional unicritical parameter space. The reverse direction in (3) is the conjugacy $\zeta^{-1}f_{\zeta c}(\zeta z) = f_c(z)$; the work is to prove that no other curve projecting dominantly to both axes can contain infinitely many simultaneous PCF points.

Ingredient I: zero height fixes the bifurcation measure

Let X be a smooth projective model of C , and let $\psi_i: X \rightarrow \mathbb{P}^1$ be the two coordinate maps, of degrees d_i . More generally, a nonconstant ψ defines the family $z^d + \psi(t)$. On $L = \psi^* \mathcal{O}_{\mathbb{P}^1}(1)$ the paper constructs semipositive adelic metrics from the iterates $f_{\psi(t)}^n(0)$. Uniform convergence of the local escape-rate metrics (Proposition 3.3) yields a height satisfying

$$\widehat{h}_{\psi}(t) = \frac{d}{d_{\psi}} \widehat{h}_{f_{\psi(t)}}(0) \quad (1)$$

away from the poles of ψ (Proposition 3.5). Hence its zero set is precisely the PCF parameters, together with those poles.

Yuan’s small-point theorem then equidistributes any nonrepeating PCF sequence against the normalized bifurcation measure (Theorems 3.1 and 3.4). A simultaneous PCF sequence is small for both ψ_1 and ψ_2 ; uniqueness of its limiting measure gives

$$d_2 \mu_{\psi_1} = d_1 \mu_{\psi_2}, \quad d_2 G_{\psi_1} = d_1 G_{\psi_2}. \quad (2)$$

The second identity is Theorem 4.1: the difference of the two potentials is harmonic and vanishes on a stable component lying over the interior of $\mathcal{M}_d$. In particular the pole sets agree, and $\psi_1(t)$ escapes $\mathcal{M}_d$ exactly when $\psi_2(t)$ does.

Ingredient II: potentials become an analytic correspondence

The Böttcher uniformization

$$\Phi: \mathbb{C} \setminus \mathcal{M}_d \xrightarrow{\sim} \mathbb{C} \setminus \overline{\mathbb{D}}, \quad \Phi(c) = c + O(1),$$

satisfies $\log |\Phi(c)| = dG(c)$. On an unbounded component of the preimage of $\mathbb{C} \setminus \mathcal{M}_d$, equation (2) and the open mapping theorem therefore strengthen equality of absolute values to

$$\Phi(\psi_1(t))^{d_2} = e^{2\pi i \eta} \Phi(\psi_2(t))^{d_1}. \quad (3)$$

Proposition 5.6 proves $\eta \in \mathbb{Q}$. If a suitable simultaneous PCF point is Misiurewicz, this follows from the rational external rays landing at its two coordinates. Otherwise one uses centers of hyperbolic components and a neighboring parabolic root, avoiding the finite branch locus of the projections. Thus a local analytic branch $h = \psi_1 \circ \psi_2^{-1}$ acts affinely on external angles.

Ingredient III: wake widths force a rotation

The remaining problem is to classify the local correspondences from (3). Rational ray pairs cut out wakes of hyperbolic components. If H has period N and width $|H|$, the wake formula gives the width of its p/q -subwake as

$$|w_{p/q}| = \frac{|H|(d^N - 1)^2}{(d - 1)(d^q - 1)}. \quad (4)$$

Theorem 5.7 compares (4) before and after applying h . Maximal $1/2$ -subwakes first show that h preserves the period of every hyperbolic component in a carefully chosen region (Proposition 5.8). Denominators of the root rays then force the angular slope to have the form $r/(d - 1)$ for a new integer $1 \leq r \leq d - 1$, and its translation to have denominator dividing $d - 1$. Applying the same restriction to the rays of angle $1/(d^N - 1)$ forces $r = d - 1$. Consequently

$$h(z) = \zeta z, \quad \zeta^{d-1} = 1, \quad (5)$$

first off $\mathcal{M}_d$ and then throughout the region by analytic continuation.

Because that region contains infinitely many PCF parameters, (5) gives infinitely many points $(\zeta c, c)$ on C . Irreducibility forces C itself to be the corresponding line. If a coordinate projection was not dominant, the curve was already one of the two PCF coordinate lines.

Why the two rigidity mechanisms fit

The proof uses arithmetic and complex dynamics in genuinely different ways. The adelic metric packages the global algebraic condition of critical height zero and makes an infinite PCF sequence visible to equidistribution. That step is insensitive to the individual orbit relations: it returns equality of measures, not an equation for C . Potential theory upgrades the measure identity to the escape-rate identity (2), but this still determines only the modulus in (3). External rays retain the missing angular information. Their landing pairs and wake widths then turn a priori analytic branches into the finite rotation group preserving f_c .

In proof-roadmap form, one first normalizes C . A nondominant projection immediately gives a coordinate line. Otherwise the two projections define non-isotrivial families, and the infinite simultaneous PCF set is a common small sequence for their heights. Equidistribution and harmonic continuation yield (2); Böttcher coordinates yield (3); rational landing rays make the phase rational; and the wake formula forces (5). Finally, infinitely many points of the line $x = \zeta y$ lie on the irreducible curve C , so the curves coincide (after replacing ζ by its inverse this is the theorem's form $y = \zeta x$).

A toy case and the hard point

When $d = 2$, the only $(d - 1)$ st root of unity is 1; hence the only dominant special curve is the diagonal $a = b$. This makes the final algebraic conclusion transparent, but removes the key difficulty: equality of Green functions controls only the modulus of a Böttcher coordinate. The external-ray landing theory, rationality of η , and the arithmetic of wake widths are what eliminate all nonlinear or fractional-angle correspondences.

The theorem concerns a fixed degree and the unicritical family over complex curves. It does not classify special subvarieties in the full moduli space of rational maps. Its equidistribution step is first proved over a number field; the main curve descends because its Zariski-dense PCF points are algebraic. The detailed landing and wake analysis is special to $\mathcal{M}_d$, so the method does not by itself extend to arbitrary polynomial families.