Derived categories of Fano varieties of lines

Alessio Bottini, Daniel Huybrechts

arXiv:2501.03534v1 (2025), 20 pages

http://arxiv.org/abs/2501.03534v1

A categorical square attached to a cubic fourfold

Let $X \subset \mathbf{P}^5$ be a smooth cubic fourfold and let F_X be its four-dimensional hyperkahler variety of lines. The Kuznetsov component

$$\mathcal{A}_X := \langle \mathcal{O}_X, \mathcal{O}_X(1), \mathcal{O}_X(2) \rangle^{\perp} \subset D^b(X)$$

is a K3 category. Its Hilbert square is the equivariant category $\mathcal{A}_X^{[2]} = (\mathcal{A}_X \boxtimes \mathcal{A}_X)^{\mathfrak{S}_2}$. Galkin's conjecture predicts $D^b(F_X) \simeq \mathcal{A}_X^{[2]}$.

Main theorem (Bottini–Huybrechts, Theorems 0.3 and 0.4)**.** *Assume that F_X admits a rational Lagrangian fibration and is generic with this property. Then there is an exact linear equivalence*

$$D^b(F_X) \simeq \mathcal{A}_X^{[2]}.$$

Equivalently, if X is in a Zariski-dense open subset of a Hassett divisor $\mathcal{C}_d$ and $d/2$ is a perfect square, the equivalence holds.

For every smooth cubic fourfold X, without this genericity assumption, there is a Hodge isometry between the natural naive weight-two structures

$$\widetilde{H}(F_X, \mathbf{Z}) \simeq \widetilde{H}(\mathcal{A}_X^{[2]}, \mathbf{Z}).$$

Here $\widetilde{H}(F_X, \mathbf{Z}) = H^2(F_X, \mathbf{Z}) \oplus U$, while $\widetilde{H}(\mathcal{A}_X^{[2]}, \mathbf{Z}) = \widetilde{H}(\mathcal{A}_X, \mathbf{Z}) \oplus^{\perp} \mathbf{Z}\delta$ with $\delta^2 = -2$. The first theorem proves the categorical conjecture on a new countable family of divisors; the second gives global cohomological evidence, but is not by itself a derived equivalence.

Ingredient 1: the twisted BKR bridge

If an equivalence $\mathcal{A}_X \simeq D^b(S, \alpha)$ identifies the K3 category with the derived category of a K3 surface twisted by $\alpha \in \mathrm{Br}(S)$, Proposition 1.2 proves

$$\mathcal{A}_X^{[2]} \simeq D^b(S^{[2]}, \alpha^{[2]}).$$

The usual Bridgeland–King–Reid kernel is the structure sheaf of the isospectral Hilbert scheme. Its restriction trivializes the difference between the pullbacks of $\alpha^{[2]}$ and $\alpha \boxtimes \alpha$, so the same kernel defines the twisted Fourier–Mukai functor. Locally it is the ordinary BKR equivalence; full faithfulness follows, and the connected Calabi–Yau target upgrades this to an equivalence.

Ingredient 2: Brauer classes survive the two flops

A Mukai flop $Z \dashrightarrow Z'$ between projective hyperkahler manifolds identifies their Brauer groups. Lemma 2.2 lifts corresponding classes α, α' from the two resolutions to the singular contraction. Consequently the Kawamata–Namikawa kernel becomes $(\alpha^{-1} \boxtimes \alpha')$-twisted and yields

$$D^b(Z, \alpha) \simeq D^b(Z', \alpha')$$

(Proposition 2.3). This is essential: the proof must transport not just the underlying fourfolds, but the precise Brauer obstruction needed by the twisted BKR step.

For a generic degree-two polarized K3 surface (S, H), let $M = M_S(0, H, -1)$ compactify the relative degree-zero Jacobian of the integral curves in $|H|$. A general length-two subscheme $\xi \subset S$ lies on a unique $C_\xi \in |H|$, and sending it to $I_{\xi/C_\xi} \otimes H|_{C_\xi}$ gives a Mukai flop $S^{[2]} \dashrightarrow M$. If M_α is the corresponding moduli space of α-twisted rank-one sheaves on those curves, the missing universal Poincare sheaf defines an obstruction $\theta \in \mathrm{Br}(M)$ and compactified autoduality gives

$$D^b(M_\alpha) \simeq D^b(M, \theta).$$

The delicate identification in Proposition 2.7 is $\theta = \pm\alpha^{[2]}$ under $\mathrm{Br}(M) \simeq \mathrm{Br}(S^{[2]})$; it is checked over the smooth relative Jacobian, where both classes pull back from the same class on the relative Hilbert scheme.

Ingredient 3: the moduli space singled out by F_X

When $d/2$ is a square and X is generic in $\mathcal{C}_d$, Proposition 2.8 produces a degree-two polarized K3 surface (S, H) and a Brauer class α such that

$$\mathcal{A}_X \simeq D^b(S, \alpha), \qquad F_X \dashrightarrow M_\alpha.$$

The first statement comes from extending the Hodge isometry induced by the birational correspondence to the index-two overlattices $\widetilde{H}(\mathcal{A}_X, \mathbf{Z})$ and $\widetilde{H}(S, \alpha, \mathbf{Z})$. The second generator of the distinguished A_2 lattice pairs to 1 with the Mukai vector, ensuring that M_α is fine and hence has a universal family.

The proof of the main equivalence is then the controlled chain

$$\begin{aligned} D^b(F_X) &\simeq D^b(M_\alpha) \simeq D^b(M, \theta) \simeq D^b(S^{[2]}, \alpha^{[2]}) \\ &\simeq D^b(S, \alpha)^{[2]} \simeq \mathcal{A}_X^{[2]}. \end{aligned}$$

Each arrow has a distinct job: birational hyperkahler invariance, relative Jacobian autoduality, identification of the obstruction across the flop, and twisted BKR. A useful toy picture is the untwisted case $\alpha = 0$, where $M_\alpha = M$, $\theta = 0$, and the chain collapses to the familiar fact that birational K3$^{[2]}$-type fourfolds have equivalent derived categories. The actual theorem must track the nonzero twist through every step.

The Hodge-theoretic companion

For Theorem 0.4, embed both lattices into $\widetilde{H}(\mathcal{A}_X, \mathbf{Z}) \oplus U$. One description is the orthogonal complement of a square-two generator $\lambda_1 \in A_2$; the other is the complement of $\gamma = e + f \in U$, also of square two, with δ sent to $e - f$. Their transcendental lattices agree by construction. Nikulin's extension theorem promotes this agreement, together with $\lambda_1 \mapsto \gamma$, to an isometry of the ambient lattices, and taking orthogonal complements gives the asserted Hodge isometry.

The categorical result remains generic: the authors can specialize the birational correspondence more broadly, but do not know how to remove the genericity/integral-curve assumptions required for compactified Jacobian autoduality. The Hodge isometry has no canonical control on the algebraic part and does not imply Galkin's conjecture for a general cubic fourfold.

This locates the remaining obstruction precisely. Proposition 2.8's twisted-K3 model and birational correspondence can, using later stability results, be specialized to every smooth cubic in $\mathcal{C}_d$ with $d/2$ a square. What is not available beyond the generic locus is the version of compactified relative-Jacobian autoduality used in Proposition 2.6 after singular or reducible curves enter the linear system. Thus the proof cannot simply promote its dense-open statement by specialization.

Comparison with the birational-Hilbert locus

Theorem 0.2 treats a different countable family. If F_X is birational to $S^{[2]}$ for a K3 surface S, a birational map between hyperkahler fourfolds factors into simple Mukai flops, hence $D^b(F_X) \simeq D^b(S^{[2]})$. The corresponding Hassett numerical condition also implies $\mathcal{A}_X \simeq D^b(S)$ for the same S; ordinary BKR then gives

$$D^b(F_X) \simeq D^b(S^{[2]}) \simeq D^b(S)^{[2]} \simeq \mathcal{A}_X^{[2]}.$$

The Lagrangian-fibration theorem is subtler because it typically produces a twisted K3 category $D^b(S, \alpha)$ rather than $D^b(S)$. The relative Jacobian and $\theta = \pm\alpha^{[2]}$ are what keep this twist from being lost.

Further integral-structure evidence

For a general K3$^{[2]}$-type hyperkahler manifold the naive lattice $H^2 \oplus U$ need not be a derived invariant; Beckmann and Taelman replace it by a half-integral B-field transform. Proposition 3.4 shows that for F_X this corrected lattice is nevertheless Hodge isometric to the naive one. Lemma 3.3 supplies the mechanism: if the half-class defining the transform is the sum of an algebraic rational class and an integral class, the algebraic transform is a Hodge isometry and the integral transform preserves the lattice. For F_X, specialization to a Pfaffian cubic gives exactly such a decomposition in terms of the Plucker polarization; parallel transport returns it to the original cubic.

The paper also tracks the sign involution of $\mathcal{A}_X^{[2]}$: it acts by -1 on $\widetilde{H}(\mathcal{A}_X, \mathbf{Z})$ and by $+1$ on the extra class δ. These results strengthen the cohomological evidence but do not remove the genericity needed for the categorical theorem.