Good Reduction of Periodic Points on Projective Varieties

Benjamin Hutz

Benjamin Hutz, arXiv:0801.3645v3 (2010), 18 pages

http://arxiv.org/abs/0801.3645v3

How much of a period survives reduction?

Let R be a complete discrete valuation ring with normalized valuation v, uniformizer π, fraction field K, and finite residue field k of characteristic p and cardinality $N\pi$. Let $\mathcal{X}/R \subset \mathbf{P}^N_R$ be smooth and projective, with generic fiber X/K nonsingular, irreducible, and of dimension d. An R-morphism $\phi_R : \mathcal{X} \to \mathcal{X}$ is a good-reduction model of its generic-fiber map ϕ; write $\bar{\phi}$ for its special fiber. Properness identifies each $P \in X(K)$ with a unique $P_R \in \mathcal{X}(R)$ and hence a reduction $\bar{P}$.

Main theorem (Hutz, Theorems 1–2 and Corollary 3)**.** *Suppose P_R has primitive period n and $\bar{P}$ has primitive period m. Either $n = m$, or*

$$n = m r_V p^e \qquad (e \geq 0),$$

where V is a $\bar{\phi}$-stable subspace of the cotangent space of the special fiber along the orbit scheme and r_V is the order of the map induced by $\bar{\phi}^m$ on V. If $d' = \dim V$, then $r_V \leq (N\pi)^{d'} - 1$; moreover e is bounded explicitly in terms of $v(p)$ (with a separate $p = 2$ formula).

Over $\mathbb{Q}$, at a prime p of good reduction,

$$e \leq \begin{cases} 1, & p \neq 2, \\ 3, & p = 2, \end{cases} \qquad n \leq \begin{cases} p^{d+1}(p^d - 1), & p \neq 2, \\ 2^{d+3}(2^d - 1), & p = 2. \end{cases}$$

Thus the lift from finite-field dynamics has only two sources of extra period: a linear action on cotangent directions and a controlled p-primary correction.

Ingredient 1: reduction and integral local coordinates

Reduction commutes with composition and iteration, so $\bar{P}$ is periodic and $m \mid n$ (Theorem 7 and Corollary 9). After replacing ϕ_R by ϕ_R^m, one may assume $\bar{P}$ is fixed.

Let $\mathcal{V}$ be the scheme-theoretic union of the n orbit points. Lemma 19 first moves the whole orbit into one affine chart by an element of $\mathrm{PGL}_{N+1}(R)$. Smoothness and a π-adic implicit function theorem then provide local parameters in which the restriction to $\mathcal{V}$ is an iterable convergent series

$$f(T) = \omega + df_P(T) + \text{terms of degree at least two}, \qquad \omega \in \mathfrak{m}^{d'},$$

with coefficients in R. This is the geometric step: without one integral coordinate neighborhood containing the orbit, the later congruence calculation would have no common domain.

Ingredient 2: remove the cotangent-order factor

The cyclic action of ϕ_R on $\mathcal{V}$ implies that the reduction of $d\phi_P$ has finite order r_V on its cotangent space. Replacing f by its r_V-th iterate makes

$$df_P \equiv I \pmod{\pi}.$$

If a residual period $s > 1$ remains, write $df_P = I + \pi^c A$. Iterating modulo the square of the ideal generated by ω gives

$$f^s(0) \equiv \big(sI + \text{terms divisible by } \pi^c\big)\omega \equiv 0 \pmod{(\omega)^2}.$$

Because one coordinate of ω generates (ω), this forces $s \equiv 0 \pmod{\pi}$, hence $p \mid s$. Repeating leaves precisely a power p^e. The order estimate for r_V comes from the resulting element of $\mathrm{GL}_{d'}(\mathbb{F}_{N\pi})$.

In the one-dimensional toy series $f(z) = a + \lambda z + O(z^2)$ with $a \in \mathfrak{m}$, first iterating through the order of $\bar{\lambda}$ gives tangent multiplier 1 modulo π; then $f^s(0) \equiv sa \pmod{a^2}$ exhibits the same forced p-divisibility. The full proof must additionally build the coordinates on $\mathcal{X}$ and restrict them to the possibly smaller orbit scheme $\mathcal{V}$.

Ingredient 3: valuation growth bounds the exponent

For the remaining p^e-cycle, set $x_k = f^{p^k}(0)$,

$$b_k = \min_i v((x_k)_i), \qquad d(f^{p^k})_0 = I + \pi^{c_k} A_k.$$

Here A_k has at least one unit entry, so c_k is the exact π-adic order of $d(f^{p^k})_0 - I$; if that derivative is I, set $c_k = \infty$. Lemma 22 derives the coupled recurrences

$$b_{k+1} \geq \min\{2b_k, b_k + (p-1)c_k, b_k + v(p)\}, \quad c_{k+1} \geq \min\{b_k, c_k + v(p), pc_k\}.$$

Before the cycle closes, $\min(b_k, c_k) \leq v(p)$. Lemma 23 shows that the lower bounds grow like powers of two for odd p and like Fibonacci numbers for $p = 2$; they therefore cross the fixed ceiling $v(p)$ after only boundedly many steps. This is the hard quantitative input and yields Theorem 2 and the displayed rational bound.

There is also an important étale extension (Theorem 4). Let ϕ be étale on a projective variety, let P be a smooth periodic point, let $Y \subset X_{\mathrm{smooth}}$ be its irreducible component, and let l be the first positive integer for which ϕ^l is an étale endomorphism of Y. If $\phi^l|_Y$ has good reduction, then the primitive period is one of

$$lm, \qquad lmr_V, \qquad lmr_V p^e,$$

with the local factors above. This removes smoothness and irreducibility of the whole variety, but it still concerns morphisms, not rational maps with indeterminacy. The bounds are local and deliberately coarse. A complementary result (Theorem 20) says that for a fixed number-field periodic point, reduction preserves its exact primitive period at all but finitely many good primes.