The intermediate type of certain moduli spaces of curves

Gavril Farkas, Alessandro Verra

arXiv:0910.3905v2 (2009), 20 pages
http://arxiv.org/abs/0910.3905v2

Two moduli spaces between uniruled and general type

Let $\overline{\mathcal{S}}_g^+$ compactify the moduli of even spin curves, let $\overline{\mathcal{M}}_{g,n}$ parametrize stable pointed curves, and let $\overline{\Theta}_{\rm null}$ be the effective-theta-characteristic divisor. On $\overline{\mathcal{M}}_{g,g}$ put

$$\overline{\mathcal{D}}_g = \big\{[C, x_1, \ldots, x_g] : h^0\big(C, \mathcal{O}_C(x_1 + \cdots + x_g)\big) \geq 2\big\}.$$

Main theorem (Farkas–Verra, Theorems 0.1, 0.3, and 0.4)**.** *The Kodaira dimension of* $\overline{\mathcal{S}}_8^+$ *is zero, while*

$$\kappa\big(\overline{\mathcal{M}}_{11,11}\big) = 19 \quad \textit{although} \quad \dim \overline{\mathcal{M}}_{11,11} = 41.$$

Moreover, for $g \leq 11$ *the effective divisor* $\overline{\mathcal{D}}_g$ *is extremal and rigid.*

Here "Calabi–Yau type" means the birational statement $\kappa = 0$, not that the coarse space is asserted to be a smooth Calabi–Yau variety.

Ingredient 1: negative covering curves detect fixed divisors

If an irreducible divisor D is swept by curves R with $R \cdot D < 0$, every effective divisor meeting R negatively contains D. Iteration makes prescribed multiples of D fixed. The task is to construct such curves with zero intersections against all competing components.

Ingredient 2: a doubly elliptic K3 pencil in genus eight

The canonical class admits the effective decomposition

$$K_{\overline{\mathcal{S}}_8^+} \equiv \tfrac{1}{2}\pi^*(\mathfrak{bn}_8) + 8\overline{\Theta}_{\rm null} + \sum_{i=1}^{4}(a_i\alpha_i + b_i\beta_i), \qquad a_i, b_i > 0,$$

where $\mathfrak{bn}_8$ is the normalized plane-septic divisor and α_i, β_i are boundary classes. Mukai's model extends a general canonical $C \in \overline{\Theta}_{\rm null}$ from its rank-three quadric to a K3 surface on a rank-four quadric. Projection from the singular locus is a degree-seven map to $\mathbf{P}^1 \times \mathbf{P}^1$; a plane pencil pulls back to $R \subset \overline{\Theta}_{\rm null}$ with

$$R \cdot \overline{\Theta}_{\rm null} = -1, \qquad R \cdot \pi^*(\overline{\mathcal{M}}_{8,7}^2) = 0, \qquad R \cdot \alpha_i = R \cdot \beta_i = 0 \ (1 \leq i \leq 4).$$

Its two reducible fibers are pairs of elliptic curves meeting in seven points; the local spin multiplicities give the negative intersection. Thus $8\overline{\Theta}_{\rm null}$ is fixed, and a covering pencil for the plane-septic divisor kills the residual moving part: $\kappa = 0$.

Ingredient 3: the pointed-curve divisor is forced to stay

For $g \leq 11$, $g \neq 10$, K3 Lefschetz pencils through the marked points give covering curves with $R \cdot \overline{\mathcal{D}}_g = -1$. Genus ten is exceptional: identify two marked points to form a nodal genus-eleven curve on a K3, blow up its node, split the resulting two-section after a double base change, and symmetrize over pairs of labels. The resulting cycle still has intersection -1, proving rigidity. In genus eleven the pencil also misses the boundary and $\phi^*(\mathfrak{bn}_{11})$, and

$$K_{\overline{\mathcal{M}}_{11,11}} \equiv \overline{\mathcal{D}}_{11} + 2\phi^*(\mathfrak{bn}_{11}) + \sum_{i,c} d_{i:c}\,\delta_{i:c}, \qquad d_{i:c} \geq 0.$$

Hence $n\overline{\mathcal{D}}_{11}$ is fixed in $|nK|$. Rational-tail K3 pencils similarly remove the $\delta_{0:c}$, leaving a class controlled by $\overline{\mathcal{M}}_{11}$.

Ingredient 4: the Mukai fibration gives exactly nineteen

A general genus-eleven curve lies on a unique degree-20 polarized K3, giving the Mukai map

$$q_{11} : \overline{\mathcal{M}}_{11,11} \dashrightarrow \overline{\mathcal{F}}_{11}, \qquad [C, x_1, \ldots, x_{11}] \longmapsto [S \supset C, \mathcal{O}_S(C)],$$

whose target has dimension 19. The residual canonical class dominates the pullback of an ample class, so $\kappa \geq 19$. Conversely K3 Lefschetz pencils form an 11-dimensional family contracted by every corresponding linear system, so $\kappa \leq 19$.

The hardest input is the rank-four quadric extension and its boundary multiplicity. The paper also proves uniruledness of $\overline{\mathcal{M}}_{5,n}$ for $n \leq 14$, $\overline{\mathcal{M}}_{7,n}$ for $n \leq 13$, and $\overline{\mathcal{M}}_{8,n}$ for $n \leq 12$, using a two-divisor criterion.

Codex mathematical paper digest, 2026-10-25. Source paper should be consulted for full statements and proofs.