

The failure of Kodaira vanishing for Fano varieties, and terminal singularities that are not Cohen-Macaulay

Burt Totaro

arXiv:1710.04364

<http://arxiv.org/abs/1710.04364v1>

Positive characteristic breaks two familiar implications

Over characteristic zero, Kodaira vanishing gives $H^i(X, K_X + L) = 0$ for an ample line bundle L and $i > 0$. In the minimal model program it also underlies the good depth properties of mild singularities. Totaro shows that even smooth Fano varieties need not satisfy this vanishing in characteristic p , and that terminal singularities need not be Cohen–Macaulay.

Main theorem (Totaro, Theorem 2.1 and Corollary 2.2). *For every prime $p \geq 3$ there is a smooth Fano variety $X/\mathbf{F}_p$ of dimension $2p + 1$ and Picard number 2 with*

$$-K_X = 2A, \quad H^1(X, A) \neq 0,$$

where A is very ample. Moreover $\chi(X, A) < 0$ for $p \geq 5$. Consequently the affine cone

$$Y = \operatorname{Spec} \bigoplus_{m \geq 0} H^0(X, mA)$$

is an isolated terminal singularity of dimension $2p + 2$ which is not Cohen–Macaulay.

Because $K_X + 3A = A$, the nonzero group is exactly a failure of Kodaira vanishing for the ample bundle $3A$. A second family (Theorem 3.1) works in every characteristic: it has dimension $2p - 1$ for $p \geq 3$, dimension 5 for $p = 2$, Picard number 2, and a very ample L with $H^1(X, K_X + L) \neq 0$.

Ingredient I: a Fano “unseparated flag variety”

Let $G = \operatorname{SL}(n)$ in characteristic p . A function on its simple roots defines a possibly non-reduced subgroup scheme $P \supset B$; Totaro takes

$$f(\alpha_1) = 1, \quad f(\alpha_2) = 0, \quad f(\alpha_i) = \infty \quad (3 \leq i \leq n - 1).$$

Although P is not smooth, $X = G/P$ is smooth, projective, rational, and has the same dimension $2n - 3$ as the flag variety $G/P_{\text{red}} = \operatorname{Fl}(1, 2, n)$. Its Picard group has basis $p\omega_1, \omega_2$, while the root calculation gives

$$-K_X = 2p\omega_1 + (n - p)\omega_2. \tag{1}$$

Taking $n = p + 2$ makes (1) equal to $2A$ for $A = p\omega_1 + \omega_2$, proving the Fano and divisibility assertions. The non-reduced stabilizer is decisive: it keeps the homogeneous space smooth but changes its line-bundle cohomology from that of an ordinary flag variety.

Ingredient II: force a cokernel in degree one

There is a Frobenius-twisted $\mathbf{P}^1$ -bundle $\pi : G/P \rightarrow G/Q = \operatorname{Gr}(2, n)$, and A has degree one on its fibres. Pulling the rank-two bundle $\pi_* A$ to G/B corresponds to an extension of B -representations

$$0 \longrightarrow k(\lambda - p\alpha) \longrightarrow M \longrightarrow k(\lambda) \longrightarrow 0, \quad \lambda = p\omega_1 + \omega_2, \alpha = \alpha_1. \tag{2}$$

Andersen’s reflection isomorphism shifts the cohomology of $\lambda - p\alpha$ by one, while Kempf vanishing applies to the dominant weights λ and $\lambda - \alpha = (p - 2)\omega_1 + 2\omega_2$. The long exact sequence collapses to

$$0 \rightarrow H^0(X, A) \rightarrow H^0(G/B, \lambda) \rightarrow H^0(G/B, \lambda - \alpha) \rightarrow H^1(X, A) \rightarrow 0, \tag{3}$$

with $H^i(X, A) = 0$ for $i \geq 2$.

The Weyl dimension formula makes the target/source ratio in (3)

$$\frac{(p-1)(p+2)^2}{4(p+1)^2} > 1 \quad (p \geq 5), \quad (4)$$

so the map cannot be surjective. For $p = 3$ the dimensions are 224 and 175 in the opposite order, but the simple socle $L(\lambda)$ of dimension 50 maps to zero because its highest weight does not occur in the target; $224 - 50 < 175$ again prevents surjectivity. This representation-theoretic non-surjectivity is the core of the vanishing counterexample. Changing to $n = p + 1$ yields Theorem 3.1; the exceptional $p = 2$ calculation instead uses Euler characteristic and Andersen’s shift directly.

Ingredient III: the cone translates cohomology into depth

For a cone over (X, A) , the vertex is terminal precisely when $A \sim_{\mathbf{Q}} a(-K_X)$ with $0 < a < 1$. Here $a = 1/2$. The same cone is Cohen–Macaulay precisely when

$$H^i(X, mA) = 0 \quad (0 < i < \dim X, m \in \mathbf{Z}). \quad (5)$$

The already established $H^1(X, A) \neq 0$ therefore proves Corollary 2.2. This is the conceptual bridge: a failure of Kodaira-type vanishing on a smooth projective Fano base becomes a depth failure at an isolated terminal vertex.

The sharp threefold example in characteristic two

Theorem 5.1 gives a different, lowest-dimensional construction. Over $\mathbf{F}_2$, let $\mathbf{Z}/2$ act on $Y = (\mathbf{G}_m)^3$ by simultaneous inversion,

$$\sigma(x_1, x_2, x_3) = (x_1^{-1}, x_2^{-1}, x_3^{-1}). \quad (6)$$

The action is free away from $(1, 1, 1)$. Fogarty’s theorem makes $X = Y/(\mathbf{Z}/2)$ non-Cohen–Macaulay because the fixed locus is nonempty of codimension three.

Terminality requires an explicit wild resolution. Blow up the fixed point, then blow up the seven exceptional directions over $\mathbf{F}_2$. On the resulting smooth Y_2 , the fixed-point scheme is the Cartier divisor

$$E_0 + 2 \sum_{j=1}^7 E_j; \quad (7)$$

hence $Y_2/(\mathbf{Z}/2)$ is smooth by the Király–Lütkebohmert criterion. Put $\tilde{X} = Y_2/(\mathbf{Z}/2)$, let $q : Y_2 \rightarrow \tilde{X}$ be the quotient, write $F_j = q(E_j)$, and let $\rho : \tilde{X} \rightarrow X$ be the resolution. Artin–Swan ramification numbers show that q is fiercely ramified along E_0 , with $q^*F_0 = E_0$, and wildly ramified along E_j , with $q^*F_j = 2E_j$ for $1 \leq j \leq 7$. Combining the different with the canonical-divisor formula for the two rounds of point blow-ups gives

$$K_{\tilde{X}} = \rho^*K_X + F_0 + \sum_{j=1}^7 F_j. \quad (8)$$

Every exceptional discrepancy is positive, so X is terminal.

Hardest point and scope

The homogeneous examples reduce their geometry to the exact sequence (3), but the hard step is detecting non-surjectivity in the small characteristic where dimensions alone do not decide it. For the threefold quotient, the subtlety is fiercer: a fixed divisor in wild characteristic can map purely inseparably, so one must distinguish fierce from wild ramification before reading discrepancies from (7). The paper constructs counterexamples; it does not rule out a dimension-dependent bound $p_0(n)$ above which smooth Fanos satisfy Kodaira vanishing, or above which klt singularities are Cohen–Macaulay. Those are its closing open questions.