

Explicit canonical heights for divisors relative to endomorphisms of projective N-space

Patrick Ingram

arXiv:2207.07206

<http://arxiv.org/abs/2207.07206v1>

The divisor-height problem

Let $f : \mathbb{P}^N \rightarrow \mathbb{P}^N$ be an endomorphism of degree $d \geq 2$ and let D be an effective divisor, both defined over $\overline{\mathbb{Q}}$. If $h(D)$ is the normalized Philippon height and f_*D is the cycle-theoretic image, the canonical height is

$$\widehat{h}_f(D) = \lim_{k \rightarrow \infty} \frac{h(f_*^k D)}{d^{kN}}. \quad (1)$$

This normalization is linear in D ; Zhang's canonical height is $\widehat{h}_f(D)/\deg D$. The computational obstacle is to control the error in (1) uniformly in the iterate and only linearly in $\deg D$.

Main theorem (Ingram, Theorem 1). *There are explicit constants C_1, C_2 , depending only on $N \geq 1$ and $d \geq 2$, such that, for f and D as above,*

$$|\widehat{h}_f(D) - h(D)| \leq (C_1 h(f) + C_2) \deg D, \quad (2)$$

where $h(f)$ is the projective height of the coefficient tuple of f . One may take

$$C_1 = 5Nd^{N+1}, \quad C_2 = 3^N N^{N+1} (2d)^{N2^{N+4}d^N}.$$

The constants are deliberately generous, but their complete effectivity is the point. Since $\deg(f_*D) = d^{N-1} \deg D$ and $\widehat{h}_f(f_*D) = d^N \widehat{h}_f(D)$, applying (2) to $f_*^k D$ gives

$$\left| \widehat{h}_f(D) - \frac{h(f_*^k D)}{d^{kN}} \right| \leq d^{-k} (C_1 h(f) + C_2) \deg D. \quad (3)$$

Thus a prescribed error tolerance determines an explicit stopping time. It also makes preperiodic hypersurfaces of a given geometric degree, defined over number fields of bounded degree, effectively enumerable, because every preperiodic divisor has canonical height zero.

Local lifts and the correct scaling

At a place v , lift f to homogeneous degree- d forms $F = (F_0, \dots, F_N)$ on $\mathbb{A}^{N+1}$. If $D = (\Phi = 0)$, define

$$F^*\Phi = \Phi \circ F, \quad F_*\Phi(Y) = \prod_{F(X)=Y} \Phi(X).$$

The second form has degree $d^N \deg \Phi$. A crucial multiplicity convention is that $F_*\Phi$ cuts out $d f_*D$, not merely f_*D ; consequently

$$F_*F^*\Phi = \Phi^{d^{N+1}}, \quad f_*f^*D = d^N D. \quad (4)$$

The resultant measures bad reduction through the lift-invariant local quantity

$$\lambda_{\text{Hom}_d^N}(f) = -\log |\text{Res}(F)| + (N+1)d^N \log \|F\|.$$

The first proof ingredient is an explicit Macaulay identity (Lemma 4): bounded polynomials $A_{i,j}$ express a power of each coordinate times $\text{Res}(F)$ as $\sum_i F_i A_{i,j}$. Together with the elementary upper estimate, it gives two-sided point escape bounds (Lemma 5), hence the local Green function

$$G_F(P) = \lim_{k \rightarrow \infty} d^{-k} \log \|F^k(P)\| \quad (5)$$

with an explicit error from $\log \|P\|$. Macaulay elimination is where the very large universal constant in (2) originates.

From points to equations of hypersurfaces

Write $m(\Phi)$ for the logarithmic Gauss norm at a non-Archimedean place and for logarithmic Mahler measure at an Archimedean place. Mahler's inequalities compare $m(\Phi)$ with coefficient norms with error linear in $\deg \Phi$. Lemma 7 turns this into a testing principle: for every P ,

$$\log |\Phi(P)| \leq m(\Phi) + \deg(\Phi) \log \|P\| + N \deg(\Phi) \log^+ |2|,$$

and some P satisfies the reverse inequality without the last term.

This bridge lets Lemma 8 compare $m(F_*\Phi)$ and $d^{N+1}m(\Phi)$ in both directions. Its harder lower bound factors $F^*F_*\Phi = \Phi\Psi$: point escape controls the norms of two points in a common fiber, while Mahler's estimate controls the auxiliary factor Ψ . The resulting errors are explicit and linear in $\deg \Phi$. A telescoping series therefore converges to the form escape rate (Lemma 9)

$$G_F(\Phi) = \lim_{k \rightarrow \infty} d^{-k(N+1)} m(F_*^k \Phi), \quad (6)$$

with $|G_F(\Phi) - m(\Phi)| \leq C(F, N, d) \deg \Phi$. It is additive in products, invariant under pullback, scales by d^{N+1} under pushforward, and has precisely the scalar and coordinate covariance needed to descend from F, Φ to f, D .

Product formula and proof roadmap

Over a number field K , sum the local quantities in (6):

$$\widehat{h}_f(D) = \sum_{v \in M_K} \frac{[K_v : \mathbb{Q}_v]}{[K : \mathbb{Q}]} G_{F,v}(\Phi). \quad (7)$$

Only finitely many summands are nonzero, and the product formula removes the dependence on the chosen lifts. Summing $m_v(\Phi)$ recovers $h(D)$; summing $\log \|F\|_v$ gives $h(f)$; and summing the local resultant term gives $(N+1)d^N h(f)$. Substitution of Lemma 9's upper and lower errors, followed by the explicit Macaulay bounds, yields the two sides of (2). Equation (4) then identifies (7) with the limit (1).

For the toy case $N = 1$ and $f(z) = z^d$, a point divisor $D = [a]$ is sent to $[a^{d^k}]$ after k iterates, so $d^{-k} h([a^{d^k}]) = h([a])$. The general argument retains this exact scaling while paying explicit errors for arbitrary coefficients, higher-dimensional fibers, and bad reduction. The same machinery constructs local Arakelov–Green pairings. Theorem 2 further shows that the critical height vanishes for postcritically finite maps, scales by n under $f \mapsto f^n$, is bounded above by an ample moduli height, and is comparable to that height along a one-parameter family when the generic critical height is nonzero; these applications are not needed for Theorem 1.