Moduli of sheaves and the Chow group of K3 surfaces

Kieran G. O'Grady

arXiv:1205.4119

http://arxiv.org/abs/1205.4119v2

Problem and setup

Let X be a projective complex K3 surface, and let $c_X \in \mathrm{CH}_0(X)$ be the Beauville–Voisin class of degree one. For $g \geq 0$, define

$$S_g(X) = \{[Z] + ac_X : Z \text{ is an effective zero-cycle of degree } g,\ a \in \mathbf{Z}\}.$$

These cones form an increasing exhaustive filtration of $\mathrm{CH}_0(X)$. For a Mukai vector $v = (r, \ell, s)$, put

$$d(v) = 1 + \frac{v^2}{2}, \qquad c_2(v) = r + \frac{\ell^2}{2} - s.$$

Let $M_v(X,H)^{\mathrm{st}}$ denote the stable locus and $\overline{M_v(X,H)^{\mathrm{st}}}$ its closure in the moduli space of semistable pure sheaves. When $r \geq 1$ and $v^2 \geq -2$, the paper sets $|v| = r^2v^2/4 + r^4/2$. A v-wall is $\mathrm{Amp}(X)_{\mathbf{R}} \cap \alpha^\perp$ for $\alpha \in \mathrm{NS}(X)$ with $-|v| \leq \alpha^2 < 0$; an open v-chamber is a connected component of the complement of these locally finite walls. The phrase *v-generic* means that $[H]$ lies in such a chamber.

Two elementary cases already display the prediction. If $r = 1$, the moduli space parametrizes $I_Z \otimes L$ with $\mathrm{length}(Z) = d(v)$ and $c_2(I_Z \otimes L) = [Z]$. If $r = 0$ and $\ell \neq 0$, a stable pure sheaf is supported on a curve $C \in |L|$; Claim 1.1 and the Severi argument below place its second Chern class in $S_{g(L)}(X) = S_{d(v)}(X)$. When $r = \ell = 0$, one has $d(v) = 1$ and the moduli space is X, with p corresponding to the skyscraper sheaf $\mathbf{C}_p$ and $c_2(\mathbf{C}_p) = -[p] \in S_1(X)$. Proposition 1.3 then supplies the reverse inclusion.

Main theorem (Theorem 0.6)**.** *Assume $M_v(X,H)^{\mathrm{st}} \neq \varnothing$. Then*

$$\{c_2(F) : [F] \in \overline{M_v(X,H)^{\mathrm{st}}}\} = \{z \in S_{d(v)}(X) : \deg z = c_2(v)\}$$

provided at least one of the following holds:

1. *$\ell = c_1(\mathcal{O}_X(H))$ is primitive and $s \geq 0$;*
2. *$\rho(X) \geq 2$, the rank r is coprime to the divisibility of ℓ, and H is v-generic;*
3. *$r \leq 2$, with H required to be v-generic when $r = 2$.*

Thus, in these cases, the possible second Chern classes depend only on their degree and the half-dimension $d(v)$ of the stable moduli space. In particular, equal-dimensional such moduli spaces produce the same set after translating degrees by a multiple of c_X.

Proof mechanism

1. Curves supply the filtration. If $f : C \to X$ is nonconstant and C is smooth of genus g, Riemann–Roch on C gives $f_* \mathrm{CH}_0(C) \subset S_g(X)$ (Claim 0.2). Degenerations of Severi varieties then produce a closed subset $\Sigma_g(X,L) \subset |L|$ of dimension at least g such that every curve $C \in \Sigma_g(X,L)$ has $\iota_* \mathrm{CH}_0(C) \subset S_g(X)$ (Proposition 1.6). This statement survives specialization in families of K3 surfaces. More precisely, Claim 1.2 expresses membership in S_g through relative Hilbert schemes (equivalently, genus-zero stable maps) as a countable union of closed parameter loci. A dense open family lies in one such closed locus, so the limiting fiber retains the same S_g bound. This specialization principle is used repeatedly after proving the assertion on a generic polarized K3.

2. Intersect a degeneracy-locus family with the right curves. For $v = (r, c_1(L), s)$ with L ample and $s \geq 0$, stability and Serre duality give $h^0(F) \geq \chi(F) = r + s$. An r-plane $U \subset H^0(F)$ defines

$$U \otimes \mathcal{O}_X \longrightarrow F, \qquad U \longmapsto C_U = V(\det \varphi_F^U) \in |L|.$$

When every U has generic rank r, this is a morphism from the whole Grassmannian. Its pullback of the hyperplane class is the Pluecker class (Lemma 2.1), so it is finite and its image has dimension at least

$$rs = \dim |L| - d(v).$$

It must meet $\Sigma_{d(v)}(X,L)$. For a curve C_U in the intersection, the exact sequence $0 \to U \otimes \mathcal{O}_X \to F \to \iota_*\xi \to 0$ gives $c_2(F) = -c_2(\iota_*\xi)$. Claim 1.1 writes $c_2(\iota_*\xi) = \iota_*\eta + ac_X$. Since $\iota_* \mathrm{CH}_0(C_U) \subset S_{d(v)}(X)$, and $S_{d(v)}(X)$ is preserved by integer multiplication (Corollary 1.7) and by adding multiples of c_X, this proves $c_2(F) \in S_{d(v)}(X)$.

3. Remove indeterminacy in the first two regimes. For primitive determinant, a generic polarized K3 has no effective divisor of small enough slope to be generated by a rank-deficient U. Proposition 2.3 therefore applies

there, and deformation plus specialization carries the containment back to every fiber (Proposition 3.1). If $\rho(X) \geq 2$ and r is coprime to the divisibility of ℓ, one twists by a sufficiently large multiple of an ample class inside the same v-chamber. The new determinant $\ell + ryh$ is primitive and the new final Mukai coordinate is positive; the primitive case applies, and tensoring back preserves the assertion (Proposition 3.3).

4. Resolve the rank-two obstruction. For $r = 2$, the degeneracy map can remain rational. After deforming to a locus where all sufficiently small effective divisors are proportional to the determinant, write $L = L_0^m$. A rank-deficient pair of sections generates L_0^a with $0 < a < m/2$. Twisting by L_0^{-a} replaces the determinant multiplicity m by $m - 2a$ and the final Mukai coordinate by

$$s_a = s - a(m - a)c_1(L_0)^2.$$

Corollary 4.6 forces $s_a \geq -1$. If $s_a \geq 0$, induction on m applies; if $s_a = -1$, Proposition 4.7 constructs the generic sheaf from a unique nontrivial extension $0 \to \mathcal{O}_X \to F \to I_Z \otimes L \to 0$, producing a birational Hilbert-scheme model and the required cycle formula. Specialization returns the result to the original K3 surface (Propositions 4.7–4.9 and the proof of Proposition 4.1).

5. Upgrade containment to equality. The construction above gives only $c_2(F) \in S_{d(v)}(X)$. Proposition 1.3 considers the map $X^{[d(v)]} \to S_{d(v)}(X)$ obtained by adding the degree-correcting multiple of c_X. A closed locus mapping onto the realized Chern classes must have dimension $2d(v)$: on a generically finite cover of the stable moduli space, the Mukai–Tyurin form is $-4\pi^2$ times the pullback of the Hilbert-scheme symplectic form. A smaller image would make the former degenerate. This contradiction proves surjectivity onto the entire degree slice.

Toy model, difficulty, and scope

For rank one, the theorem is essentially the definition of $S_{d(v)}(X)$: the subscheme Z is visible directly. Higher rank replaces it by a degeneracy curve carrying the residual sheaf, while the symplectic argument recovers every permitted rational-equivalence class rather than merely the classes constructed by that curve.

The hardest point is controlling the indeterminacy of the degeneracy map; the separate rank-two induction is precisely what is needed when slope arguments do not make every r-plane generically generating. The paper conjectures the equality without the theorem's extra hypotheses, and it leaves the analogous question $c_2(F) \in S_{d(F)}(X)$ for arbitrary simple sheaves open. Its final section also asks which correspondences preserve the filtration and relates $S_g(X)$ to Voisin's product filtration, but these directions are not inputs to Theorem 0.6. All statements here concern projective complex K3 surfaces and the closure of the stable locus, not every semistable sheaf in an unrelated component.