

Dynamical canonical heights for Jordan blocks, arithmetic degrees of orbits, and nef canonical heights on abelian varieties

Shu Kawaguchi, Joseph H. Silverman

arXiv:1301.4964v2

<http://arxiv.org/abs/1301.4964v2>

Exponential growth of an orbit

Let K be a global field in the paper's sense (in particular, $\text{char } K = 0$), let X/K be normal and projective, and let $f: X \rightarrow X$ be a K -morphism. For a Weil height h_H attached to an ample divisor, put $h_H^+ = \max\{h_H, 1\}$. The arithmetic degree, when the limit exists, is

$$\alpha_f(x) = \lim_{n \rightarrow \infty} h_H^+(f^n(x))^{1/n};$$

it is independent of the chosen ample height and records only exponential height growth.

Main theorem (Kawaguchi–Silverman, Theorem 2). *For every $x \in X(\overline{K})$ the limit defining $\alpha_f(x)$ exists. As x varies, the values $\alpha_f(x)$ form a finite set of algebraic integers.*

The result allows non-dominant morphisms and singular X (subject to normality). Its engine is a canonical-height construction that retains the nilpotent part of f^* rather than diagonalizing it away.

Ingredient I: canonical heights for a Jordan block

Suppose $D_0, \dots, D_\ell \in \text{Div}(X)_{\mathbb{C}}$ satisfy

$$f^* D_0 \sim \lambda D_0, \quad f^* D_k \sim D_{k-1} + \lambda D_k \quad (1 \leq k \leq \ell), \quad |\lambda| > 1. \quad (1)$$

Writing Λ for the corresponding lower Jordan block, Theorem 12 constructs a unique vector height $\hat{\mathbf{h}} = (\hat{h}_0, \dots, \hat{h}_\ell)^t$ with

$$\hat{\mathbf{h}} = \mathbf{h} + O(1), \quad \hat{\mathbf{h}} \circ f = \Lambda \hat{\mathbf{h}}. \quad (2)$$

The point is that $\mathbf{E} = \mathbf{h} \circ f - \Lambda \mathbf{h}$ is bounded: its coordinates are heights of linearly trivial divisors. Since $\|\Lambda^{-n}\| = O(n^\ell |\lambda|^{-n})$, the correction series

$$\hat{\mathbf{h}} = \mathbf{h} + \sum_{n \geq 0} \Lambda^{-n-1} \mathbf{E} \circ f^n$$

converges uniformly. This proves existence, (2), and the $O(1)$ normalization; the same exponential estimate forces uniqueness. Without $|\lambda| > 1$, Theorem 12(a) still bounds the raw vector height by $O(n^\ell \max\{|\lambda|, 1\}^n)$.

A two-dimensional model

For $f^*D_0 \sim \lambda D_0$ and $f^*D_1 \sim D_0 + \lambda D_1$, iteration of (2) gives

$$\widehat{h}_0(f^n x) = \lambda^n \widehat{h}_0(x), \quad \widehat{h}_1(f^n x) = \lambda^n \widehat{h}_1(x) + n\lambda^{n-1} \widehat{h}_0(x).$$

Thus the nilpotent direction contributes a polynomial factor n , but the n th root still sees only $|\lambda|$. This elementary observation is the mechanism behind the general arithmetic-degree theorem.

Ingredient II: reduce all height growth to finitely many blocks

Lemma 18 produces a monic $P_f(t) \in \mathbb{Z}[t]$ such that $P_f(f^*)$ annihilates $\text{Pic}(X)$. Indeed, $\text{Pic}^0(X)$ is an abelian variety (use the characteristic polynomial on a Tate module), while $\text{NS}(X)$ is finitely generated. For an ample H , the cyclic space

$$V = \text{Span}_{\mathbb{Q}}\{H, f^*H, \dots, (f^*)^{\deg P_f - 1}H\} \subset \text{Pic}(X)_{\mathbb{Q}}$$

is therefore finite-dimensional and f^* -invariant. Put $f^*|_V$ in Jordan form.

For the blocks with $|\lambda| > 1$, use Ingredient I. Choose the first canonical-height coordinate that is nonzero at x . Its exact iterate formula gives the lower bound $\underline{\alpha}_f(x) \geq |\lambda|$; Lemma 17 permits this comparison with an arbitrary complex divisor height. Expressing H in the Jordan basis gives the reverse bound: all other terms are at most a polynomial times $|\lambda|^n$, while blocks of modulus at most one contribute only $O((1+\varepsilon)^n)$. Hence $\alpha_f(x) = |\lambda|$. If every expanding canonical-height coordinate vanishes, the same upper bound gives $\alpha_f(x) = 1$.

Consequently every arithmetic degree lies in $\{1\}$ together with the moduli of finitely many eigenvalues of $f^*|_V$. Those eigenvalues are roots of the monic integral polynomial P_f , yielding both integrality and finiteness.

Ingredient III: nef heights on abelian varieties

For an abelian variety $A/\overline{\mathbb{Q}}$ and a nonzero nef $D \in \text{Div}(A)_{\mathbb{R}}$, Theorem 4 identifies a unique proper abelian subvariety $B_D \subsetneq A$ such that

$$\{x : \widehat{q}_{A,D}(x) = 0\} = B_D(\overline{\mathbb{Q}}) + A(\overline{\mathbb{Q}})_{\text{tors}}. \quad (3)$$

Fix an ample H . Under the Rosati identification, nefness makes Φ_D positive semidefinite, so Proposition 25 factors $\Phi_D = \gamma'\gamma$ with $\gamma' = \gamma \in \text{End}(A)_{\mathbb{R}}$. The height pairings then give $\widehat{q}_{A,D}(x) = \widehat{q}_{A,H}(\gamma x)$. Because γ need not be an algebraic endomorphism, Lemma 29 is essential: on $A(\overline{\mathbb{Q}})_{\mathbb{Q}}$ its kernel is the common kernel of finitely many rational endomorphisms. Taking the connected component and then the divisible hull yields exactly (3).

Finally, a Perron–Frobenius argument supplies a nonzero nef eigenclass F with $f^*F \equiv \delta_f F$. If the orbit of x is Zariski dense, (3) and finite generation over a number field force $\widehat{q}_{A,F}(x) > 0$; its functional equation gives $\alpha_f(x) \geq \delta_f$, while the general upper bound gives equality (Corollary 31).

Scope and bottleneck

The theorem concerns morphisms, not rational self-maps with indeterminacy. The clean $O(1)$ normalization in (2) uses linear equivalence; the paper only states, without proof, a weaker algebraic-equivalence variant. The hard conceptual step is separating polynomial Jordan growth from its exponential rate while controlling every ample height through one finite Picard-space calculation.