Obstruction classes for moduli spaces of sheaves and Lagrangian fibrations

Dominique Mattei, Reinder Meinsma

Dominique Mattei and Reinder Meinsma, arXiv:2404.16652v2 (2025), 30 pages
https://arxiv.org/abs/2404.16652v2

The universal-sheaf obstruction

Let S be a complex projective K3 surface. Its Mukai lattice is

$$\widetilde{H}(S,\mathbf{Z}) = H^0(S,\mathbf{Z}) \oplus H^2(S,\mathbf{Z}) \oplus H^4(S,\mathbf{Z}), \qquad (r,c,s)\cdot(r',c',s') = c\cdot c' - rs' - r's,$$

and its algebraic part is $N(S) = H^0(S,\mathbf{Z}) \oplus \mathrm{NS}(S) \oplus H^4(S,\mathbf{Z})$. For a primitive Mukai vector $v \in N(S)$ and a v-generic polarization H, write $M = M_H(v)$ for the moduli space of H-Gieseker-stable sheaves of vector v. It is generally only a coarse moduli space. There is, however, a unique class $\alpha \in \mathrm{Br}(M)$ for which a $(1\boxtimes\alpha)$-twisted universal sheaf exists on $S\times M$; M is fine exactly when $\alpha = 0$.

Set

$$d = \mathrm{div}_{N(S)}(v) := \gcd\{v\cdot x : x \in N(S)\}.$$

The paper extends Căldăraru's calculation for isotropic Mukai vectors to the higher-dimensional case.

Main theorem (Mattei–Meinsma, Theorems 1.2 and 4.5)**.** *Let $v \in N(S)$ be primitive with $v^2 > 0$, let H be a v-generic polarization, and put $M = M_H(v)$. There is a short exact sequence*

$$0 \longrightarrow \langle\alpha\rangle \longrightarrow \mathrm{Br}(M) \longrightarrow \mathrm{Br}(S) \longrightarrow 0,$$

where α is the obstruction class. Moreover, $\mathrm{ord}(\alpha) = d$*. Consequently M is fine if and only if $d = 1$ (Corollary 4.6).*

Thus the theorem does more than detect non-fineness: it measures the entire discrepancy between the two Brauer groups by one explicitly controlled cyclic class.

Ingredient 1: the lattice defect

For a hyperkähler manifold of $\mathrm{K3}^{[n]}$-type the group $H^2(M,\mathbf{Z})$ need not be unimodular, so one cannot simply identify its Brauer group with $\mathrm{Hom}(T(M),\mathbf{Q}/\mathbf{Z})$. The correct object is

$$T'(M) := H^2(M,\mathbf{Z})/\,\mathrm{NS}(M), \qquad \mathrm{Br}(M) \simeq T'(M)\otimes\mathbf{Q}/\mathbf{Z}.$$

The Mukai Hodge isometry $v^\perp \simeq H^2(M,\mathbf{Z})$ induces the central lattice calculation (Lemma 4.1)

$$0 \longrightarrow T'(M) \longrightarrow T(S)^* \longrightarrow \mathbf{Z}/d\mathbf{Z} \longrightarrow 0.$$

If $u \in \widetilde{H}(S,\mathbf{Z})$ satisfies $u\cdot v \equiv 1 \pmod d$ and $w \in T'(M)$ maps to $du \in T(S)^*$, then the resulting Brauer sequence has kernel generated by w/d. At this stage w/d is only the lattice-theoretic candidate for the geometric obstruction.

Ingredient 2: a canonical Fourier–Mukai class

A quasi-universal sheaf of similitude ρ gives a normalized cohomological Fourier–Mukai morphism $\varphi : \widetilde{H}(S,\mathbf{Q}) \to H^*(M,\mathbf{Q})$. Lemma 4.3 shows that the degree-two component $[\varphi(u)]_2$ maps to w/d in $\mathrm{Br}(M)$. This class is independent of both choices: changing the quasi-universal sheaf changes $[\varphi(u)]_2$ by a rational Néron–Severi class, while changing u changes it by integral and algebraic classes. This is the bridge from the cyclic lattice quotient to a class that can be compared with α.

Ingredient 3: deform to a fine fibre

Proposition 3.17 embeds the data in a smooth connected family $\mathcal{S} \to T$ with constant integral cohomology and compatible v-generic polarizations. The original surface is one fibre, while another fibre S_1 satisfies $\mathrm{div}_{N(S_1)}(v) = 1$. The associated relative moduli space $\mathcal{M} \to T$ carries a relative twisted universal sheaf, but the obstruction vanishes over S_1 because that fibre is fine.

Choose an α^{-1}-twisted locally free sheaf F of rank ρ on $\mathcal{M}$, and restrict the twisted universal sheaf $\mathcal{E}$ and F to $\mathcal{E}_t, F_t$ on M_t. Put $U_t = \mathcal{E}_t \otimes p^*F_t$; let φ_t be its normalized cohomological transform. On the fine fibre $\mathcal{E}_1$ is an ordinary universal sheaf; write Φ_1', φ_1' for its Fourier–Mukai and cohomological transforms. Then

$$\varphi_1(x) = \varphi_1'(x)\frac{\mathrm{ch}(F_1^\vee)}{\rho}, \qquad \alpha|_M = \frac{c_1(F_1)}{\rho},$$

the second equality being Căldăraru's deformation formula.

Here the two choices of Mukai class must be separated. Since the full Mukai lattice is unimodular and v is primitive, choose $u_0 \in \widetilde{H}(S, \mathbf{Z})$ with $u_0 \cdot v = 1$ exactly. On S_1 the paper constructs a line bundle L with

$$u_1 := v(L), \qquad u_1 \cdot v \equiv 1 \pmod{d};$$

it does not assert equality. Replace L by a sufficiently positive twist and reset $u_1 = v(L)$ (the congruence is preserved); boundedness then makes $G := \Phi_1'(L)$ locally free, so $[\varphi_1'(u_1)]_2 = c_1(G)$ is integral. Because $\mathrm{div}_{N(S_1)}(v) = 1$, choose $x_1 \in N(S_1)$ with $x_1 \cdot v = (u_1 - u_0) \cdot v$. Then $z = u_1 - u_0 - x_1 \in v^\perp$; the term $[\varphi_1'(z)]_2$ is integral by the Mukai isometry, while $[\varphi_1'(x_1)]_2$ is algebraic. Thus u_0 and u_1 define the same zero Brauer class on M_1. This is the independence mechanism of Lemma 4.3, which also identifies $[\varphi_0(u_0)]_2$ with w/d on the original fibre.

Now the exact pairing may legitimately be used: $[\varphi_1'(u_0)]_0 = -u_0 \cdot v = -1$. Taking degree two in the displayed transform identity, discarding the integral/algebraic term, and transporting through the family gives $[\varphi_0(u_0)]_2 = c_1(F_1)/\rho = \alpha$. Hence the lattice generator w/d is precisely the obstruction.

A numerical model and the hard point

Suppose $\mathrm{NS}(S) = \mathbf{Z}L$ with $L^2 = 2m > 0$ and take $v = (2, L, 0)$. Then v is primitive, $v^2 = 2m$, and $\mathrm{div}_{N(S)}(v) = \gcd(2, 2m) = 2$. For a generic polarization the obstruction therefore has exact order two: a twisted universal sheaf exists, but an ordinary universal sheaf does not. By contrast, any primitive positive-square vector of divisibility one gives a fine moduli space.

The technically delicate step is not the final gcd computation but the deformation that keeps v algebraic and the polarization generic while reaching a divisibility-one fibre, together with the exact-versus-congruent comparison above.

Lagrangian-fibration applications (statements only)

The following records the paper's other headline results; unlike the obstruction theorem above, their proofs are not unpacked here. Let $H \subset S$ be a smooth irreducible curve with primitive class and $H^2 = 2g - 2 \geq 2$, put

$$v_k = (0, H, k + 1 - g), \qquad P_k = \overline{\mathrm{Pic}}^{\,k}(\mathcal{C}/|H|) = M(v_k).$$

The support map $P_k \to |H|$ is a Beauville–Mukai Lagrangian fibration. Proposition 5.6 gives a short exact sequence

$$0 \longrightarrow C_k \longrightarrow \mathrm{Sha}(S, H) \longrightarrow \mathrm{Br}(P_k) \longrightarrow 0, \qquad |C_k| = \frac{\mathrm{div}(H)}{\gcd(\mathrm{div}(H), k + 1 - g)},$$

where $C_k = A/\ker(\zeta_H)$ in the paper's notation. Under this surjection, Proposition 5.8 identifies the torsor class $[P_1]$ with the universal-sheaf obstruction $\alpha_k \in \mathrm{Br}(P_k)$.

If $U \subset \mathrm{NS}(S)$ (equivalently, S admits an elliptic fibration with a section), Theorem 5.15 says that P_k and P_ℓ are birational exactly when their obstruction classes have the same order, equivalently

$$\gcd(\mathrm{div}(H), k + 1 - g) = \gcd(\mathrm{div}(H), \ell + 1 - g).$$

In the same setting, Theorem 5.13 says that a $2n > 2$-dimensional moduli space is fine exactly when it is birational to $S^{[n]}$, and birationality to $S^{[n]}$ implies derived equivalence. The converse derived-Torelli direction fails broadly: Theorem 5.17 constructs, for every $n \geq 3$, two $2n$-dimensional moduli spaces of sheaves on the same K3 surface whose transcendental lattices are Hodge isometric but whose derived categories are not equivalent. The positive-square theorem above does not cover the isotropic case $v^2 = 0$, which is earlier work recalled in the paper.