Arithmetic degrees and Zariski dense orbits of cohomologically hyperbolic maps

Yohsuke Matsuzawa, Long Wang

arXiv:2212.05804

http://arxiv.org/abs/2212.05804v2

Arithmetic growth from a cohomological gap

Let X be a smooth projective variety over $\overline{\mathbf{Q}}$ and $f\colon X \dashrightarrow X$ a dominant rational self-map. Write $X_f(\overline{\mathbf{Q}})$ for the points whose forward orbit avoids the indeterminacy locus. For an ample divisor H, the lower arithmetic degree is

$$\underline{\alpha}_f(x) = \liminf_{m\to\infty} \max\{1, h_H(f^m(x))\}^{1/m}.$$

The dynamical degrees $\delta_i(f)$ measure the exponential growth of codimension-i intersections. The map is p-cohomologically hyperbolic if $\delta_p(f) > \delta_i(f)$ for every $i \neq p$, $1 \leq i \leq \dim X$. Put $N = \dim X$ and use the boundary conventions $\delta_0(f) = \delta_{N+1}(f) = 1$.

An orbit is called *generic* when it is infinite and meets every proper closed subset only finitely. This is stronger than merely being Zariski dense.

Main theorem (Matsuzawa–Wang, Theorem A and Corollary 1.8)**.** *If f is p-cohomologically hyperbolic and $x \in X_f(\overline{\mathbf{Q}})$ has generic orbit, then*

$$\underline{\alpha}_f(x) \geq \frac{\delta_p(f)}{\delta_{p-1}(f)}.$$

Moreover, there is a sequence $x_i \in X_f(\overline{\mathbf{Q}})$, each satisfying the dynamical Mordell–Lang property, for which $\underline{\alpha}_f(x_i)$ converges and

$$\lim_i \underline{\alpha}_f(x_i) \geq \frac{\delta_p(f)}{\delta_{p-1}(f)}.$$

In particular, if $p = 1$, then every generic orbit has an arithmetic degree and $\alpha_f(x) = \delta_1(f)$; the sequence above converges to $\delta_1(f)$.

For $p = 1$ the paper also proves that such points exist: some $x \in X_f(\overline{\mathbf{Q}})$ has Zariski-dense orbit and $\alpha_f(x) = \delta_1(f)$ (Theorem B).

Ingredient 1: a big divisor produces a height recurrence

Resolve simultaneously the indeterminacy of f^n and f^{2n}. On the resulting smooth model, Proposition 3.5 combines log-concavity of the dynamical degrees with a volume estimate to prove that, for every $\varepsilon \in (0,1)$ and all sufficiently large n, a divisor built from the pullbacks of H at times $0, n, 2n$ is big. Outside its augmented base locus, functoriality of heights gives the concrete inequality

$$\frac{h_H(f^{2n}(z))}{\delta_{p+1}(f)^n} + \frac{h_H(z)}{\delta_{p-1}(f)^n} \geq \frac{\varepsilon^n \delta_p(f)^n}{\delta_{p-1}(f)^n \delta_{p+1}(f)^n} h_H(f^n(z)) - C_n. \tag{1}$$

This is the key replacement for the height functoriality that a rational, rather than regular, self-map lacks.

Ingredient 2: solve the recurrence along a generic orbit

Put $g = f^n$. Cohomological hyperbolicity makes the middle coefficient in (1) eventually larger than the sum of the two reciprocal endpoint coefficients. Lemma 3.1 then turns (1) into

$$\underline{\alpha}_{f^n}(y) \geq \frac{\varepsilon^n \delta_p(f)^n}{2\delta_{p-1}(f)^n}$$

whenever the full f^n-orbit of y remains in the relevant open set. A generic orbit is eventually contained in every such open set. Taking n-th roots, then $n \to \infty$ and $\varepsilon \to 1$, removes the factor $2^{1/n}$ and proves the theorem's ratio.

This recurrence is also the useful toy model: if three successive heights at times $0, n, 2n$ obey (1), the dominant characteristic growth is forced by δ_p/δ_{p-1}; the geometric work is precisely to make that one-dimensional comparison valid off a proper closed set.

Ingredient 3: adelic points supply DML orbits

Proposition 3.2, using Xie's adelic-topology results, says that for any nonempty open $U \subset X$ and any infinite-order f, a nonempty adelic open set of points has its entire orbit in U, has infinite orbit, and satisfies dynamical Mordell–Lang. Applying this to the open sets from (1) gives the sequence x_i. The DML property matters because a Zariski-dense DML orbit is automatically generic.

Ingredient 4: growth forces the orbit closure to be all of X

For a point supplied by the adelic construction, let Z be a periodic irreducible component of its orbit closure and let $c = \operatorname{codim}_X Z$. On a suitable open set an iterate of f is quasi-finite at the generic point of Z. If g is the induced map on Z, the iterate and restriction formulas (Lemmas 2.7 and 2.8), the standard arithmetic-degree upper bound (Proposition 2.6), and the restriction comparison (Lemma 2.3) give

$$\underline{\alpha}_f(x)^k \leq \delta_1(g) \leq \delta_{1+c}(f)^k.$$

For $p = 1$, the constructed lower bound is arbitrarily close to $\delta_1(f)$, while $\delta_2(f) < \delta_1(f)$. Hence $c = 0$, proving Theorem B. Applying this to a plane map with transcendental $\delta_1(f) > \delta_2(f) = 5$ yields Theorem D: arithmetic degrees of rational maps need not be algebraic.

The large-topological-degree branch

The paper also treats N-cohomologically hyperbolic maps. Theorem E says that if x has Zariski-dense orbit and that orbit meets every closed subset of codimension at least two only finitely, then $\underline{\alpha}_f(x) \geq \delta_N(f)/\delta_{N-1}(f)$. Theorem F produces a nonempty open $U \subset X$ for which the preperiodic points $x \in U(\overline{\mathbf{Q}}) \cap X_f(\overline{\mathbf{Q}})$ whose entire orbit stays in U form a set of bounded height. These results use a top-degree height inequality together with weak dynamical Mordell–Lang.

Hard point and scope

The hardest step is the bigness calculation behind (1): on a common resolution, mixed intersection terms must be controlled uniformly by degrees of iterates, and log-concavity must leave exactly the required spectral gap. The conclusion for an arbitrary prescribed orbit assumes *genericity*, not only Zariski density. The separate existence argument manufactures DML points so that this stronger condition follows; the paper does not assert that every Zariski-dense orbit of an arbitrary rational map is generic.