

Canonical heights and preperiodic points for certain weighted homogeneous families of polynomials

Patrick Ingram

arXiv:1510.08807

<http://arxiv.org/abs/1510.08807v3>

A height gap in a non-isotrivial family

Let K be a number field. A polynomial family is *homogeneous of weight e* if $e \geq 2$ and

$$f_t(z) = F(z^e, t),$$

where $F(X, Y)$ is a homogeneous binary form divisible by neither X nor Y ; “weighted homogeneous” means homogeneous for some e . The model example is $f_t(z) = z^d + t$ (with $e = d$). Write $\hat{h}_{f_t}$ for the canonical height, $h_{M_d}(f_t)$ for the height of the moduli point, and $\Re_{f_t}$ for the minimal resultant.

Main theorem (Ingram, Theorem 1). *For every $s \geq 0$ there is $\varepsilon > 0$, depending on K , the family, and the chosen ample class on M_d , such that if f_t has at most s places of bad reduction and $z \in K$ is not preperiodic, then*

$$\hat{h}_{f_t}(z) \geq \varepsilon \max\{h_{M_d}(f_t), \log |\text{Norm } \Re_{f_t}|\}. \quad (1)$$

This verifies, for these families and with bounded bad support, the dynamical analogue of Lang’s height-gap conjecture. It also explains why height lower bounds and uniform bounds for rational preperiodic points are linked: a nonzero point of very small canonical height is ruled out uniformly once the parameter is arithmetically complicated.

Ingredient I: local escape and tiny residue tubes

After a finite extension and a fixed linear conjugacy, the family may be taken monic without changing canonical heights. It then factors as

$$f_t(z) = \prod_{i=1}^{d/e} (z^e - \beta_i t).$$

For each place v , Lemma 5 gives constants independent of t such that outside the scale $|z|_v^e \asymp |t|_v$ one has $G_{f_t, v}(z) = \log |z|_v + O_v(1)$. More sharply, outside a fixed finite set of places,

$$e \nmid v(t) \implies G_{f_t, v}(z) \geq \frac{1}{d} \log^+ |t|_v \quad (2)$$

for every $z \in K$. Thus a preperiodic point can exist only when $|z|_v^e = |t|_v$ at such a denominator place.

The local Arakelov–Green function

$$g_{f_t, v}(x, y) = -\log |x - y|_v + G_{f_t, v}(x) + G_{f_t, v}(y)$$

measures how strongly two orbit points are forced together relative to their escape. Lemma 7 is the key combinatorial estimate. Given at least $d + 3$ orbit indices, either two iterates coincide, or a fixed positive fraction of the indices has pairwise

$$g_{f_t, v}(f_t^i(z), f_t^j(z)) \geq \delta \log^+ |t|_v - c_v, \quad (3)$$

where $\delta = \min\{1/e, 1 - 1/e - 1/d\} > 0$ when $d > e$. The proof partitions the nonescaping iterates among at most d small tubes around the roots of $z^e = \beta_i t$, plus one escaping class. Two points in the same tube are v -adically so close that the $-\log |x - y|_v$ term supplies (3).

Ingredient II: one pair works at every bad place

Let S contain the archimedean places and all places with $|t|_v > 1$. Start with a sufficiently long initial orbit segment. Apply Lemma 7 successively at every $v \in S$, retaining at least a factor $1/(d+2)$ of the indices each time. Unless an equality of iterates has already proved preperiodicity, two indices $i \neq j$ survive and satisfy (3) simultaneously at every place in S . Lemma 6 supplies a uniform lower bound at all remaining places.

Summing over all v now uses the product formula to cancel $-\log |f_t^i(z) - f_t^j(z)|_v$. Since $\widehat{h}_{f_t}(f_t^m(z)) = d^m \widehat{h}_{f_t}(z)$, the result is

$$\widehat{h}_{f_t}(z) \geq \varepsilon_S h(t) - C_S. \quad (4)$$

The constants deteriorate exponentially with $\#S$, which is exactly why Theorem 1 fixes a bound on the bad-reduction support.

Ingredient III: parameter height controls the target

Lemma 8 closes the gap between (4) and (1). Functoriality of heights for the curve $t \mapsto [f_t] \in \mathcal{M}_d$ gives $h_{\mathcal{M}_d}(f_t) \ll h(t)$. Writing an integral model explicitly also bounds the valuation of its resultant and yields

$$\log |\text{Norm } \mathfrak{R}_{f_t}| \ll h(t), \quad (5)$$

with family-dependent constants. The finitely many small-height parameters and the case $d = e$ are handled separately (the latter is the previously known family $z^d + bt$). Equations (4)–(5), together with the orbit-repetition alternative, prove the stated height gap.

A local toy model

Take $f_t(z) = z^d + t$ and a nonarchimedean v with $|t|_v > 1$ and $d \nmid v(t)$. Because $v(z) \in \mathbb{Z}$, equality $|z|_v^d = |t|_v$ is impossible. If $|z|_v^d > |t|_v$, the leading term dominates and the orbit escapes. If $|z|_v^d < |t|_v$, then $|f_t(z)|_v = |t|_v$, after which the leading term dominates and the orbit again escapes. This is (2) in its cleanest form. The general proof replaces the single boundary circle by finitely many very small root tubes and must choose the same pair of orbit points at every bad place.

Generic base change and uniform boundedness

The paper also shows how to remove the bound on $\#S$ after a suitably generic base change. Put $N_e = 5, 4, 3$ for $e = 2, 3, \geq 4$, respectively, and call $\varphi: \mathbb{P}_K^1 \rightarrow \mathbb{P}_K^1$ e -general when it has at least N_e distinct K -rational finite poles (poles in $\mathbb{A}^1(K)$) whose orders are prime to e . Assuming the *abc* conjecture for K , Theorem 2 proves an absolute version of (1) for $t = \varphi(s)$. The point is that *abc* forces a positive proportion of $h(\varphi(s))$ to come from denominator valuations not divisible by e , where (2) applies.

Unconditionally, Lemma 9 and the Darmon–Granville theorem show that, outside finitely many $s \in \mathbb{P}^1(K)$, some such valuation must occur. Lemma 5(C) then rules out every finite K -rational preperiodic point. This proves Theorem 3 and hence uniform boundedness for the base-changed family, although over the thin parameter set $\varphi(\mathbb{P}^1(K))$.

Hard point and scope

The decisive step is the iterated local pigeonhole argument: local escape alone gives a good pair at one place, whereas the product formula needs one pair that is good at all bad places. The shrinking of the index set records the precise cost of synchronizing those local choices.

The theorem concerns weighted homogeneous polynomial families over a number field; it is not a height gap for arbitrary rational maps. The unrestricted base-change height inequality is conditional on *abc*, while the finiteness of parameters carrying rational preperiodic points is unconditional. The paper proves existence of effective-looking constants but does not make the final uniform bound numerically sharp.