

Singularities of Foliations and Good Moduli Spaces of Algebraic Stacks

Federico Bongiorno

arXiv:2603.24997

<http://arxiv.org/abs/2603.24997v1>

From stack singularities to quotient geometry

Let $\mathcal{X}$ be a normal algebraic stack of finite type over an algebraically closed field k of characteristic zero, with affine diagonal and finite generic stabilizers. A smooth presentation $p : X \rightarrow \mathcal{X}$ has relative tangent foliation

$$\mathcal{F} = T_{X/\mathcal{X}} \subset T_X. \quad (1)$$

The stack is *relatively (log-)canonical* at x when $\mathcal{F}$ is Gorenstein there and every divisorial valuation centered over a lift of x has nonnegative (log-)discrepancy. Smooth ascent and descent make this independent of the chosen presentation (Definition 2.11).

Main theorem (Bongiorno, Main Theorem). *For $x \in \mathcal{X}(k)$:*

1. *if $\mathcal{X}$ is relatively log-canonical at x , its stabilizer is a finite extension of an algebraic torus, and $\mathcal{X}$ admits a good moduli space étale locally at x ;*
2. *if $\mathcal{X}$ is relatively canonical at x , the stable locus for that local good moduli space is nonempty.*

Here a point is stable for $q : \mathcal{U} \rightarrow Q$ when its moduli fiber is just that point. The result makes an MMP-type discrepancy condition control two quotient-theoretic features: reductivity of stabilizers and, under the stronger canonical condition, existence of closed generic orbits.

Semistable reduction turns the foliation into a torus model

Finite generic stabilizers first give a dense smooth Deligne–Mumford gerbe $\mathcal{U}^\circ \rightarrow B^\circ$ over a smooth affine scheme. This is a rational first integral $\mathcal{X} \dashrightarrow B$. Resolving its indeterminacy and applying functorial semistable reduction produce a logarithmic alteration

$$\pi : (\mathcal{Z}^{\text{ss}}, \Delta^{\text{ss}}) \longrightarrow \mathcal{X}, \quad g^{\text{ss}} : (\mathcal{Z}^{\text{ss}}, \Delta^{\text{ss}}) \longrightarrow (B^{\text{ss}}, \Delta_B^{\text{ss}}). \quad (2)$$

On an atlas Z^{ss} , the pulled-back presentation foliation $\mathcal{F}^{\text{ss}}$ and the logarithmic relative tangent sheaf satisfy

$$\mathcal{F}^{\text{ss}} \subset T_{g^{\text{ss}}} \quad (3)$$

with the same generic rank. Semistability makes $T_{g^{\text{ss}}}$ log-regular and gives its stabilizer Lie algebra a faithful semisimple normal representation (Proposition 2.10). Log-canonical singularities persist under the alterations. Off the logarithmic boundary, the point lies over the relatively regular gerbe locus, and Lemmas 2.12, 2.13, and 2.5 upgrade (3) to equality. On the boundary, log-canonicity and the equal-rank criterion of Lemma 2.8 give the same equality; a strict inclusion would expose a divisor with negative log-discrepancy. Proposition 3.9 is the technical core: after alteration, the original infinitesimal stack directions are exactly those of a semistable, hence toroidal, map.

Why additive stabilizers are impossible

Let G be the stabilizer of x and suppose its identity component contains $\mathbb{G}_a$. The group acts on the proper fiber $V = \pi^{-1}(x)$. Lemmas 4.1–4.2 ensure that this additive subgroup fixes a point $v \in V$: pass to the proper coarse space, choose an orbit of minimal dimension, and use that a nontrivial $\mathbb{G}_a$ -orbit cannot be both affine and proper.

At v , differentiating the $\mathbb{G}_a$ -action gives a nilpotent endomorphism in the normal representation of $\mathcal{F}^{\text{ss}}$. By (3) and Proposition 3.9, this is the faithful semisimple normal representation of $T_{g^{\text{ss}}}$. Semisimplicity forces the nilpotent operator to vanish, while faithfulness forbids that vanishing. Thus G° contains no $\mathbb{G}_a$; Lemma 4.3 says G° is a torus, so G is a finite extension of a torus. The Alper–Hall–Rydth slice theorem then supplies the claimed étale-local good moduli space.

What the canonical strengthening adds

Canonical singularities control the exceptional boundary more sharply. For the resolved map $g : (\mathcal{Z}, \Delta_{\mathcal{Z}}) \rightarrow B$, Proposition 3.6 proves that $g(\Delta_{\mathcal{Z}})$ is not dense: if a boundary valuation dominated the generic point of B , the relative derivation sending a uniformizer to 1 would contradict invariance of that divisor. Over a dense open $W \subset B$ disjoint from the boundary image, $g^{-1}(W) \rightarrow W$ is a gerbe. A closed $w \in W$ therefore has a unique closed point z' upstairs. Moreover $g^{-1}(W)$ lies in the original Deligne–Mumford gerbe locus $\mathcal{U}^\circ$, where the modification is an isomorphism. Thus $x' = \sigma(z')$ has finite stabilizer, and properness makes it closed. In this local good-moduli quotient, a closed point with the minimal stabilizer dimension (here zero) is stable: its moduli fiber is the singleton $\{x'\}$. Hence the stable locus is nonempty.

For a smooth presentation, Corollary B consequently identifies $\mathcal{F}$ generically, after an étale localization, with $T_{U/Q}$ for an affine quotient $U \rightarrow Q$. This is an étale-local form of the expected quotient for algebraically integrable canonical foliations.

The two weight models and the limits

Let $\mathbb{G}_m$ act on $\mathbb{A}^2$. With weights $(1, 1)$, the vector field $x\partial_x + y\partial_y$ is log-canonical at the origin, but generic orbits are not closed; $[\mathbb{A}^2/\mathbb{G}_m] \rightarrow \text{Spec } k$ has no stable point. With weights $(1, -1)$, the vector field $x\partial_x - y\partial_y$ is canonical, the invariant xy gives the good moduli space $[\mathbb{A}^2/\mathbb{G}_m] \rightarrow \mathbb{A}^1$, and the locus $xy \neq 0$ is stable. The examples show why log-canonicity yields a good quotient while canonicity is needed for stability.

The theorem is local in the étale topology and does not construct a global projective quotient. Its characteristic-zero, affine-diagonal, and finite generic-stabilizer hypotheses are essential inputs to the alteration, semistable-reduction, and slice arguments. Corollary B applies to foliations arising from stack presentations; it is not a general global solution of the Cascini–Spicer conjecture.