



< Previous

✓

✓

✓

✓

✓

✓

✓

✓

✓

Next >

## 8. Kolmogorov-Smirnov Test Statistic Pivotal Under Null

[🔖](#) Bookmark this page

[Start of transcript. Skip to the end.](#)



So It turns out that this result is also true.  
So what did we want?  
Why did we use the Donsker's theorem?  
Because Donsker's theorem tells us  
that no matter what the true  $f$  is,  
asymptotically,  
or even what  $f_0$  is, asymptotically,  
square root of  $n$ ,  $f_n$  minus  $f$  not, if  $f$  not  
equal to  $f$ ,  
will converge to something which is

▶

0:00 / 0:00

▶

1.50x

🔊

🔍

CC

🗣️

Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

### CDF as a Random Function

3 points possible (graded)  
Let  $X$  be a random variable with invertible cdf  $F_X$ . Define another random variable  $Y = F_X(X)$ . Find the cdf  $F_Y$  of  $Y$ .

For  $t < 0$ :

$F_Y(t) =$

For  $t \geq 1$ :

$F_Y(t) =$

For  $0 \leq t < 1$ :

$F_Y(t) =$

(What is the distribution of  $Y$ ?)

STANDARD NOTATION

Submit

You have used 0 of 3 attempts

Let  $X_1, \dots, X_n$  be iid samples with unknown cdf  $F_X$ . For simplicity, restrict to the cases when  $F_X$  is invertible.

Recall the goal of the Kolmogorov-Smirnov Test goodness of fit test is to decide between the hypotheses

$$\begin{aligned} H_0 &: F_X = F^0 \\ H_1 &: F_X \neq F^0. \end{aligned}$$

Recall also the Kolmogorov-Smirnov test statistic:

$$T_n = \sqrt{n} \sup_{t \in \mathbb{R}} |F_n(t) - F^0(t)|$$

Assuming  $H_0$  is true, then  $T_n$  becomes

$$T_n = \sqrt{n} \sup_{t \in \mathbb{R}} |F_n(t) - F_X(t)|$$

We will see that under the null hypothesis, the distribution of  $T_n$  does not depend on the distribution of the data  $X_i$ , i.e.  $T_n$  is pivotal, and this is true for any  $n$ , not only for large  $n$ .

The trick is to make a change of variables. Let  $\tilde{t} = F_X(t)$ , then  $t = F_X^{-1}(\tilde{t})$ . We have

$$\begin{aligned} T_n &= \sqrt{n} \sup_{t \in \mathbb{R}} |F_n(t) - F_X(t)| \\ &= \sqrt{n} \sup_{t \in \mathbb{R}} \left| \left( \frac{1}{n} \sum_{i=1}^n \mathbf{1}(X_i \leq t) \right) - F_X(t) \right| \quad (\text{definition of empirical cdf}) \\ &= \sqrt{n} \sup_{t \in \mathbb{R}} \left| \left( \frac{1}{n} \sum_{i=1}^n \mathbf{1}(F_X(X_i) \leq F_X(t)) \right) - F_X(t) \right| \quad (\text{apply } F_X \text{ to both sides of inequality}) \\ &= \sqrt{n} \sup_{\tilde{t} \in (0,1)} \left| \left( \frac{1}{n} \sum_{i=1}^n \mathbf{1}(Y_i \leq \tilde{t}) \right) - \tilde{t} \right| \quad \text{where } Y_i \sim \text{Unif}(0,1). \end{aligned}$$

## Discussion

Hide Discussion

**Topic:** Unit 4 Hypothesis testing:Lecture 16: Goodness of Fit Tests Continued: Kolmogorov-Smirnov test, Kolmogorov-Lilliefors test, Quantile-Quantile Plots / 8. Kolmogorov-Smirnov Test Statistic Pivotal Under Null

Add a Post

Show all posts	by recent activity
<div>Any hint for the excersice?</div>	4

< Previous

Next >

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

---

## Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

---

## Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)

