

My M&M OCD

Yoni

13 04, 2025

Intro

Objective of Simulation

The goal of this simulation is to test the statistics of M&M and other stacks even Chocolate lentils by color.

I wanted to know what is the chance of my finishing the package of M&M without mixing any color in one bite, eating them 2 by 2

In addition, here are some BI incite that needed to be checked:

1. What is the probability of M&M packages packaged fairly?
2. What is the probability of M&M packages packaged without one color?
3. How does the size of the package or number of colors affect this probability?

Method

I do not have an inner information of how does (M&M)[<https://www.mms.com/en-us>] make their delicious snacks nor we know how they make sure each package have fair amount of each color. Therefore, the method I chose is based of simulation of some M&M bags, according to the most common sizes of packages.

Each time we sample x lentils(units of M&M), name them by colors (V1,V2...), and see the results for many packages as a statistic data.

my hypothesis is that the probability of perfect package (aka a package with all colors number been even) is very small, at least for a standard 6 colors pack.

Parameters

Basic parameters:

```
#parameters
n<- 800           #numbers of bags per sample
n_color<- 6       #unique colors of M&M
gram<- 0.91       #weight of one M&M
bag_g<- 250       #common weight of M&M package
n_unit<- bag_g/gram #M&M per packagenm,
av_per_color= n_unit/n_color
paste0("The avarage number of lentils per color is ", round(av_per_color,2))
```

```
## [1] "The avarage number of lentils per color is 45.79"
```

Creating of the Sample

General Sample

In order to test the theoretical data, we need to simulate it using customize functions. here are there:

- Create_bag- function to create one snack package for chosen package size and number of colors.
- sample_MnM- function to create n bags from the Create_bag function.

```
## [1] "One bag of 100:"
```

```
##      1  2  3  4  5  6
## [1,] 14 18 12 20 14 23
```

```
## [1] "3 bags of 100:"
```

```
##      Red Blue Green Orange Yellow Brown
## Bag_1  20  16   17   14   16   18
## Bag_2  16  23   11   16   16   18
## Bag_3  14  14   18   21   15   18
```

Preview Graph

Now will be creating n bugs of M&M
columns:

1. V1:V6- the number of lentils per color
2. even_count- how many evens colors there are
3. even_evens- are the uneven colors even
4. Variance- variance of lentils per color
5. low_col- sum true if one color's count is lower than $\frac{2}{3}$ of expected value
6. min- the lowest color in each row


here are the first rows:


```
## # A tibble: 6 x 12
##   Red   Blue Green Orange Yellow Brown even_count even_evens low_col Variance
##   <int> <int> <int>  <int>  <int>  <int>      <dbl> <lgl>      <dbl>    <dbl>
## 1    57    42    41    45    47    43         1 FALSE         0     34.6
## 2    34    58    48    43    46    46         5 FALSE         0     60.2
## 3    43    45    51    40    44    51         2 TRUE          0     19.9
## 4    47    44    45    47    38    53         2 TRUE          0     23.9
## 5    63    48    39    47    40    38         3 FALSE         0     88.6
## 6    46    39    54    47    42    47         3 FALSE         0     26.2
## # i 2 more variables: min <int>, all_even <lgl>
```

plot the M&M sample sample

```
## [1] "summary of all colors Distribution:"
```

##	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	Var
## Red	28	41	46	45.58875	50	66	37.84693
## Blue	30	42	46	45.91125	50	67	37.18986
## Green	30	41	46	45.62125	50	69	36.92896
## Orange	28	42	46	45.95500	50	67	36.96668
## Yellow	27	42	46	45.67875	50	64	37.41482
## Brown	27	41	46	45.78375	50	65	39.91438





Statistics Checking of the Simulation

Test Expected Value


to see is the μ of the lentils per color are fair, we will test it per column with t.test for each color.

Here is the result, none of them bellow 5% P. value

```
##      Red      Blue      Green      Orange      Yellow      Brown
## "36.10%" "56.63%" "43.92%" "43.62%" "61.50%" "98.64%"
```


Now we will do the same checking for 2 samples, to see whether there is correlation between each 2 colors distribution.

for each row i and column j, 1) if i==j, this it the check from before of the expected value to n_unit/n_color
 2) if i!=j, this is two samples test of same expected value hypothesis



Now here Is visualization of the actual data per color

```
## 'stat_bin()' using 'bins = 30'. Pick better value with 'binwidth'.
```



Variance Distribution Checking


We know that the distribution of variance is approximately Gamma distribution:

$$f(x) = \frac{1}{(\Gamma(\alpha)\theta^\alpha)} x^{\alpha-1} e^{-x/\theta}$$


We can see that the variance distribution is Gamma like with shape and rate as seen below

```
## [1] "The parameters of the gamma shaped variance is shape 2.574 and rate 0.057"
```

Density Plot with Gamma Distribution



Are All Even in the Sample?



n*m types of snacks

I will create a function that create sample for each number of colors and package size we want, and then calculate some interesting parameters


```
color_op<- 2:8
grams_op<- c(25,45,150,250,330,500,750,1000)
n_unit_op<- grams_op/gram
nn=800
```

We will make the multiple sample. Here is some random rows:


```
##   n_unit n_color even_count even_evens   var_col all_even low_color
## 1  549.5      5  0.500500  0.0965000 115.46125  0.02375  0.0000
## 2   27.5      8  0.496250  0.0590625   3.34125  0.00750  0.5225
## 3  362.6      3  0.510000  0.1750000 120.92750  0.13500  0.0000
## 4   27.5      5  0.505000  0.0980000   5.52000  0.03250  0.3225
## 5  824.2      2  0.510625  0.2543750 431.33312  0.26500  0.0000
##   smallest_col
## 1             79
## 2             0
## 3            93
## 4             0
## 5           366
```


Deep Insight on the Data

here are some insights:





does the Uneven Colors Even




Here is probability of all even, and whether there is pattern.

```
## <Guides[1] ggproto object>
##
## colour : "none"
```







As we can see, only the small package (less than 50 lentils) have high probability of at least one color to appear severely lower.

Therefore, splitting package by color on the big ones should be relatively even.

using regression for correlation check

```
##
## Call:
## lm(formula = mega_snack_2$even_count ~ mega_snack_2$n_color +
##     mega_snack_2$n_unit)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.5332 -0.1744 -0.0091  0.1409  0.4983
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    5.452e-01  2.161e-02  25.235  <2e-16 ***
## mega_snack_2$n_color -6.000e-03  3.679e-03  -1.631   0.103
## mega_snack_2$n_unit  -2.751e-06  2.059e-05  -0.134   0.894
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.2462 on 1117 degrees of freedom
## Multiple R-squared:  0.002392,    Adjusted R-squared:  0.0006057
```

```
## F-statistic: 1.339 on 2 and 1117 DF,  p-value: 0.2625

##
## Call:
## lm(formula = mega_snack_2$all_even ~ mega_snack_2$n_color + mega_snack_2$n_unit +
##     mega_snack_2$color_No2)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.18758 -0.07151 -0.04450  0.02295  0.81752
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      1.761e-01  1.837e-02   9.589  < 2e-16 ***
## mega_snack_2$n_color -3.482e-02  2.910e-03 -11.966  < 2e-16 ***
## mega_snack_2$n_unit  -4.762e-06  1.628e-05  -0.292    0.77
## mega_snack_2$color_No2TRUE  8.125e-02  1.176e-02   6.908 8.21e-12 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.1948 on 1116 degrees of freedom
## Multiple R-squared:  0.1461, Adjusted R-squared:  0.1438
## F-statistic: 63.66 on 3 and 1116 DF,  p-value: < 2.2e-16
```

From the second regression we can suggest that more colors is correlated with 3.3% less probability of all colors even, while even numbers of colors is correlated with 7.7% more probability of all colors eve, regardless of any package size.

Conclusions

Data Structure

The simulation created a random samples of snack packs, which was proven to be statistically random. We created with “sample_MnM” one sample with specific size and numbers of colors, and then “mega_snack” that create costume samples and check the relevand indicators fot this project.

We I out that:

- Small packages often lack at least one color, and sometimes contain only one color.
- As the number of colors increases, the chance that all colors have even counts drops significantly.
- For medium to large packages, the probability of any one color being significantly underrepresented (less than $\frac{2}{3}$ of its expected amount) is near zero.

Main Q: Eating M&M by Two

Although there is no clear pattern to the right M&M package for all the colors to have even count, different approach might find a clear reason for more or less couples of M&M. Here is what I did found:

The general probability of all colors to be even in 6 colored pack is 1.5% for small 50g package 2.1% for big 1000g package, and overall 1.5%, which is more than I expected.

For 5 colored pack like Skittles the average is about 2.9%

For 2 colored pack the average is 24.5%, so for 2 colored marshmallow bag this will be the statistics.

Summary

To sum it up, for each medium pack the probability of all even colors is 1.4%, or 1 in a 73 packs of 250g. So I might need to change my snack preference to marshmallow if I want to keep this method.

This project help me implement the method of simulation on a need— as silly as in might be— and check the business question with wider sight in order to understand the hole story.

I applied:

- Simulation logic
- Exploratory analysis
- Hypothesis testing
- Distribution checks
- Outlier detection
- Visualization using

In addition, I created the infrastructure for similar questions with different parameters to be checked in a reusable, structured way.