

# Expectation and variance

## 1 Expectation

Expectation

Properties of expectation

Example - Expectation of coin related random variables

Example - Maximization of expected profit

## 2 Variance, covariance, correlation coefficient

Variance, covariance, correlation coefficient

Properties of variance and covariance

Jensen's inequality

Mean and variance of the sum of random variables

Mean and variance of the weighted sum of random variables

Example - How to compute the variance

Example - Variance of coin related random variables

## 3 Standardization and reverse standardization

How to measure the typical size of error or deviation from mean

Standardization and reverse standardization

Example - Standardization

## Expectation

## Definition


$$\mathbb{E}(X) = \sum_{x_i} x_i \times P(X = x_i) = \sum_{x_i} x_i \times p_{x_i}$$

## Interpretation 1

$$\underbrace{\mathbb{E}(X)}_{\text{Expected payoff}} = \sum_{x_i} \underbrace{x_i}_{\text{Payoff}} \times \underbrace{p_{x_i}}_{\text{Probability}}$$

## Interpretation 2

$$\underbrace{\mathbb{E}(X)}_{\text{Area under curve}} = \sum_{x_i} \underbrace{x_i}_{\text{Height}} \times \underbrace{P(X = x_i)}_{\text{Width}}$$



## Properties of expectation

## Expectation as a linear operator

$$(1) \quad \mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$$

$$(2) \quad \mathbb{E}(aX) = a\mathbb{E}(X)$$

$$(3) \quad \mathbb{E}(a) = a$$

## Change of variable - Recycle, save the earth

$$(4) \quad \mathbb{E}[g(X)] = \sum_{x_i} g(x_i)p_{x_i}$$

$$(5) \quad \mathbb{E}[g(X, Y)] = \sum_{x_i} \sum_{y_j} g(x_i, y_j)p_{x_i, y_j}$$

## Product of independent random variables

$$(6) \quad \mathbb{E}[XY] = \mathbb{E}[X] \mathbb{E}[Y] \quad \text{if } X \text{ and } Y \text{ are independent}$$

$$(7) \quad \mathbb{E}[g(X)h(Y)] = \mathbb{E}[g(X)] \mathbb{E}[h(Y)] \quad \text{if } X \text{ and } Y \text{ are independent}$$

## No free lunch

$$(8) \quad X \geq 0 \quad \Rightarrow \quad \mathbb{E}(X) \geq 0$$


$$(9) \quad X \geq Y \quad \Rightarrow \quad \mathbb{E}(X) \geq \mathbb{E}(Y)$$

$$(10) \quad |\mathbb{E}(X)| \leq \mathbb{E}(|X|)$$

## Cauchy-Schwartz inequality

$$(11) \quad \mathbb{E}|XY| \leq (\mathbb{E}X^2)^{1/2}(\mathbb{E}Y^2)^{1/2}$$

$$\begin{aligned}
\mathbb{E}(X + Y) &= \sum_{x_i} \sum_{y_j} (x_i + y_j) p_{x_i, y_j} \\
&= \sum_{x_i} \sum_{y_j} x_i p_{x_i, y_j} + \sum_{x_i} \sum_{y_j} y_j p_{x_i, y_j} \\
&= \sum_{x_i} x_i \left( \sum_{y_j} p_{x_i, y_j} \right) + \sum_{y_j} y_j \left( \sum_{x_i} p_{x_i, y_j} \right) \\
&= \sum_{x_i} x_i p_{x_i} + \sum_{y_j} y_j p_{y_j} \\
&= \mathbb{E}(X) + \mathbb{E}(Y)
\end{aligned}$$



$$\begin{aligned}
\mathbb{E}[g(X)] &= \sum_{g_k} g_k P(g(X) = g_k) \\
&= \sum_{g_k} g_k \left( \sum_{x_i \text{ with } g(x_i)=g_k} p_{x_i} \right) \\
&= \sum_{g_k} \left( \sum_{x_i \text{ with } g(x_i)=g_k} g(x_i) p_{x_i} \right) \\
&= \sum_{x_i} g(x_i) p_{x_i}
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}[XY] &= \sum_{x_i} \sum_{y_j} x_i y_j p_{x_i, y_j} = \sum_{x_i} \sum_{y_j} x_i y_j p_{x_i} p_{y_j} \\
&= \left( \sum_{x_i} x_i p_{x_i} \right) \left( \sum_{y_j} y_j p_{y_j} \right) = \mathbb{E}[X] \mathbb{E}[Y]
\end{aligned}$$

$$X \geq 0 \Rightarrow x_i \geq 0 \Rightarrow \mathbb{E}(X) = \sum_{x_i} x_i p_{x_i} \geq 0$$

$$\begin{aligned}
X \geq Y &\Rightarrow X - Y \geq 0 \Rightarrow \mathbb{E}(X - Y) \geq 0 \\
&\Rightarrow \mathbb{E}(X) - \mathbb{E}(Y) \geq 0 \Rightarrow \mathbb{E}(X) \geq \mathbb{E}(Y)
\end{aligned}$$

#### Proof of Cauchy-Schwartz inequality

If  $\mathbb{E}X^2 = 0$ ,  $X = 0$  with probability 1 and hence  $\mathbb{E}|XY| = 0$ . By the same token, if  $\mathbb{E}Y^2 = 0$ , then  $\mathbb{E}|XY| = 0$ . In these two extreme cases (11) holds trivially. So, without loss of generality we assume that  $\mathbb{E}X^2 > 0$  and  $\mathbb{E}Y^2 > 0$ .

With  $t = -\mathbb{E}|XY|/\mathbb{E}Y^2$

$$\begin{aligned}
(|X| + t|Y|)^2 = X^2 + 2t|XY| + t^2Y^2 &\geq 0 \Rightarrow \mathbb{E}X^2 + 2t\mathbb{E}|XY| + t^2\mathbb{E}Y^2 \geq 0 \\
&\Rightarrow (\mathbb{E}X^2)(\mathbb{E}Y^2) \geq (\mathbb{E}|XY|)^2
\end{aligned}$$

Example - Expectation of coin related random variables

Distribution	Expectation	Variance
$B(p)$	$p$	$pq$
$B(n, p)$	$np$	$npq$
$Geo(p)$	$\frac{1}{p}$	$\frac{q}{p^2}$
$NB(r, p)$	$\frac{r}{p}$	$\frac{rq}{p^2}$

$$X \sim B(p) \Rightarrow \mathbb{E}(X) = 1 \times p + 0 \times q = p$$

$$X \sim B(n, p) \Rightarrow X = \sum_{i=1}^n X_i, \quad X_i \text{ iid } B(p)$$

$$\Rightarrow \mathbb{E}(X) = \sum_{i=1}^n \mathbb{E}(X_i) = np$$

$$1 + x + x^2 + \dots = \frac{1}{1-x} \quad \begin{array}{l} \text{Diff wrt } x \\ \Rightarrow \end{array} \quad 1 + 2x + 3x^2 + \dots = \frac{1}{(1-x)^2}$$

$$\quad \begin{array}{l} \text{Let } x = q \\ \Rightarrow \end{array} \quad 1 + 2q + 3q^2 + \dots = \frac{1}{p^2}$$

$$\Rightarrow X \sim Geo(p) \Rightarrow \mathbb{E}(X) = \left( \sum_{k=1}^{\infty} kq^{k-1} \right) p = \frac{1}{p}$$

$$X \sim NB(r, p) \Rightarrow X = \sum_{i=1}^r X_i, \quad X_i \text{ iid } Geo(p)$$

$$\Rightarrow \mathbb{E}(X) = \sum_{i=1}^r \mathbb{E}(X_i) = \frac{r}{p}$$

**Example - Maximization of expected profit**

A newsboy purchases papers at 10 cents and sells them at 15 cents. However, he is not allowed to return unsold papers. If his daily demand  $X$  is  $B(n, p)$  with  $n = 10$ ,  $p = 0.4$ , approximately how many papers should he purchase so as to maximize his expected profit?

With  $t$  purchase, his profit  $Y(t)$  and expected profit  $f(t) = EY(t)$  are

$$Y(t) = \begin{cases} 5t & \text{if } X \geq t \\ 5X - 10(t - X) & \text{if } X < t \end{cases}$$


$$f(t) = 5tP(X \geq t) + \sum_{i=0}^{t-1} (15i - 10t)P(X = i)$$

To find an optimal  $t_0$  that maximize  $f(t)$ , differentiate  $f(t)$  discretely: Find  $t_0$  with

$$\begin{aligned} f(t_0) - f(t_0 - 1) &= 15P(X \geq t_0) - 10 \geq 0 \\ f(t_0 + 1) - f(t_0) &= 15P(X \geq t_0 + 1) - 10 \leq 0 \end{aligned}$$

or

$$P(X \leq t_0 - 1) \leq \frac{1}{3} \quad \text{and} \quad P(X \leq t_0) \geq \frac{1}{3}$$



```

n=10;
p=0.4;

i=0:n;
pmf=binomial(n,i).*(p.^i).*((1-p).^(n-i));
cdf=cumsum(pmf);

% Find the smallest t0 such that P(X<=t0) exceeds 1/3
t=find(cdf>=1/3,1);
t0=t-1 % i starts from 0, but MATLAB index starts from 1

% Plot of CDF of X, the daily demand
bar(i,cdf); grid on; hold on;
plot(-1:0.1:n+1,1/3,'--b')
plot([t0 t0 0],[0 cdf(t0+1) cdf(t0+1)], '--r')
title('CDF of X, the daily demand')
xlabel('Daily demand'); ylabel('CDF');

% Plot of CDF of X
% Critical level 1/3
% F(t0) >= 1/3

```



## Variance, covariance, correlation coefficient

## Variance

$$\text{Var}(X) = \mathbb{E}(X - \mathbb{E}X)^2 = \mathbb{E}X^2 - (\mathbb{E}X)^2$$

## Covariance

$$\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}X)(Y - \mathbb{E}Y)] = \mathbb{E}(XY) - (\mathbb{E}X)(\mathbb{E}Y)$$

## Correlation coefficient

$$-1 \leq \rho = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)}} \leq 1$$

$$\begin{aligned} \text{Var}(X) &= \mathbb{E}(X - \mathbb{E}X)^2 \\ &= \mathbb{E}(X^2 - 2(\mathbb{E}X)X + (\mathbb{E}X)^2) \\ &= \mathbb{E}X^2 - 2(\mathbb{E}X)\mathbb{E}X + (\mathbb{E}X)^2 \\ &= \mathbb{E}X^2 - (\mathbb{E}X)^2 \end{aligned}$$

$$\begin{aligned} \text{Cov}(X, Y) &= \mathbb{E}[(X - \mathbb{E}X)(Y - \mathbb{E}Y)] \\ &= \mathbb{E}[XY - (\mathbb{E}Y)X - (\mathbb{E}X)Y + (\mathbb{E}X)(\mathbb{E}Y)] \\ &= \mathbb{E}(XY) - (\mathbb{E}Y)(\mathbb{E}X) - (\mathbb{E}X)(\mathbb{E}Y) + (\mathbb{E}X)(\mathbb{E}Y) \\ &= \mathbb{E}(XY) - (\mathbb{E}Y)(\mathbb{E}X) \end{aligned}$$

$$\begin{aligned} |\text{Cov}(X, Y)| &\leq \mathbb{E}|(X - \mathbb{E}X)(Y - \mathbb{E}Y)| \\ &\leq (\mathbb{E}(X - \mathbb{E}X)^2)^{1/2}(\mathbb{E}(Y - \mathbb{E}Y)^2)^{1/2} = \sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)} \end{aligned}$$

## Properties of variance and covariance

- (1)  $Var(X) = Cov(X, X)$
- (2)  $Cov(aX + bY, Z) = aCov(X, Z) + bCov(Y, Z)$
- (2)  $Cov(Z, aX + bY) = aCov(Z, X) + bCov(Z, Y)$
- (3)  $Cov(X, Y) = Cov(Y, X)$
- (4)  $Cov(X, a) = Cov(a, X) = 0$
- (5)  $Cov(X, Y) = 0$  if  $X$  and  $Y$  are independent

$$Var(X) = \mathbb{E}(X - \mathbb{E}X)^2 = \mathbb{E}[(X - \mathbb{E}X)(X - \mathbb{E}X)] = Cov(X, X)$$

$$\begin{aligned}
 Cov(aX + bY, Z) &= \mathbb{E}(aX + bY - a\mathbb{E}X - b\mathbb{E}Y)(Z - \mathbb{E}Z) \\
 &= \mathbb{E}[a(X - \mathbb{E}X) + b(Y - \mathbb{E}Y)](Z - \mathbb{E}Z) \\
 &= a\mathbb{E}[(X - \mathbb{E}X)(Z - \mathbb{E}Z)] + b\mathbb{E}[(Y - \mathbb{E}Y)(Z - \mathbb{E}Z)] \\
 &= aCov(X, Z) + bCov(Y, Z)
 \end{aligned}$$

$$Cov(X, Y) = \mathbb{E}[(X - \mathbb{E}X)(Y - \mathbb{E}Y)] = \mathbb{E}[(Y - \mathbb{E}Y)(X - \mathbb{E}X)] = Cov(Y, X)$$

$$Cov(X, a) = \mathbb{E}[(X - \mathbb{E}X)(a - a)] = 0$$

$$\begin{aligned}
 X, Y \text{ independent} &\Rightarrow \mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y) \\
 &\Rightarrow Cov(X, Y) = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y) = 0
 \end{aligned}$$

## Jensen's inequality

## Definition

$\varphi : \mathbf{R} \rightarrow \mathbf{R}$  is **convex** if for any  $x, y$  and  $0 < \lambda < 1$

$$\varphi(\lambda x + (1 - \lambda)y) \leq \lambda\varphi(x) + (1 - \lambda)\varphi(y)$$

$\varphi : \mathbf{R} \rightarrow \mathbf{R}$  is **strictly convex** if for any  $x, y$  and  $0 < \lambda < 1$

$$\varphi(\lambda x + (1 - \lambda)y) < \lambda\varphi(x) + (1 - \lambda)\varphi(y)$$

## Jensen's inequality

If  $\varphi$  convex,  $\mathbb{E}\varphi(X) \geq \varphi(\mathbb{E}X)$

If  $\varphi$  strictly convex,  $\mathbb{E}\varphi(X) = \varphi(\mathbb{E}X) \Leftrightarrow X = \mathbb{E}X$


## Example

$$\varphi(x) = x^2 \quad \Rightarrow \quad \mathbb{E}X^2 \geq (\mathbb{E}X)^2$$

$$\varphi(x) = |x| \quad \Rightarrow \quad \mathbb{E}|X| \geq |\mathbb{E}X|$$

$$\varphi(x) = e^x \quad \Rightarrow \quad \mathbb{E}e^X \geq e^{\mathbb{E}X}$$

$$\varphi(x) = -\log x \quad \Rightarrow \quad \mathbb{E} \log X \leq \log \mathbb{E}X \quad \text{for } X > 0$$



$$\varphi(X) \geq \alpha(X - \mu) + \varphi(\mu) \quad \xRightarrow{\text{Take expectation}} \quad \mathbb{E}\varphi(X) \geq \alpha\mathbb{E}(X - \mu) + \varphi(\mu) = \varphi(\mu) = \varphi(\mathbb{E}X)$$

## Mean and variance of the sum of random variables

In general

$$\begin{aligned}\mathbb{E}\left(\sum_{i=1}^n X_i\right) &= \sum_{i=1}^n \mathbb{E}(X_i) \\ \text{Var}\left(\sum_{i=1}^n X_i\right) &= \sum_{i=1}^n \text{Var}(X_i) + \sum_{i \neq j} \text{Cov}(X_i, X_j) \\ &= \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{1 \leq i < j \leq n} \text{Cov}(X_i, X_j)\end{aligned}$$

If  $X_i$  are independent

$$\begin{aligned}\mathbb{E}\left(\sum_{i=1}^n X_i\right) &= \sum_{i=1}^n \mathbb{E}(X_i) \\ \text{Var}\left(\sum_{i=1}^n X_i\right) &= \sum_{i=1}^n \text{Var}(X_i)\end{aligned}$$

If  $X_i$  are iid

$$\begin{aligned}\mathbb{E}\left(\sum_{i=1}^n X_i\right) &= \sum_{i=1}^n \mathbb{E}(X_i) = n\mathbb{E}(X_1) \\ \text{Var}\left(\sum_{i=1}^n X_i\right) &= \sum_{i=1}^n \text{Var}(X_i) = n\text{Var}(X_1)\end{aligned}$$

## Mean and variance of the weighted sum of random variables

In general

$$\begin{aligned}
\mathbb{E}\left(\sum_{i=1}^n a_i X_i\right) &= \sum_{i=1}^n a_i \mathbb{E}(X_i) \\
\text{Var}\left(\sum_{i=1}^n a_i X_i\right) &= \sum_{i=1}^n a_i^2 \text{Var}(X_i) + \sum_{i \neq j} a_i a_j \text{Cov}(X_i, X_j) \\
&= \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{1 \leq i < j \leq n} a_i a_j \text{Cov}(X_i, X_j)
\end{aligned}$$

If  $X_i$  are independent

$$\begin{aligned}
\mathbb{E}\left(\sum_{i=1}^n a_i X_i\right) &= \sum_{i=1}^n a_i \mathbb{E}(X_i) \\
\text{Var}\left(\sum_{i=1}^n a_i X_i\right) &= \sum_{i=1}^n a_i^2 \text{Var}(X_i)
\end{aligned}$$

If  $X_i$  are iid

$$\begin{aligned}
\mathbb{E}\left(\sum_{i=1}^n a_i X_i\right) &= \sum_{i=1}^n a_i \mathbb{E}(X_i) = \left(\sum_{i=1}^n a_i\right) \mathbb{E}(X_1) \\
\text{Var}\left(\sum_{i=1}^n a_i X_i\right) &= \sum_{i=1}^n a_i^2 \text{Var}(X_i) = \left(\sum_{i=1}^n a_i^2\right) \text{Var}(X_1)
\end{aligned}$$

Example - How to compute the variance

Let  $Var(X) = 2$ ,  $Var(Y) = 2$ ,  $Var(Z) = 3$ ,  $Cov(X, Y) = 0.25$ . If  $Z$  is independent to both  $X$  and  $Y$ , compute the variance of  $V$ , where  $V$  is given by

$$V = X + 2Y - 3Z - 2$$

$$\begin{aligned}
 Var(V) &\stackrel{(1)}{=} Cov(V, V) = Cov(X + 2Y - 3Z - 2, X + 2Y - 3Z - 2) \\
 &\stackrel{(2)}{=} \begin{array}{cccc} Cov(X, X) & +2Cov(X, Y) & -3Cov(X, Z) & -2Cov(X, 1) \\ 2Cov(Y, X) & +4Cov(Y, Y) & -6Cov(Y, Z) & -4Cov(Y, 1) \\ -3Cov(Z, X) & -6Cov(Z, Y) & +9Cov(Z, Z) & +6Cov(Z, 1) \\ -2Cov(1, X) & -4Cov(1, Y) & +6Cov(1, Z) & +4Cov(1, 1) \end{array} \\
 &\stackrel{(3)}{=} \begin{array}{cccc} Cov(X, X) & +2Cov(X, Y) & -3Cov(X, Z) & -2Cov(X, 1) \\ 2Cov(X, Y) & +4Cov(Y, Y) & -6Cov(Y, Z) & -4Cov(Y, 1) \\ -3Cov(X, Z) & -6Cov(Y, Z) & +9Cov(Z, Z) & +6Cov(Z, 1) \\ -2Cov(X, 1) & -4Cov(Y, 1) & +6Cov(Z, 1) & +4Cov(1, 1) \end{array} \\
 &= \begin{array}{cccc} Cov(X, X) & +4Cov(X, Y) & -6Cov(X, Z) & -4Cov(X, 1) \\ & +4Cov(Y, Y) & -12Cov(Y, Z) & -8Cov(Y, 1) \\ & & +9Cov(Z, Z) & +12Cov(Z, 1) \\ & & & +4Cov(1, 1) \end{array} \\
 &\stackrel{(4)}{=} \begin{array}{ccc} Cov(X, X) & +4Cov(X, Y) & -6Cov(X, Z) \\ & +4Cov(Y, Y) & -12Cov(Y, Z) \\ & & +9Cov(Z, Z) \end{array} \\
 &\stackrel{(5)}{=} \begin{array}{ccc} Cov(X, X) & +4Cov(X, Y) & \\ & +4Cov(Y, Y) & \\ & & +9Cov(Z, Z) \end{array} \\
 &\stackrel{(1)}{=} \begin{array}{ccc} Var(X) & +4Cov(X, Y) & \\ & +4Var(Y) & \\ & & +9Var(Z) \end{array} \\
 &= \begin{array}{ccc} 2 & +4 \cdot 0.25 & \\ & +4 \cdot 2 & \\ & & +9 \cdot 3 \end{array} \\
 &= 38
 \end{aligned}$$

Example - Variance of coin related random variables

Distribution	Expectation	Variance
$B(p)$	$p$	$pq$
$B(n, p)$	$np$	$npq$
$Geo(p)$	$\frac{1}{p}$	$\frac{q}{p^2}$
$NB(r, p)$	$\frac{r}{p}$	$\frac{rq}{p^2}$

$$\begin{aligned}
 X \sim B(p) &\Rightarrow \mathbb{E}(X^2) = \mathbb{E}(X) = p \\
 &\Rightarrow \text{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}X)^2 = p - p^2 = p(1 - p) = pq
 \end{aligned}$$

$$\begin{aligned}
 X \sim B(n, p) &\Rightarrow X = \sum_{i=1}^n X_i, \quad X_i \text{ iid } B(p) \\
 &\Rightarrow \text{Var}(X) = \sum_{i=1}^n \text{Var}(X_i) = npq
 \end{aligned}$$

$$\begin{aligned}
 1 + x + x^2 + \dots &= \frac{1}{1-x} & \text{Diff} \Rightarrow_x & 1 + 2x + 3x^2 + \dots = \frac{1}{(1-x)^2} \\
 & & \text{Diff} \Rightarrow_x & 2 \cdot 1 + 3 \cdot 2x + 4 \cdot 3x^2 \dots = \frac{2}{(1-x)^3} \\
 \text{Let } x &= q & \Rightarrow & 2 \cdot 1 + 3 \cdot 2q + 4 \cdot 3q^2 \dots = \frac{2}{p^3}
 \end{aligned}$$

$$\begin{aligned}
 X \sim Geo(p) &\Rightarrow \mathbb{E}[X(X-1)] = \left( \sum_{k=1}^{\infty} k(k-1)q^{k-2} \right) qp = \frac{2q}{p^2} \\
 &\Rightarrow \mathbb{E}X^2 = \mathbb{E}[X(X-1) + X] = \mathbb{E}[X(X-1)] + \mathbb{E}X = \frac{2q}{p^2} + \frac{1}{p} \\
 &\Rightarrow \text{Var}(X) = \mathbb{E}X^2 - (\mathbb{E}X)^2 = \frac{2q}{p^2} + \frac{1}{p} - \frac{1}{p^2} = \frac{q}{p^2}
 \end{aligned}$$

$$\begin{aligned}
 X \sim NB(r, p) &\Rightarrow X = \sum_{i=1}^r X_i, \quad X_i \text{ iid } Geo(p) \\
 &\Rightarrow \text{Var}(X) = \sum_{i=1}^r \text{Var}(X_i) = \frac{rq}{p^2}
 \end{aligned}$$



How to measure the typical size of error or deviation from mean

First try

$$\underbrace{\mathbb{E}}_{\text{Average}} \underbrace{(X - \mathbb{E}X)}_{\text{Error}}$$

However, this try is fertile:  $\mathbb{E}(X - \mathbb{E}X) = \mathbb{E}X - \mathbb{E}X = 0$ .


Second try

$$\underbrace{\mathbb{E}}_{\text{Average}} \underbrace{|X - \mathbb{E}X|}_{\text{Error}}$$

Due to computational difficulties, this measure of typical error size is not popular.

Standard way to measure typical error size

$$\underbrace{SD(X)}_{\text{Standard deviation}} = \sqrt{\underbrace{\mathbb{E}}_{\text{Average}} \underbrace{(X - \mathbb{E}X)^2}_{\text{Error}}} = \sqrt{\text{Var}(X)}$$



## Standardization and reverse standardization

## Mean and variance lemma for standardization and reverse standardization

$$\begin{aligned}\mathbb{E}(aX + b) &= a\mathbb{E}(X) + b \\ \text{Var}(aX + b) &= \text{Var}(aX) = a^2\text{Var}(X)\end{aligned}$$

## Standardization

If  $X$  has mean  $\mu$  and standard deviation  $\sigma$ , then

$$\frac{X - \mu}{\sigma} \text{ has mean 0 and standard deviation 1}$$


## Reverse standardization

If  $X$  has mean 0 and standard deviation 1, then

$$\mu + \sigma * X \text{ has mean } \mu \text{ and standard deviation } \sigma$$

$$\begin{aligned}\text{Var}(aX + b) &= \text{Cov}(aX + b, aX + b) \\ &= a^2\text{Var}(X) + 2ab\text{Cov}(X, 1) + b^2\text{Var}(1) \\ &= a^2\text{Var}(X)\end{aligned}$$

$$\text{Var}(aX) = \text{Cov}(aX, aX) = a^2\text{Var}(X)$$



### Example - Standardization

We flip a fair coin many times.

$X_i$                        $i^{th}$  flip record, where  $H$  and  $T$  are recorded as 1 and 0

$Y_i := 2X_i - 1$        $i^{th}$  flip record, where  $H$  and  $T$  are recorded as 1 and  $-1$

Calculate the mean and variance of the following related random variables, i.e., fill up blanks of the below table.

Random variable	Mean	Variance
$Y_i$		
$\sum_{i=1}^n Y_i$		
$\frac{\sum_{i=1}^n Y_i}{\sqrt{n}}$		

$$X_i \text{ iid } B(0.5) \quad \Rightarrow \quad \mathbb{E}X_i = 0.5, \text{ } Var(X_i) = 0.5 * (1 - 0.5) = 0.25$$

$$\Rightarrow \quad Y_i \text{ iid with } \mathbb{E}Y_i = 0, \text{ } Var(Y_i) = 1$$

$$\Rightarrow \quad \mathbb{E} \left( \sum_{i=1}^n Y_i \right) = 0, \text{ } Var \left( \sum_{i=1}^n Y_i \right) = n$$

$$\text{Standardization} \Rightarrow \quad \frac{\sum_{i=1}^n Y_i}{\sqrt{n}} \text{ has mean 0, variance 1}$$

Random variable	Mean	Variance
$Y_i$	0	1
$\sum_{i=1}^n Y_i$	0	$n$
$\frac{\sum_{i=1}^n Y_i}{\sqrt{n}}$	0	1