

The microbiome affects hormonal therapy in breast cancer

**저자
(Authors)** Jeongshin An, Jong Bin Kim, Eun Yeol Yang, Jinho Yang, Won-Hee Lee, Jong-kyu Kim, HyunGoo Kim, Se Hyun Paek, Jun Woo Lee, Joohyun Woo, Hyungju Kwon, Woosung Lim, Nam Sun Paik, Yoon-Keun Kim, Byung-In Moon

**출처
(Source)** [대한임상종양학회 학술대회지](#) , 2019.2, 201-203 (3 pages)

**발행처
(Publisher)** [대한종양외과학회](#)
Korea society of Surgical Oncology

URL <http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE09326315>

APA Style Jeongshin An, Jong Bin Kim, Eun Yeol Yang, Jinho Yang, Won-Hee Lee, Jong-kyu Kim, HyunGoo Kim, Se Hyun Paek, Jun Woo Lee, Joohyun Woo, Hyungju Kwon, Woosung Lim, Nam Sun Paik, Yoon-Keun Kim, Byung-In Moon (2019). The microbiome affects hormonal therapy in breast cancer. [대한임상종양학회 학술대회지](#), 201-203.

**이용정보
(Accessed)** 이화여자대학교
203.255.***.68
2020/05/18 04:02 (KST)

저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되는 저작물은 DBpia와 구독계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

Copyright information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.

OP1-4

The microbiome affects hormonal therapy in breast cancer

Jeongshin An¹, Jong Bin Kim², Eun Yeol Yang¹, Jinho Yang³, Won-Hee Lee³,
Jong-kyu Kim¹, HyunGoo Kim¹, Se Hyun Paek¹, Jun Woo Lee¹, Joohyun Woo¹,
Hyungju Kwon¹, Woosung Lim¹, Nam Sun Paik¹, Yoon-Keun Kim³, Byung-In Moon¹

¹Department of Surgery, Ewha Womans University, Korea,

²Research Center for Cellular Homeostasis, Ewha Womans University, Korea,

³Department of Research, MD Healthcare, Korea

Background/Aims

Microbiomes are associated with breast cancer, and breast cancer is closely related to estrogen. Among symbiotic bacteria in the human body, microbiomes which produce estrogen-metabolizing enzymes called estrobolome. This study identified the microbiome in breast cancer and normal group and found microbiome which was abundant in the breast cancer group but poor in the normal group. We hypothesized that these bacteria would potentially affect the hormone treatment of breast cancer and tamoxifen treatment. We report the bacteria and its mechanism by which affect tamoxifen efficacy.

Methods

The 347 urine samples in female (127 breast cancer patients and 220 normal individuals) were collected. The mean age in urine samples 51.8 in cancer groups and 59.2 years in control groups. These urine samples were analyzed by NGS using a universal bacterial primer of 16S rDNA. Metagenomic analysis of urine samples was performed at phyla to genus-level. Target bacteria which is abundant in breast cancer of luminal type was selected. Tamoxifen or K.pneumoniae-derived EVs plus tamoxifen was used in the MCF7 cells, and we compared cell survival, cell cycle, and signaling molecules in each group. Human estrogen receptor-positive cells (MCF7) were cultured and bacterial extracellular vesicles (EVs) were isolated from Klebsiella pneumoniae, the target bacteria. For cell-cycle analysis, the cells were treated with distilled water as a control, EVs of K.pneumoniae at 10, 100, 1000 ng/ml for 72 h. Quantitative Real-Time PCR and western blotting for signaling molecule analysis were performed after treatment of EVs in breast cancer cells.

Results

Microbiome between breast cancer and normal group was showed the significantly different. Among these results, Klebsiella was more abundant in luminal A type, estrogen-receptor positive breast cancer and less in the normal group. The efficacy of tamoxifen was verified by comparing the results between cells treated with tamoxifen alone and those treated with tamoxifen and bacterial EVs. When K.

pneumoniae-derived EVs were used alone, the growth of MCF7 cells was not suppressed significantly. However, K. pneumoniae EVs co-treated with tamoxifen inhibited the cell growth of MCF7 by two-folds compared with tamoxifen treatment alone. When K. pneumoniae EVs and tamoxifen were used together, cyclin E2 transcription was down-regulated, and protein levels of P21

Conclusions

In conclusion, these results show that K.pneumoniae may affect the patients who have estrogen-receptor positive breast cancer according to metagenomic analysis. K. pneumoniae-derived EVs enhanced the anti-estrogen effect of tamoxifen through down-regulation of survival signaling molecules in estrogen positive breast cancer cells.



