



Sirt1-Mediated Anti-Aging Effects of Houttuynia cordata Extract in a High Glucose-Induced Endothelial Cell-Aging Model

저자 (Authors)	Gi Dae Kim
출처 (Source)	Preventive Nutrition and Food Science 25(1) , 2020.3, 108-112 (5 pages)
발행처 (Publisher)	한국식품영양과학회 The Korean Society of Food Science and Nutrition
URL	http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE09323092
APA Style	Gi Dae Kim (2020). Sirt1-Mediated Anti-Aging Effects of Houttuynia cordata Extract in a High Glucose-Induced Endothelial Cell-Aging Model. Preventive Nutrition and Food Science, 25(1), 108-112.
이용정보 (Accessed)	이화여자대학교 203.255.***.68 2020/05/18 04:04 (KST)

저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되는 저작물은 DBpia와 구독계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

Copyright Information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.

Sirt1-Mediated Anti-Aging Effects of *Houttuynia cordata* Extract in a High Glucose-Induced Endothelial Cell-Aging Model

Gi Dae Kim

Department of Food and Nutrition, Kyungnam University, Gyeongsang 51767, Korea

ABSTRACT: *Houttuynia cordata* (HC) is a herb widely used in traditional Asian medicine as an ingredient in complex prescriptions. HC is known for its anti-leukemic, anti-oxidative, and anti-inflammatory properties. However, its anti-vascular endothelial aging efficacy and the underlying mechanisms are not fully understood. In this study, we investigated the anti-aging effects of HC in a high glucose (HG)-induced endothelial cell (EC)-aging model. Treatment with HC (40 µg/mL) increased migration of ECs, and increased phosphorylation of extracellular signal-regulated kinases and p38 in a dose-dependent manner. Following HG treatment (30 mM), HC significantly decreased the number of senescence-associated β -galactosidase positive cells, which are the biomarkers for aging, in a dose-dependent manner. Based on levels of phosphorylation, HC (40 µg/mL) was shown to increase expression of sirtuin1 (Sirt1) and endothelial nitric oxide synthase (eNOS) by 74.4% and 328.2%, respectively. Furthermore, treatment of HG-induced senescent ECs with HC (40 µg/mL) significantly increased nitric oxide production ($P<0.05$). These results demonstrate that HC both increases EC migration and regulates the Sirt1/eNOS pathway, suggesting HC has potential for protecting ECs against HG-induced aging.

Keywords: anti-aging, endothelial cell, *Houttuynia cordata*, Sirt1/eNOS

INTRODUCTION

Senescence of vascular endothelial cells (ECs) can contribute to the development of atherosclerosis in humans (Minamino and Komuro, 2007; Wu et al., 2019). Many studies have shown that high glucose (HG) can induce EC aging (Matsui-Hirai et al., 2011; Hayashi et al., 2014; Lin et al., 2019). Moreover, EC dysfunctions can lead to nitric oxide (NO) deficiency (Thoonen et al., 2015). NO is synthesized from L-arginine by NO synthase, of which there are three important isoforms: the endothelial nitric oxide synthase (eNOS), neuronal NOS, and inducible NOS (iNOS). For example, eNOS has been shown to play a vital role in the development of diabetic cardiovascular complications by exacerbating oxidative stress (Kayama et al., 2015). In addition, sirtuin1 (Sirt1), a nicotinamide adenine dinucleotide (NAD⁺)-dependent histone deacetylase, is involved in regulation of metabolism, cell survival, differentiation, and aging. Sirt1 is a key regulator of vascular EC aging. Previous studies have shown that Sirt1 activation can protect against vascular endothelial dysfunctions, whilst controlling eNOS generation (Hu et al., 2017).

Houttuynia cordata (HC) Thunberg, a member of the Saururaceae family, is a herb used for traditional healing in Southeast Asia. Recently, HC has been shown to be a rich source of naturally occurring polysaccharides and flavonoids (Lu et al., 2006a). Hence, HC is used for immune stimulation and chemotherapy in alternative medicine. Furthermore, HC exhibits various pharmacological properties, such as anti-leukemic (Chang et al., 2001), anti-oxidative (Hsu et al., 2016), and anti-inflammatory (Lu et al., 2006b) properties. Several studies have investigated HC, however only a few have evaluated the role of HC in preventing EC aging. To the best of our knowledge, this is the first study to show suppressive effects of HC on aging in a HG-induced aging model. This was achieved by using human umbilical vein ECs and evaluating the underlying mechanisms.

MATERIALS AND METHODS

Reagents

HC was obtained from Dr. Park at Kyungnam University, where it was extracted, separated, and subjected to qual-

Received 3 December 2019; Accepted 12 February 2020; Published online 31 March 2020

Correspondence to Gi Dae Kim, Tel: +82-55-249-2176, E-mail: gidaekim@kyungnam.ac.kr

Author information: Gi Dae Kim (Professor)

Copyright © 2020 by The Korean Society of Food Science and Nutrition. All rights Reserved.

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ity control, as described previously (Shon et al., 2014). 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and gelatin were purchased from Sigma-Aldrich Co. (St. Louis, MO, USA). The senescence-associated β -galactosidase (SA- β -gal) kit was purchased from Abcam (Cambridge, UK) and the NO assay kit was purchased from Thermo Fisher Scientific (Vienna, Austria). p-p38, p-Sirt1, and p-eNOS antibodies were purchased from Cell Signaling Technology (Danvers, MA, USA), and the β -actin and p-extracellular signal-regulated kinases (p-ERK) antibodies were purchased from Santa Cruz Biotechnology (Dallas, TX, USA). Horseradish peroxidase (HRP)-conjugated anti-mouse and anti-rabbit antibodies were purchased from GeneTex Inc. (Irvine, CA, USA).

EC culture

Human umbilical vein endothelial cells were obtained from the American Type Culture Collection (Manassas, VA, USA). Cells were cultured at 37°C with 5% CO₂ in endothelial growth medium-2 (EGM-2; Lonza, Walkersville, MD, USA) supplemented with 10% fetal bovine serum (FBS). ECs were cultured for 48 h at 37°C with 5% CO₂ in EGM-2 (control) or HG 30 mM medium, with or without addition of different concentrations of HC (10~40 μ g/mL).

Cell viability assay

Cells were cultured at 37°C for 72 h in EGM-2 medium supplemented with 2% FBS and various concentrations of HC, and were treated with MTT solution for 4 h. Resulting formazan deposits were dissolved with dimethyl sulfoxide, where the absorbance was measured at 570 nm using a VersaMax ELISA microplate reader (Molecular Devices, Sunnyvale, CA, USA).

Scratch-wound migration assay

Cells were wounded, and then culture media was replaced with fresh media containing various concentrations of HC. Cells were maintained for 24 to 48 h. When at full migration, cells were imaged using a microscope (Olympus Optical Co., Ltd., Tokyo, Japan). The migrated cells were counted using an optical microscope at 200 \times magnification, and quantified manually.

SA- β -gal staining

To determine the number of senescent cells, SA- β -gal assays were performed using the SA- β -gal kit (Abcam), according to the manufacturer's instructions. Cells were fixed for 5 min in β -gal fixative, washed with PBS, and then stained using β -gal fixative solution at 37°C. This process was performed until β -gal staining was visible in either the experimental or control plate. SA- β -gal positive cells were observed via microscopy, with >500 cells counted using three independent fields.

Western blot analysis

Cells were harvested and lysed in protein-extraction solution (Intron Biotechnology, Inc., Gyeonggi, Korea) containing protease and phosphatase inhibitors (10 min at 4°C). Total protein concentrations in the supernatants were measured using Bradford assays. After heating at 95°C for 5 min, protein samples (30 μ g) were separated by 8~12% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Proteins were then transferred onto polyvinylidene fluoride membranes (Millipore, Bedford, MA, USA) at 100 V for 60 min. Membranes were incubated in 5% bovine serum albumin (BSA) in Tris-buffered saline (TBS) solution supplemented with 0.05% TBS with Tween-20 (TBST) (30 min at room temperature), then incubated in 5% BSA in TBST supplemented with primary antibodies (1:200 to 1:1,000) (overnight at 4°C). Next, membranes were inoculated with either HRP-conjugated goat anti-rabbit or anti-mouse antibodies (1:5,000) for 1 h, and protein bands were detected using an enhanced chemiluminescence detection kit (Intron Biotechnology, Inc.) and a LAS-1000 Imager (Fuji Film Corp., Tokyo, Japan). Semi-quantitative analysis of the densitometry results was performed using Image J (National Institutes of Health, Bethesda, MD, USA).

Measurement of NO production

NO concentrations were determined by detecting the concentration of nitrite using NO assay kits, according to manufacturer's instructions. Optical densities were determined at 560 nm using a microplate reader, and NO concentrations were calculated using a calibration curve.

Statistical analysis

Results are expressed as mean \pm standard deviation (SD). Statistical significance was determined by one-way analysis of variance (ANOVA). Post hoc comparisons between groups were performed using Bonferroni multiple comparisons tests. All calculations were performed using SPSS in Windows (v.23.0; IBM Corp., Armonk, NY, USA), and *P*-values <0.05 were considered statistically significant.

RESULTS AND DISCUSSION

HC cytotoxicity

To determine the cytotoxicity of HC on ECs, cells were incubated with various concentrations of HC (0 to 100 μ g/mL) for 72 h. HC showed significant cytotoxicity (*P*<0.05) at concentrations >50 μ g/mL at all-time points of treatment (Fig. 1). Therefore, HC concentrations of \leq 50 μ g/mL were considered non-toxic to ECs and were used in subsequent experiments.

HC increases cell migration.

The mitogen-activated protein kinase (MAPK) pathway regulates key processes involved in vascular EC develop-




Fig. 1. Effects of *Houttuynia cordata* (HC) on cell viability. Cells were treated with HC (0 to 100 $\mu\text{g/mL}$) for 72 h. Cell viability was calculated as the percentage of viable cells cultured in the absence of HC. Results show mean \pm SD. * $P<0.05$.

ment, such as cell proliferation, invasion, metastasis and survival, and angiogenesis (Pišlar et al., 2015). The MAPK involves ERK, c-Jun N-terminal kinase, and p38 (Kim and Choi, 2010). ERK and p38 are important signaling proteins involved in wound healing, whereas ERK plays an important role in controlling cell migration, proliferation, differentiation, and survival (Roskoski, 2012). In a recent study, the p38 MAPK signaling pathway was suggested to be involved in wound healing (Loughlin and Artlett, 2011; Kim et al., 2019; Wang et al., 2019). In the present study, we examined protein expression of ERK and p38 to identify the signaling pathways involved in HC-regulated EC migration. HC treatment (40 $\mu\text{g/mL}$) increased EC migration 1.5-fold relative to the control ($P<0.05$) (Fig. 2A). Western blot analysis showed that HC treatment (20 to 40 $\mu\text{g/mL}$) significantly increased phosphorylation of ERK in ECs ($P<0.05$). Furthermore, HC increased phosphorylation of p38 in a concentration-dependent manner; in particular, treatment with 40 $\mu\text{g/mL}$ HC, increased activation of p38 >2.5-fold higher than the control ($P<0.01$) (Fig. 2B). These results show that




Fig. 2. Effects of *Houttuynia cordata* (HC) on cell migration. (A) Representative images show migration of control and HC-treated cells. Cell migration was expressed as the percentage of migration observed for control cells. (B) Representative blots of levels of p-extracellular signal-regulated kinases (p-ERK) and p-p38 phosphorylation. Quantitative analysis of p-ERK/ β -actin and p-p38/ β -actin. Results show mean \pm SD. * $P<0.05$ and ** $P<0.01$.

HC regulates phosphorylation of ERK and p38 in a concentration-dependent manner. From these results of ERK and p38 MAPK activation, we concluded that HC induces EC migration and enhances its function in wound closure.

HC suppresses aging in HG-induced ECs.

To investigate the inhibitory effects HC on EC aging, cells were treated with glucose (30 mM) and various concentrations of HC (10 to 40 $\mu\text{g/mL}$). A trend of delayed aging was noted following treatment with increasing con-

centrations of HC. In particular, treatment with 40 $\mu\text{g/mL}$ HC significantly decreased the abundance of the aging biomarker (SA- β -gal positive cells) by 10.2% compared with cells in the control group (31.5%) ($P < 0.01$) (Fig. 3A). To determine if HC regulates expression of Sirt1, a protein that regulates aging, phosphorylation of Sirt1 and eNOS was investigated following treatment with HC. We showed HC (20 and 40 $\mu\text{g/mL}$) significantly increased the expression of Sirt1 by 58.5% and 74.4%, respectively, compared with cells in the control group ($P < 0.05$). Following treatment with 40 $\mu\text{g/mL}$ HC, expression of eNOS




Fig. 3. Effects of *Houttuynia cordata* (HC) on the aging of cells cultured in high glucose (HG) medium (30 mM). (A) Representative images senescence-associated β -galactosidase (SA- β -gal) positive cells (blue) following culture in media containing HG, or HG and HC (10 to 40 $\mu\text{g/mL}$) for 48 h. The number of SA- β -gal positive cells are expressed as a percentage of those in the control group. (B) Representative blots showing levels of p-Sirt1 and p-eNOS phosphorylation. Phosphorylation was quantified relative to β -actin. (C) NO production in cells in each treatment group. Results show mean \pm SD of five independent experiments. * $P < 0.05$ and ** $P < 0.01$.

was also significantly increased by over 3-fold HC compared with the control group ($P < 0.01$) (Fig. 3B). Moreover, treatment with 40 $\mu\text{g/mL}$ HC significantly increased NO production ($P < 0.05$) in HG-induced aged ECs (Fig. 3C).

Following treatment with high concentrations of glucose, we observed a large number of ECs expressing the aging biomarker SA- β -gal, a major regulator of HG-induced aging (Sun et al., 2015). In previous studies, eNOS activation has been reported to control NO-mediated delays to cell aging in human ECs (Hayashi et al., 2008; Lee et al., 2017). Therefore, stimulating NO production may protect cells against oxidative stress and decrease the time to recovery. In this study, treatment of HG-treated ECs with 40 $\mu\text{g/mL}$ HC significantly increased NO levels, which suggests that HC enhances to EC-mediated wound healing, and could therefore be used to treat aging in HG-induced ECs by mediating the Sirt1/eNOS pathway.

ACKNOWLEDGEMENTS

This study was funded by the Kyungnam University Foundation Grant, No 20170139.

AUTHOR DISCLOSURE STATEMENT

The author declares no conflict of interest.

REFERENCES

- Chang JS, Chiang LC, Chen CC, Liu LT, Wang KC, Lin CC. Anti-leukemic activity of *Bidens pilosa* L. var. minor (Blume) Sherff and *Houttuynia cordata* Thunb.. *Am J Chin Med*. 2001. 29:303-312.
- Hayashi T, Kotani H, Yamaguchi T, Taguchi K, Iida M, Ina K, et al. Endothelial cellular senescence is inhibited by liver X receptor activation with an additional mechanism for its atheroprotection in diabetes. *Proc Natl Acad Sci USA*. 2014. 111:1168-1173.
- Hayashi T, Yano K, Matsui-Hirai H, Yokoo H, Hattori Y, Iguchi A. Nitric oxide and endothelial cellular senescence. *Pharmacol Ther*. 2008. 120:333-339.
- Hsu CC, Yang HT, Ho JJ, Yin MC, Hsu JY. *Houttuynia cordata* aqueous extract attenuated glycation and oxidative stress in heart and kidney of diabetic mice. *Eur J Nutr*. 2016. 55:845-854.
- Hu T, Chen Y, Jiang Q, Lin J, Li H, Wang P, et al. Overexpressed eNOS upregulates SIRT1 expression and protects mouse pancreatic β cells from apoptosis. *Exp Ther Med*. 2017. 14:1727-1731.
- Kayama Y, Raaz U, Jagger A, Adam M, Schellinger IN, Sakamoto M, et al. Diabetic cardiovascular disease induced by oxidative stress. *Int J Mol Sci*. 2015. 16:25234-25263.
- Kim EK, Choi EJ. Pathological roles of MAPK signaling pathways in human diseases. *Biochim Biophys Acta*. 2010. 1802:396-405.
- Kim J, Kim B, Kim SM, Yang CE, Song SY, Lee WJ, et al. Hypoxia-induced epithelial-to-mesenchymal transition mediates fibroblast abnormalities via ERK activation in cutaneous wound healing. *Int J Mol Sci*. 2019. 20:2546. <https://doi.org/10.3390/ijms20102546>
- Lee HY, Zeeshan HMA, Kim HR, Chae HJ. Nox4 regulates the eNOS uncoupling process in aging endothelial cells. *Free Radic Biol Med*. 2017. 113:26-35.
- Lin X, Li S, Wang YJ, Wang Y, Zhong JY, He JY, et al. Exosomal Notch3 from high glucose-stimulated endothelial cells regulates vascular smooth muscle cells calcification/aging. *Life Sci*. 2019. 232:116582. <https://doi.org/10.1016/j.lfs.2019.116582>
- Loughlin DT, Artlett CM. Modification of collagen by 3-deoxyglucosone alters wound healing through differential regulation of p38 MAP kinase. *PLoS One*. 2011. 6:e18676. <https://doi.org/10.1371/journal.pone.0018676>
- Lu HM, Liang YZ, Chen S. Identification and quality assessment of *Houttuynia cordata* injection using GC-MS fingerprint: a standardization approach. *J Ethnopharmacol*. 2006a. 105:436-440.
- Lu HM, Liang YZ, Yi LZ, Wu XJ. Anti-inflammatory effect of *Houttuynia cordata* injection. *J Ethnopharmacol*. 2006b. 104:245-249.
- Matsui-Hirai H, Hayashi T, Yamamoto S, Ina K, Maeda M, Kotani H, et al. Dose-dependent modulatory effects of insulin on glucose-induced endothelial senescence *in vitro* and *in vivo*: a relationship between telomeres and nitric oxide. *J Pharmacol Exp Ther*. 2011. 337:591-599.
- Minamino T, Komuro I. Vascular cell senescence: contribution to atherosclerosis. *Circ Res*. 2007. 100:15-26.
- Pišlar A, Perišić Nanut M, Kos J. Lysosomal cysteine peptidases — molecules signaling tumor cell death and survival. *Semin Cancer Biol*. 2015. 35:168-179.
- Roskoski R Jr. ERK1/2 MAP kinases: structure, function, and regulation. *Pharmacol Res*. 2012. 66:105-143.
- Shon MS, Lee Y, Song JH, Park T, Lee JK, Kim M, et al. Anti-aging potential of extracts prepared from fruits and medicinal herbs cultivated in the Gyeongnam area of Korea. *Prev Nutr Food Sci*. 2014. 19:178-186.
- Sun Y, Hu X, Hu G, Xu C, Jiang H. Curcumin attenuates hydrogen peroxide-induced premature senescence via the activation of SIRT1 in human umbilical vein endothelial cells. *Biol Pharm Bull*. 2015. 38:1134-1141.
- Thoonen R, Cauwels A, Decaluwe K, Geschka S, Tainsh RE, Delanghe J, et al. Cardiovascular and pharmacological implications of haem-deficient NO-unresponsive soluble guanylate cyclase knock-in mice. *Nat Commun*. 2015. 6:8482.
- Wang T, Li X, Fan L, Chen B, Liu J, Tao Y, et al. Negative pressure wound therapy promoted wound healing by suppressing inflammation via down-regulating MAPK-JNK signaling pathway in diabetic foot patients. *Diabetes Res Clin Pract*. 2019. 150:81-89.
- Wu X, Zheng W, Jin P, Hu J, Zhou Q. Role of IGFBP1 in the senescence of vascular endothelial cells and severity of aging-related coronary atherosclerosis. *Int J Mol Med*. 2019. 44:1921-1931.