



## Thirteen Years' Experience of Pediatric Abdominal Mass and Analysis of Associated Factors Affecting Repeated Operation

---

저자 (Authors)	Hee-Beom Yang, Hyun-Young Kim, Sung Eun Jung
출처 (Source)	<a href="#">대한임상종양학회 학술대회지</a> , 2019.2, 222-222 (1 pages)
발행처 (Publisher)	<a href="#">대한종양외과학회</a> Korea society of Surgical Oncology
URL	<a href="http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE09326328">http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE09326328</a>
APA Style	Hee-Beom Yang, Hyun-Young Kim, Sung Eun Jung (2019). Thirteen Years' Experience of Pediatric Abdominal Mass and Analysis of Associated Factors Affecting Repeated Operation. <a href="#">대한임상종양학회 학술대회지</a> , 222-222.
이용정보 (Accessed)	이화여자대학교 203.255.***.68 2020/05/18 04:02 (KST)

---

### 저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되는 저작물은 DBpia와 구독계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

### Copyright Information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.

OP3-5

## Thirteen Years' Experience of Pediatric Abdominal Mass and Analysis of Associated Factors Affecting Repeated Operation

Hee-Beom Yang<sup>1</sup>, Hyun-Young Kim<sup>2</sup>, Sung Eun Jung<sup>2</sup>

<sup>1</sup>Department of Pediatric Surgery, Seoul National University Children's Hospital, Korea,

<sup>2</sup>Department of Pediatric Surgery, Seoul National University College of Medicine, Korea

### Background/Aims

Malignancies in pediatric population were quite different from those in adults. Advanced disease requiring neoadjuvant chemotherapy is frequent in pediatrics. This study was aimed to investigate the clinical characteristics of abdominal tumor and to analyze the associated factors affecting repeated operation for abdominal malignancies in pediatric population.

### Methods

Retrospective chart review was performed from January 2005 to December 2017 for patients who underwent surgery for abdominal tumor in Seoul National University Children's Hospital. Patients were screened through operation name and diagnosis, and selected with pathologic results and operation note. Patients' characteristics, tumor location, tumor size, method of operation, intent of operation, total number of operation, pathologic result, secondary tumor, and follow-up data were investigated. P-values < 0.05 were considered statistically significant. This study was approved by the institutional review board.

### Results

All operations, 16,719 operations during study period were included. Total 15,648 operations irrelevant to abdominal tumor were excluded. After detailed review of pathologic results and operation note, 792 operations in 662 patients were identified for abdominal tumor. Median age was 4 years old. The number of male was 252 (38.1%). The number of malignancy was 340 (51.4%). Operation for primary biopsy accounted for 63 (9.5%). The number of open surgery was 443 (67.0%). The number of patients who underwent surgery at least 2 times was 119 (18.0%). Mature teratoma and ovarian cyst were common benign tumor. Neuroblastoma and hepatoblastoma were common malignant tumor. Total 59 diseases for benign tumor, and 38 diseases for malignant tumor were identified. The number of patients who occurred second primary tumor irrelevant to primary malignancies was 23. Focal nodular hyperplasia in liver and renal cell carcinoma were common secondary tumor. Patients who underwent repeated operations were more aged (5 year vs 3.2 year,  $p=0.007$ ). Tumor location was significantly different, more in ovary and pancreas for non-repeated group and more in retroperitoneum in repeated group ( $p=0.014$ ). Diagnosis was not different significantly ( $p=0.183$ ). Secondary tumors were more in repeated group than in non-repeated group (15.6% vs 3.3%,  $p<0.001$ ).

### Conclusions

Various tumors were occurred in pediatric population. Common tumor were mature teratoma and neuroblastoma. Pediatric tumor is common in female due to high incidence of ovarian tumor. Repeated operation for abdominal malignancies in pediatric population was associated with tumor location, not diagnosis, and development of secondary tumor. Long term surveillance for detecting secondary tumor is needed in patients who underwent repeated operation for abdominal malignancies.