

Expression of Flavonoid and Stilbene Synthesis Genes in Grape Berries is Affected by High Temperature

저자 (Authors)	Seon Ae Kim, Sang-Keun Oh, Soon Young Ahn, Hae Keun Yun
출처 (Source)	한국원예학회 학술발표요지 , 2019.5, 33-33(1 pages) HORTICULTURE ABSTRACTS , 2019.5, 33-33(1 pages)
발행처 (Publisher)	한국원예학회 Korean Society For Horticultural Science
URL	http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE08739996
APA Style	Seon Ae Kim, Sang-Keun Oh, Soon Young Ahn, Hae Keun Yun (2019). Expression of Flavonoid and Stilbene Synthesis Genes in Grape Berries is Affected by High Temperature. 한국원예학회 학술발표요지 , 33-33
이용정보 (Accessed)	이화여자대학교 203.255.***.68 2020/05/18 04:00 (KST)

저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되는 저작물은 DBpia와 구독계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

Copyright Information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.

Different Vegetative Growth Stages of Kimchi cabbage (*Brassica rapa* L.) Exhibit Specific Glucosinolate Composition and Content

Byeong Wook Jeon¹, Man-Ho Oh², Eun Ok Kim³, Hyoung Seok Kim³, and Won Byoung Chae^{4*}

¹Vegetable Research Division, National Institute of Horticultural and Herbal Science, Rural Development Administration, Wanju 55365, Korea, ²Department of Biological Sciences, College of Biological Sciences and Biotechnology, Chungnam National University, Daejeon 34134, Korea, ³Natural Products Research Center, Korea Institute of Science and Technology (KIST), Gangneung 25451, Korea, ⁴Korea Program on International Agriculture, Rural Development Administration, Jeonju 54875, Korea

This study aimed to simultaneously investigate the changes in growth characteristics and glucosinolate (GL) content during different growth stages in Kimchi cabbage. Two Kimchi cabbage cultivars ‘Chuweol’ and ‘HwiparamGold’ were grown in the field and plant characteristics such as leaf length, number and fresh weight, and GL contents were measured weekly from 2 to 9 weeks after transplanting (WAT). The only significant difference between the two cultivars for either plant growth or GL content was observed for GL contents during 3 and 7 WAT. Leaf length increased until 4 WAT and then remained unchanged, exhibiting logarithmic growth. The fresh weight and number of leaves increased linearly until 9 WAT. Five GLs (two aliphatic GLs: progoitrin and gluconapin, two indole GSLs: glucobrassicin and neoglucobrassicin, and one aromatic GL: gluconasturtiin) of the nine GLs investigated in this study (glucoiberin, progoitrin, glucoraphanin, sinigrin, gluconapin, glucobrassicin, gluconasturtiin, 4-methoxy glucobrassicin and neoglucobrassicin) were detected in the two cultivars. The contents of these five GLs were similar in the two cultivars during 2 WAT, but gluconapin and gluconasturtiin increased more dramatically than the others. The increasing pattern of total GL more closely resembled the leaf growth pattern than the fresh weight. Our results suggest that the change in total GL content positively correlates to leaf length, and the increase in total GL content is attributed to the increase in the amount of gluconasturtiin and gluconapin during the autumn growing season.

T. 063-238-6614, chaeddang@korea.kr



성명: 채원병
학사: 2001 경희대학교 원예학과
석사: 2003 경희대학교 원예학과
박사: 2012 미국 University of Illinois
Urbana-Champaign, Dept. of Crop Sciences
2006-2016 농촌진흥청 국립원예특작과학원, 농업연구사
2017-2019 농촌진흥청 국외농업기술과, 농업연구사
2019-현재 농촌진흥청 국립원예특작과학원, 농업연구사

Expression of Flavonoid and Stilbene Synthesis Genes in Grape Berries is Affected by High Temperature

Seon Ae Kim^{1,2†}, Sang-Keun Oh^{3†}, Soon Young Ahn¹, and Hae Keun Yun^{1*}

¹Department of Horticulture and Life Science, Yeungnam University, Gyeongsan 38541, Korea, ²Apple Research Institute, National Institute of Horticultural and Herbal Science, RDA, Gunwi 39000, Korea, ³Department of Applied Biology, Chungnam National University, Daejeon 34134, Korea

Increased temperatures caused by climate change inhibit berry skin coloration during the ripening season of grape cultivation. In this study, we analyzed the expression patterns of flavonoid synthesis genes involved in color and stilbene compound synthesis genes at the transcript level in ‘Campbell Early’ grapes subjected to different temperatures. The expression of *chalcone isomerase (CHI)*, *leucoanthocyanidin dioxygenase (LDOX)*, and *trans-cinnamate 4-monooxygenase (CYP73A)* genes increased in all temperature conditions; however, their expression was lower at 30–35°C than 25°C, and was much lower at 35°C. Indeed, the expression of most tested genes was lower at 35°C than other temperatures. Additionally, subjecting grapes to high temperatures in the initial stage of veraison significantly reduced the synthesis and accumulation of anthocyanins. Moreover, the expression of the *resveratrol synthase (STS1)* gene was gradually induced at all temperature treatments, but decreased at 48 h and the expression at 25°C was inhibited when compared to expression at 35°C. Expression of *stilbe synthase1a-1 (STS11)*, *STS1a-2 (STS12)*, and *STS1a-3 (STS13)* genes had similar expression patterns at all temperatures, which decreased with increasing temperature and was suppressed at 35°C compared to 25°C. These results can be used to understand the response mechanisms of grapes to high temperature stress at the molecular level, as well as to provide information for grape breeding programs and viticulture to overcome obstacles caused by high temperatures associated with global climate change.

T. 053-810-2942, haekeun@ynu.ac.kr



성명: 윤해근
학사: 1989 충남대학교 응용생물학과
석사: 1991 충남대학교 대학원 응용생물학과
박사: 1996 충남대학교 대학원 응용생물학과
1994-1996 한국생명공학연구원 연구원
1997-1999 국립원예특작과학원 박사후연구원
1999-2009 농촌진흥청, 국립원예특작과학원 농업연구사
2002-2003 미국 플로리다주립대학교 Viticulture
Research Center 박사후연구원
2009-현재 영남대학교 원예생명과학과 교수