

Airway-gut Axis: Persistent Allergic Responses in Mouse Airway Affects Intestinal Microbial **Ecosystems**

Chan Min Jung, Sang Sun Yoon 저자

(Authors)

한국미생물학회 학술대회논문집 , 2019.4, 120-120(1 pages) 출처

(Source)

한국미생물학회 발행처

The Microbiological Society of Korea (Publisher)

http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE08756797 **URL**

Chan Min Jung, Sang Sun Yoon (2019). Airway-gut Axis: Persistent Allergic Responses in Mouse Airway Affects Intestinal Microbial Ecosystems. 한국미생물학회 학술대회논문집, 120-120 APA Style

이용정보 이화여자대학교 203.255.***.68 2020/05/18 04:01 (KST) (Accessed)

저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되 는 저작물은 DBpia와 구독계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에 서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

Copyright Information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.



D035

Drosophila melanogaster-based Identification of the Pseudomonas aeruginosa Genes for Polymicrobial Interaction with Staphylococcus aureus

Yeon-Ji Yoo, Shin-Yae Choi, Hye-Jeong Jang, In-Young Chung, and You-Hee Cho* Department of Pharmacy, College of Pharmacy and Institute of Pharmaceutical Sciences, CHA University

Microorganisms are widespread in natural habitats and human tissues where they coexist and constitute polymicrobial communities. It remains unclear how such polymicrobial interactions are orchestrated to survive the dynamic host environments. Here, we established a *Drosophila melanogaster*-based infection model to assess the key aspects of polymicrobial interaction between the two important opportunistic pathogens, Pseudomonas aeruginosa and Staphylococcus aureus that commonly coexist in the mucosal layers. This system revealed that the polymicrobial infection enhanced the virulence of P. aeruginosa, not that of S. aureus. This virulence enhancement was still evident in the interaction between the P. aeruginosa lasRmvfR mutant and a virulence-attenuated S. aureus mutant (m6), suggesting that the virulence enhancement did not require the LasR-MvfR quorum-sensing circuitry. To identify the genes required for the virulence enhancement in P. aeruginosa by S. aureus, we have screened ~1,000 transposon clones of the lasRmvfR mutant and isolated three mutants (11E3, 11G10, and 16E12), whose virulence was not enhanced at all by m6 in the Drosophila polymicrobial infection. Their transposon insertion sites were determined: a hypothetical gene (PA14_24300) under the control of MvfR, dgt for the dGTPase, and fleQ for the regulator of flagellar synthesis. These results suggest unexpected interspecies interactions during the polymicrobial infection caused by P. aeruginosa and S. aureus.

D036

Airway-gut Axis: Persistent Allergic Responses in Mouse Airway Affects Intestinal Microbial Ecosystems Chan Min Jung and Sang Sun Yoon*

Department of Microbiology and Immunology, Yonsei University College of Medicine

Asthma and allergic airway diseases are common in Westernized country and rapidly increasing their patient population in developing country. A hallmark of allergic airway disease is airway hyperresponsiveness, which is characterized by eosinophilic inflammation and airway remodeling. The host-microbiome interactions have been considered to modulate host immune mechanisms, especially in the mucosal surfaces. Microbiome analysis by next-generation sequencing allows us to gain more information between asthma and commensal microbes. Airway, once believed to be the sterile site, have an estimated number of 10-100 bacteria per 1,000 human cells. However there is little known about the correlation between allergic airway diseases and mucosal microbiome. Herein, we asked whether persistently applied allergic stimulations to the respiratory system would affect intestinal microbial ecosystems. In mouse model, HDM was treated intranasally for 8 consecutive weeks and samples were collected from cecum, feces, lung and airway secretion of each mice. We focused on microbiome change from chronic asthma mouse models induced by house dust mite(HDM), using 16S rRNA gene sequencing. There was an increase of Proteobacteria ratio in airway secretion, however, no significant change was seen in fecal samples. Addressing this question would help us provide clues to understand how an airway-gut axis operates.

[Supported by a grant from the National Research Foundation (NRF-2017M3A9F3041233).]