

## NOX4 affects the stage of neural differentiation in the hippocampus through independent with an inflammatory contribution

---

저자 (Authors)	Payanart Jiranugrom, Sunmi Yoon, Sun Shin Yi
출처 (Source)	<a href="#">한국실험동물학회 학술발표대회 논문집</a> , 2020.2, 130-130 (1 pages)
발행처 (Publisher)	<a href="#">한국실험동물학회</a> Korean Association for Laboratory Animal Science
URL	<a href="http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE09319095">http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE09319095</a>
APA Style	Payanart Jiranugrom, Sunmi Yoon, Sun Shin Yi (2020). NOX4 affects the stage of neural differentiation in the hippocampus through independent with an inflammatory contribution. <a href="#">한국실험동물학회 학술발표대회 논문집</a> , 130-130.
이용정보 (Accessed)	이화여자대학교 203.255.***.68 2020/05/18 03:54 (KST)

---

### 저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되는 저작물은 DBpia와 구독 계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

### Copyright Information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.

## PS-D-09

## NOX4 affects the stage of neural differentiation in the hippocampus through independent with an inflammatory contribution

Payanart Jiranugrom<sup>1,2</sup>, Sunmi Yoon<sup>1</sup>, Sun Shin Yi<sup>1\*</sup><sup>1</sup>Department of Biomedical Laboratory Science, Soonchunhyang University, Asan, Republic of Korea<sup>2</sup>Department of chemical engineering, King Mongkut's University of Technology Thonburi, Bangkok, Thailand

It has been known that reactive oxygen species (ROS) regulate diverse complex cellular processes including angiogenesis, inflammation, differentiation, and proliferation. The family of NADPH oxidases (Nox) consists of 7 iso-forms with tissue- and cell type-specific expression profiles and Nox4 is constitutively active and predominantly produces H<sub>2</sub>O<sub>2</sub> among these. It is known that the difference in the amount of hippocampal neurogenesis in the brain is compassionate, especially the continuous supply of high-fat diet (HF) has a very negative effect on hippocampal neurogenesis. Therefore, if NOX4 is specifically removed, it is expected that the expression of ROS to the brain would be limited, which could result in an increase of neurogenesis in NOX4 KO mice. To this end, Normal chow (NC) and HF were supplied to WT and NOX4 KO mice for seven weeks from 7 weeks, and their body weight and food intake were tracked. The volume of white adipose tissue (WAT) in animals, cell proliferation (Ki67) in walrus, hippocampal neurogenesis (DCX) and distribution of microglia (microglia) were observed by immunohistochemistry (IHC) after sacrificing animals. The results showed that NOX4 KO animals had a significant difference in weight gain over seven weeks in comparison to WT animals in the HF supply group, but there was no difference in NC. Interestingly, while the number of Ki67 positive cells in the hippocampus in NOX4 KO has been confirmed to increase compared to WT animals in the NC and HF, DCX has been confirmed to be very low in NOX4 KO. Although NOX4 is a critical gene for inflammatory control mechanisms, it is believed that it affects the stage of neural differentiation in the hippocampus through independent channels in the neurogenesis of the hippocampus.

\*Corresponding author : Sun Shin Yi

Keywords : NOX4, Inflammation, Cell proliferation, Neurogenesis, Hippocampus

## PS-D-10

## A novel gene expression signature for gastric cancer stemness

Hyeokwon An, Siyou Kim, Jongwon Kwon, Junwon Park\*

Division of Biomedical Convergence, College of Biomedical Science, Kangwon National University, Chuncheon, 24341, Republic of Korea

Gastric cancer (GC) is the leading cause of cancer-associated death in Korea. In our previous study, we established GC stem cell-like murine cell line, namely S1M, lacking Smad4, p53, and E-cadherin. We identified 295 differentially upregulated genes in S1M cells from multi-omics analyses. Here, we tried to explore a novel gene expression signature associated with GC stemness by in-vivo transplantations of S1M cells in immunocompetent mice. From the 295 gene expression profile, we selected 30 candidate genes based on their clinical relevance for GC patients. S1M cells were transduced with cas9 and lentiviral gRNAs for these 30 genes and transplanted into subcutaneous tissues and spleens of immunocompetent mice. Genomic DNA was collected from the subcutaneous tumors or metastatic lesions in livers and the integrated gRNA were retrieved by PCR. The proportional changes of each gRNA in the retrieved gRNAs were evaluated by quantitative real-time PCR analysis. As a result, gRNAs for several genes including Spp1, Il1rl1 and Tmsb4x showed statistically significant reductions of proportions in both subcutaneous tumors and liver metastases, implying that these genes might contribute to GC tumorigenesis and metastasis. Especially, knockout of Tmsb4x in S1M cells showed a greater suppressive effect on the drug resistance and stem cell sphere forming ability in-vitro. Collectively, we discover the novel gene expression signature for GC stemness.

\*Corresponding author : Junwon Park

Keywords : Immunocompetent, Gastric cancer, Metastasis

## PS-D-11

## Effects of e2f3 gene deletion in mice

Ki-Hoan Nam\*, Hae-Rim Kim, Eun-Kyeong Kim, San-Mi Cho, Ki-Hoon Lee, Young-Won Seo, Young-Suk Won, Hoyoung Kang, won-Kee Yoon, Hyoung-Chin Kim

한국생명공학연구원 실험동물자원센터

E2F transcription factors (E2Fs) are found in most cell types of the body and contribute to cell cycle progression, cell proliferation, differentiation, and apoptosis processes. These E2F transcription factors are potent regulators of a variety of genes, including cell-cycle checkpoints controlling genes in mammalian cells. Until now, eight members have been characterized, E2F1 to 8, and they are generally classified into transcriptional activators (E2F1-3) and repressors (E2F4-8). Some of these E2F family members have also a key role, in myeloid development and cardiac neovascularization. For transcriptional regulation, dimerization partner (DP) family and retinoblastoma (RB) family collaborate with E2Fs, by binding to E2F proteins. DP proteins identified DP1-3 form a heterodimer with an E2F, and allow it to bind to target promoters. A member of the E2F family, E2f3 plays a critical role in cell cycle and proliferation by targeting downstream, retinoblastoma (RB) a tumor suppressor family protein. The aim of this study, was to investigate the effects of E2F3 deficiency in vivo. We examined phenotypic abnormalities, by mutation of the E2f3 gene in mice. Complete deletion of the E2f3 was fully penetrant, in the pure C57BL/6N background. The E2f3 heterozygous mouse embryo developed normally without fatal abnormality. However, they exhibited altered phenotypes in body weight, growth rate, skeletal perfection, and grip strength ability. Findings suggest that E2F3 has a critical role in muscle and bone development, and affect normal mouse growth.

\*Corresponding author : Ki-Hoan Nam

Keywords : E2f3, Gene deletion, Phenotype

## PS-D-12

## The effect of Importin-11 is in the normal embryonic development

Eun-Kyeong Kim<sup>1</sup>, Ju-Young Lee<sup>1</sup>, Sang-Mi Cho<sup>1</sup>, Hae-Rim Kim<sup>1</sup>, Won-Kee Yoon<sup>1</sup>, Hyoung-Chin Kim<sup>1</sup>, Bae-Hwan Kim<sup>2\*</sup>, Ki-Hoan Nam<sup>1\*</sup><sup>1</sup>Laboratory Animal Resource Center, KRIBB<sup>2</sup>Department of Public Health, College of Natural Science, Keimyung University

Importin-11 (Ipo11) is a new member of the human importin family of transport receptors which are well-known to mediate the nucleocytoplasmic transport of protein and RNA cargos. Despite its role in the transport of some proteins, we found that deletion of importin-11 affects normal embryonic development and govern embryo-lethality in mice. In this study, we for the first time produced a mouse line containing complete null mutation in Ipo11 gene utilized by gene trapping method. The Ipo11 deficient mouse embryos showed an embryonic lethal phenotype. The Ipo11 deficient embryos showed a reduced size at embryonic day 10.5 (E10.5) when compared with Ipo11 wild type or Ipo11 heterozygous embryos and died by E11.5. Whereas Ipo11 heterozygous mice were healthy and fertile, and there was no detectable abnormality in their appearance when reviewed. In the X-gal staining with the Ipo11 homozygous or Ipo11 heterozygous embryos, strong X-gal staining positivity was detected systematically in the whole mount embryos at E10.5, although almost no X-gal positive signal was detected at E9.5, indicating that the embryos die soon after the stage of Ipo11 expression in the embryos. These results indicate that Ipo11 is critical for the normal embryonic development in mice indicating its novel role.

\*Corresponding author : Bae-Hwan Kim, Ki-Hoan Nam

Keywords : Ipo11, Gene deletion, Mouse